

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 40 (1965-1966)

Artikel: A 3-Fold Vector Product in R8.
Autor: Zvengrowski, Peter
DOI: <https://doi.org/10.5169/seals-30632>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A 3-Fold Vector Product in R^8

By PETER ZVENGROWSKI

An r -fold vector product in R^n is, according to ECKMANN [2], a continuous map

$$X : R^{rn} = R^n \times R^n \times \dots \times R^n \rightarrow R^n$$

- satisfying (i) $X(a_1, a_2, \dots, a_r) \cdot a_i = 0$, $1 \leq i \leq n$,
(ii) $\|X(a_1, a_2, \dots, a_r)\|^2 = |a_i \cdot a_j|$.

It can be deduced from [2] and [4] that R^n admits an r -fold vector product in precisely the following cases:

- a) $r = 1$ and n is even,
- b) $r = n - 1$,
- c) $r = 2$ and $n = 3, 7$,
- d) $r = 3$ and $n = 8$.

Explicit formulas for these vector products are known in all but the last case, and the purpose of this note is to give a simple formula for the last case, taking advantage of the CAYLEY numbers. Such a formula was useful to the author in examining homotopy classes of vector fields on S^7 (cf. [5]).

1. The Cayley Numbers

We take the CAYLEY numbers C to be the 8-dimensional algebra over the reals having basis vectors e_1, e_2, \dots, e_8 and suitably defined products. In particular, $e_1 = 1$ is the unit and for $i, j > 1$ one has $e_i e_j = \begin{cases} -1, & i = j \\ -e_j e_i, & i \neq j. \end{cases}$

The subspace of C orthogonal to e_1 is called the pure CAYLEY numbers. Conjugation is defined by writing any CAYLEY number x as $x = r e_1 + x'$, where x' is pure, and setting $\bar{x} = r e_1 - x'$. The CAYLEY numbers inherit the norm and scalar product of R^8 , and

$$x\bar{x} = \bar{x}x = x \cdot x = |x|^2.$$

Lemma 1.1. For any CAYLEY numbers a, b , one has $\bar{\bar{a}} = a$, $\bar{ab} = \bar{b}\bar{a}$, and $a\bar{b} + b\bar{a} = 2(a \cdot b)$.

Proof: By linearity it suffices to prove these when a, b are the units e_1, \dots, e_8 , and these cases follow from the above remarks on multiplication.

Corollary: $\bar{a}b + \bar{b}a = 2(\bar{a} \cdot \bar{b}) = 2(a \cdot b)$.

The multiplication in C can be explicitly defined by regarding C as a 2-dimensional algebra over the quaternions H , with basis $1, k$, and setting

$$(q_1 + q_2k)(q'_1 + q'_2k) = (q_1q'_1 - \bar{q}'_2q_2) + (q'_2q_1 + q_2\bar{q}'_1)k.$$

We shall need this formula to prove the next lemma.

Lemma 1.2. For any CAYLEY numbers a, b, c ,

$$a(\bar{b}c) + b(\bar{a}c) = 2(a \cdot b)c.$$

Proof: Let $a = a_1 + a_2k$, $b = b_1 + b_2k$, $c = c_1 + c_2k$, where $a_1, a_2, b_1, b_2, c_1, c_2$ are quaternions. By direct computation,

$$\begin{aligned} a(\bar{b}c) + b(\bar{a}c) &= (b_1\bar{a}_1 + a_1\bar{b}_1)c_1 + c_1(\bar{a}_2b_2 + \bar{b}_2a_2) + \\ &\quad + [c_2(\bar{a}_1b_1 + \bar{b}_1a_1) + (b_2\bar{a}_2 + a_2\bar{b}_2)c_2]k. \end{aligned}$$

Applying Lemma 1.1 and its corollary, this equals

$$\begin{aligned} &2[(b_1 \cdot a_1)c_1 + c_1(a_2 \cdot b_2) + (c_2(a_1 \cdot b_1) + (b_2 \cdot a_2)c_2)k] \\ &= 2(a_1 \cdot b_1 + a_2 \cdot b_2)(c_1 + c_2k) \\ &= 2(a \cdot b)c. \end{aligned}$$

According to [1], the subalgebra of C generated by any two elements (and their conjugates) is associative. In particular, for any CAYLEY numbers a, b, c , the product $a(bc)a$ is well defined, and according to [3], we have $a(bc)a = (ab)(ac)$.

Lemma 1.3. For any CAYLEY numbers a, b, c , one has

- (i) $a \cdot (a(\bar{b}c)) = |a|^2(b \cdot c)$,
- (ii) $c \cdot (a(\bar{b}c)) = |c|^2(a \cdot b)$, and
- (iii) $b \cdot (a(\bar{b}c)) = -|b|^2(c \cdot a) + 2(a \cdot b)(b \cdot c)$.

Proof of (i): $2a \cdot (a(\bar{b}c)) = a(\overline{a(\bar{b}c)}) + (a(\bar{b}c))\bar{a} = a((\bar{c}b)\bar{a}) + (a(\bar{b}c))\bar{a}$. Since a , \bar{a} , $\bar{c}b$, and $\bar{c}\bar{b} = \bar{b}c$ generate an associative subalgebra, this equals

$$a(\bar{c}b)\bar{a} + a(\bar{b}c)\bar{a} = a(\bar{c}b + \bar{b}c)\bar{a} = 2a(b \cdot c)\bar{a} = 2|a|^2(b \cdot c).$$

Proof of (ii): $2c \cdot (a(\bar{b}c)) = c((\bar{c}b)\bar{a}) + (a(\bar{b}c))\bar{c} = 2r$, say, where r must be a real number. Then making use of the preceding results without specific mention, we have

$$\begin{aligned} rc &= c((\bar{c}b)\bar{a})c + (a(\bar{b}c))|c|^2 \\ &= (c(\bar{c}b))(\bar{a}c) + (a(\bar{b}c))|c|^2 \\ &= |c|^2(b(\bar{a}c) + a(\bar{b}c)) \\ &= 2|c|^2(a \cdot b)c. \end{aligned}$$

Proof of (iii): $2b \cdot (a(\bar{b}c)) = b((\bar{c}b)\bar{a}) + (a(\bar{b}c))\bar{b}$

$$\begin{aligned} &= b[(-\bar{b}c + 2b \cdot c)\bar{a}] + [a(-\bar{c}b + 2b \cdot c)]\bar{b} \\ &= 2(b \cdot c)(b\bar{a} + a\bar{b}) - b((\bar{b}c)\bar{a}) - (a(\bar{c}b))\bar{b} \\ &= 4(b \cdot c)(a \cdot b) - 2r, \text{ for some real number } r. \end{aligned}$$

Computing rb as in (ii), one finds $r = |b|^2(c \cdot a)$.

2. Applications to Vector Products

A 2-fold vector product in R^7 is obtained (cf. [2]) by identifying R^7 with the pure CAYLEY numbers, and for any two vectors b, c in R^7 putting $X_2(b, c) = bc + b \cdot c$. We shall now define a 3-fold vector product in R^8 , which in a certain sense generalizes X_2 .

Theorem 2.1. Let a, b, c be vectors in R^8 , regarded as the CAYLEY numbers. Then

$$X_3(a, b, c) = -a(\bar{b}c) + a(b \cdot c) - b(c \cdot a) + c(a \cdot b)$$

defines a 3-fold vector product in R^8 .

Proof: Lemma 1.3 will be frequently used without specific mention, and we write $X = X_3(a, b, c)$. The continuity of X_3 is obvious. Thus $a \cdot X = -|a|^2(b \cdot c) + |a|^2(b \cdot c) - (a \cdot b)(c \cdot a) + (a \cdot c)(a \cdot b) = 0$, and the proofs that $b \cdot X = 0$, $c \cdot X = 0$, are entirely similar. Finally, writing $d = (a(\bar{b}c))$, we have

$$\begin{aligned}
|X|^2 &= X \cdot X = |d|^2 + |a|^2(b \cdot c)^2 + |b|^2(a \cdot c)^2 + |c|^2(a \cdot b)^2 - 2(a \cdot d)(b \cdot c) \\
&\quad + 2(b \cdot d)(c \cdot a) - 2(c \cdot d)(a \cdot b) + (-2 + 2 - 2)(a \cdot b)(b \cdot c)(c \cdot a) \\
&= |a|^2|b|^2|c|^2 + |a|^2(b \cdot c)^2 + |b|^2(a \cdot c)^2 + |c|^2(a \cdot b)^2 \\
&\quad - 2|a|^2(b \cdot c)^2 - 2|b|^2(a \cdot c)^2 + 4(a \cdot b)(b \cdot c)(c \cdot a) - 2|c|^2(a \cdot b)^2 \\
&\quad - 2(a \cdot b)(b \cdot c)(c \cdot a) \\
&= |a|^2|b|^2|c|^2 - |a|^2(b \cdot c)^2 - |b|^2(a \cdot c)^2 - |c|^2(a \cdot b)^2 + 2(a \cdot b)(b \cdot c)(c \cdot a) \\
&= \begin{vmatrix} |a|^2 & a \cdot b & a \cdot c \\ b \cdot a & |b|^2 & b \cdot c \\ c \cdot a & c \cdot b & |c|^2 \end{vmatrix}.
\end{aligned}$$

When restricted to the STIEFEL manifold $V_{7,2}$, the 2-fold product X_2 gives a cross section $X_2: V_{7,2} \rightarrow V_{7,3}$. Similarly, X_3 gives a cross section $X_3: V_{8,3} \rightarrow V_{8,4}$. Then the fact that $X_3(e_1, b, c) = bc + b \cdot c = X_2(b, c)$, where $(e_1, b, c) \in V_{8,3}$ (and hence b, c are pure), implies that the following diagram commutes:

$$\begin{array}{ccc}
V_{7,2} & \xrightarrow{i} & V_{8,3} \\
\downarrow X_2 & & \downarrow X_3 \\
V_{7,3} & \xrightarrow{i'} & V_{8,4}.
\end{array}$$

Further details on these vector products can be found in [5].

The University of Illinois

REFERENCES

- [1] DICKSON L. E.: *Linear Algebras*, Cambridge Univ. Press (1914).
- [2] ECKMANN B.: *Stetige Lösungen linearer Gleichungssysteme*, Comment. Math. Helv. 15, 358–366 (1942).
- [3] TODA H., SAITO Y. and YOKOTA I.: *Note on the Generator of $\pi_7(SO(n))$* , Kyoto Univ. Memoirs Vol. XXX Math., No. 3, 227–230 (1957).
- [4] WHITEHEAD G. W.: *Note on Cross Sections of Stiefel Manifolds*, Comment. Math. Helv. 37, 239–240 (1963).
- [5] ZVENGROWSKI P.: *Vector Fields and Vector Products*, Chapter 4: Vector Products, Thesis, Univ. of Chicago (1965).

(Received April 5, 1965)