

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 44 (1969)

**Artikel:** On a Special Class of Hamiltonian Graphs.  
**Autor:** Chartrand, Gary / Kronk, Hudson V.  
**DOI:** <https://doi.org/10.5169/seals-33757>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 15.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## On a Special Class of Hamiltonian Graphs

GARY CHARTRAND<sup>1)</sup> and HUDSON V. KRONK

One of the most basic questions asked about a graph (finite, undirected, without loops or multiple edges) is whether its structure is such that it can be traversed or traced in a certain manner. Undoubtedly, the two most important classes of graphs dealing with traversability are the eulerian graphs and the hamiltonian graphs. A graph  $G$  is *eulerian* if it has a closed path (called an eulerian path) containing every edge of  $G$  exactly once and every vertex of  $G$  at least once, while  $G$  is *hamiltonian* if it has a closed path containing every vertex of  $G$  exactly once, i.e. if it has a hamiltonian cycle.

A graph  $G$  is said to be *randomly eulerian from a vertex  $v$*  if the following procedure always results in an eulerian path. Begin at the given vertex  $v$  and traverse any incident edge. On arriving at a vertex, choose any incident edge which has not yet been traversed. When no new edges are available the procedure terminates. These graphs have also been referred to as arbitrarily traversable from  $v$  and arbitrarily traceable from  $v$  and have been investigated by BÄBBLER [1], HARARY [3], and ORE [4].

This suggests the following concept. We define a graph  $G$  to be *randomly hamiltonian from the vertex  $v$*  if the following procedure always results in a hamiltonian cycle. Begin at the vertex  $v$  and proceed to any adjacent vertex. On arriving at a vertex, select any adjacent vertex not previously encountered. When no new vertices remain, then an edge exists between the final vertex chosen and  $v$ , and the procedure terminates. Thus in a graph  $G$  which is randomly hamiltonian from a vertex  $v$ , any path beginning at  $v$  can be extended to a hamiltonian cycle. Graphs which are randomly hamiltonian from every vertex were characterized in [2] and are called simply randomly hamiltonian graphs.

It is the object of this article to present a characterization of graphs which are randomly hamiltonian from a vertex, and thereby provide a classification of all such graphs.

It is convenient to introduce notation for several types of graphs which are encountered throughout the course of this article. The complete graph with  $p$  vertices is denoted by  $K_p$ , while  $C_p$  represents the cycle with  $p \geq 3$  vertices. The complete bipartite graph  $K(m, n)$  is the graph with  $p = m + n$  vertices whose vertex set  $V$  can be partitioned as  $V_1 \cup V_2$  such that  $|V_1| = m$ ,  $|V_2| = n$ , and vertices  $u$  and  $v$  are adjacent if and only if  $u \in V_i$  and  $v \in V_j$ ,  $i \neq j$ . It was shown in [2] that a graph  $G$  with  $p \geq 3$  vertices is randomly hamiltonian if and only if it is one of the graphs  $K_p$ ,  $C_p$ , and  $K(p/2, p/2)$ .

<sup>1)</sup> Research supported in part by a Faculty Research Fellowship from Western Michigan University.

We express a graph  $G$  as  $H+v$  provided  $v$  is a vertex of  $G$  adjacent to all other vertices of  $G$ , where then  $H$  is the graph obtained from  $G$  by the removal of  $v$  and all edges incident with  $v$ . For example, the graph  $C_n+v$  is often referred to as the wheel  $W_n$ . The graphs  $K(3,3)+v$  and  $W_5=C_5+v$  are illustrated in Figure 1. In each case, the graph is randomly hamiltonian from  $v$ .



Figure 1  
Two graphs which are randomly hamiltonian from the vertex  $v$

Of course, if a graph  $G$  is randomly hamiltonian from a vertex, then  $G$  is hamiltonian and therefore has a hamiltonian cycle. Thus whenever we have a graph  $G$  with  $p$  vertices which is randomly hamiltonian from a vertex we assume the existence of a hamiltonian cycle  $C$  whose vertices are labeled cyclically  $v_1, v_2, \dots, v_p$ . Each edge of  $G$  then either belongs to  $C$ , and is called a *cycle edge* of  $G$ , or joins two non-consecutive vertices of  $C$  and is called a *diagonal*.

If  $G$  is a graph which is randomly hamiltonian from some vertex (and which contains a hamiltonian cycle  $C$  labeled as earlier indicated), then any cycle of  $G$  containing exactly one diagonal of  $G$  is called an *outer cycle* of  $G$ . An *outer  $n$ -cycle* has length  $n$ , and an outer 3-cycle is also referred to as an *outer triangle*.

We now present the main result of the paper.

**THEOREM.** *A graph  $G$  is randomly hamiltonian from a vertex  $v$  if and only if  $G$  is randomly hamiltonian or  $G=H+v$ , where  $H$  is randomly hamiltonian.*

*Proof.* If  $G$  is a randomly hamiltonian graph containing a vertex  $v$  or if  $G$  is expressible as  $H+v$ , where  $H$  is randomly hamiltonian, then it is easily observed that  $G$  is randomly hamiltonian from  $v$ .

Conversely, let  $G$  be a graph with  $p$  vertices which is randomly hamiltonian from the vertex  $v$ . Thus  $G$  contains a hamiltonian cycle  $V$  whose vertices we label cyclically as  $v=v_1, v_2, \dots, v_p$ .

Suppose that  $G$  is not randomly hamiltonian so that  $G$  is none of the graphs  $K_p, C_p, K(p/2, p/2)$ . In particular, this implies that  $G$  contains diagonals so that  $G$  necessarily contains outer cycles. Hence the vertex  $v$  belongs to one or more outer cycles.

Let  $n$  be the length of the smallest outer cycle containing  $v$ . We first show that there exists an outer  $n$ -cycle containing  $v$  but in which  $v$  is not the endpoint of the associated diagonal. Suppose that the vertices of an outer  $n$ -cycle are  $v_1, v_2, \dots, v_n$ . Consider the path which commences at  $v_1$ , proceeds to  $v_n$  along the diagonal  $v_1v_n$ , and encounters in succession the vertices  $v_{n+1}, v_{n+2}, \dots, v_p$ . Since  $G$  is randomly hamiltonian from  $v=v_1$  and  $v$  belongs to no outer  $k$ -cycle,  $k < n$ , the diagonal  $v_pv_{n-1}$  must be present in  $G$ . Hence  $v$  belongs to the outer  $n$ -cycle whose vertices are  $v_p, v_1, \dots, v_{n-1}$ . In a similar way, one can show that if  $v_{p-n+2}v_1$  is a diagonal of  $G$ , then  $v_{p-n+3}v_2$  is a diagonal of  $G$ .

Thus we may assume the existence of an outer  $n$ -cycle whose vertices are  $v_m, v_{m+1}, \dots, v_p, v_1, \dots, v_{k-1}$ , where  $m=p-n+k$  and  $3 \leq k \leq n$ . We now show that the diagonals  $v_{m-1}v_{k-2}$  and  $v_{m+1}v_k$  are present in  $G$  in addition to  $v_mv_{k-1}$ . We begin a path at  $v=v_1$  and proceed along  $C$  to  $v_p, v_{p-1}, \dots, v_m$ . Following along the diagonal  $v_mv_{k-1}$  to  $v_{k-1}$  and then taking  $v_k, v_{k+1}, \dots, v_{m-1}$ , we see that  $v_{m-1}v_{k-2}$  is a diagonal of  $G$  since  $G$  is randomly hamiltonian from  $v$  and  $v$  belongs to no outer  $t$ -cycle,  $t < n$ . Similarly, by applying the preceding arguments to the path  $v_1, v_2, \dots, v_{k-1}, v_m, v_{m-1}, v_{m-2}, \dots, v_k$ , we observe that  $v_{m+1}v_k$  is a diagonal of  $G$ .

We now prove that  $n < 5$ , for suppose, to the contrary, that  $n \geq 5$ . We have already seen that there exists an outer  $n$ -cycle whose vertices are  $v_m, v_{m+1}, \dots, v_p, v_1, \dots, v_{k-1}$ , where  $m=p-n+k$  and  $3 \leq k \leq n$ , and, in addition, the edges  $v_{m-1}v_{k-2}$  and  $v_{m+1}v_k$  belong to  $G$ . Furthermore, since  $n \geq 5$ ,  $v_1$  is not adjacent to both  $v_m$  and  $v_{k-1}$ . Let us say that  $v_1$  is not adjacent to  $v_{k-1}$ , the other case being handled analogously. We now construct a path which begins at  $v=v_1$  and takes in succession  $v_p, v_{p-1}, \dots, v_{m+1}$ . We then proceed to  $v_k$  via the diagonal  $v_{m+1}v_k$  and move along  $C$  in the order  $v_{k+1}, v_{k+2}, \dots, v_{m-1}$ . On reaching  $v_{m-1}$ , we next take  $v_{k-2}$  (which is different from  $v$ ),  $v_{k-1}$ , and then  $v_m$ . Since  $G$  is randomly hamiltonian from  $v$ , there exists either a vertex not yet encountered which is adjacent to  $v_m$  or the edge  $v_mv$  which completes a hamiltonian cycle. In either case, there exists an edge  $v_mu$ , where  $u$  is one of the vertices  $v_{m+2}, v_{m+3}, \dots, v_p, v_1, \dots, v_{k-3}$ , which determines an outer cycle containing  $v$  having length less than  $n$ , and this is a contradiction.

We now show that  $n \neq 4$ . To prove this, we assume  $n=4$  so that  $v$  belongs to an outer 4-cycle but not an outer triangle. From what we have shown above, we may assume, without loss of generality, that  $v_p, v_1, v_2, v_3$  are the vertices of an outer 4-cycle. Since  $G$  is randomly hamiltonian from  $v=v_1$ , the path  $v_1, v_2, v_3, v_p, v_{p-1}, v_{p-2}, \dots, v_4$ , which contains all vertices of  $G$ , implies that  $v_1v_4$  is an edge of  $G$ . The path  $v_1, v_4, v_3, v_p, v_{p-1}, v_{p-2}, \dots, v_5$  contains all the vertices of  $G$  with the exception of  $v_2$ ; hence  $v_2v_5$  is an edge of  $G$ . Next the path  $v_1, v_2, v_5, v_4, v_3, v_p, v_{p-1}, v_{p-2}, \dots, v_6$  contains all the vertices of  $G$  and, as such, implies that  $v_1v_6$  is an edge of  $G$ . Continuing inductively, it is now easily verified that all edges of the type  $v_1v_{2m}$  belong to  $G$  as do all edges of the type  $v_2v_{2m+1}$ . From this it now follows that every two vertices  $v_a$

and  $v_\beta$ , where  $\alpha$  and  $\beta$  are of opposite parity, are adjacent. To see this, let  $v_{2r}$  and  $v_{2s+1}$  be two non-consecutive vertices of  $C$ , where  $v_{2r}$  is different from  $v_2$  and  $v_{2s+1}$  is not  $v_1$ . There are two cases to consider according to whether the path  $v_{2r}, v_{2r+1}, \dots, v_{2s+1}$  does or does not contain the vertex  $v$ . We treat here only the latter case, the former case being handled in a similar manner. We construct a path which begins at  $v_1$ , proceeds along a diagonal to  $v_{2s}$ , then along  $C$  to the vertices  $v_{2s-1}, v_{2s-2}, \dots, v_{2r+1}$ , from where we move to  $v_2$  by way of the diagonal  $v_2 v_{2r+1}$ . Next we proceed to  $v_{2r-1}$  via the diagonal  $v_2 v_{2r-1}$  and then take  $v_{2r-2}, v_{2r-3}, \dots, v_3, v_p, v_{p-1}, \dots, v_{2s+1}$ , which produces a path failing only to contain  $v_{2r}$ . Since  $G$  is randomly hamiltonian from  $v$ , the edge  $v_{2r} v_{2s+1}$  must be present in  $G$ . Finally, we show that if  $\alpha$  and  $\beta$  are of the same parity, then  $v_\alpha$  and  $v_\beta$  are not adjacent. We consider here only the case where  $\alpha$  and  $\beta$  are odd, the other case following similarly. Assume, to the contrary, that the vertices  $v_{2r+1}$  and  $v_{2s+1}$  are adjacent, where  $2r+1 < 2s+1$ , say. The path  $v_1, v_{2s+2}, v_{2s+3}, \dots, v_p, v_3, v_4, \dots, v_{2r+1}, v_{2s+1}, v_{2s}, \dots, v_{2r+2}$  fails only to contain the vertex  $v_2$ ; thus  $v_2 v_{2r+2}$  is an edge of  $G$ . From this we see that the path  $v_1, v_p, v_{p-1}, \dots, v_{2r+2}, v_2, v_{2r+1}, v_{2r}, \dots, v_3$ , which contains all vertices of  $G$ , implies that  $v_1 v_3$  is a diagonal of  $G$ . However, this contradicts the fact that  $v_1$  belongs to no outer triangle. Hence,  $v_\alpha$  and  $v_\beta$  are adjacent if and only if  $\alpha$  and  $\beta$  are of opposite parity. This implies that  $p$  is even since  $v_1 v_p$  is an edge of  $G$ . Furthermore, by letting  $V_1 = \{v_{2n} \mid n=1, 2, \dots, p/2\}$  and  $V_2 = \{v_{2n-1} \mid n=1, 2, \dots, p/2\}$ , we see that  $G$  is the graph  $K(p/2, p/2)$ , which, as noted earlier, is randomly hamiltonian. However, this is a contradiction since it is contrary to our assumption that  $G$  is not randomly hamiltonian.

We now arrive at the conclusion that the only possible value is  $n=3$ ; thus  $v$  belongs to an outer triangle. From methods similar to those we have already employed, it is immediately established that  $G$  contains the edges  $v_1 v_3, v_2 v_p$ , and  $v_1 v_{p-1}$ . Thus  $v_1$  is adjacent to each of the vertices  $v_2, v_3$ , and  $v_p$ . However,  $v_1$  is necessarily adjacent to all other vertices of  $G$ , for if  $v_1 v_k$  is an edge of  $G$ ,  $3 \leq k < p-1$ , then so too is  $v_1 v_{k+1}$  an edge of  $G$  since the path  $v_1, v_k, v_{k+1}, \dots, v_2, v_p, v_{p-1}, \dots, v_{k+1}$  contains all vertices of  $G$  and therefore  $v_1$  is adjacent to  $v_{k+1}$ . The result then follows by induction.

Hence we may express  $G$  as  $H+v$ . The only remaining detail now is to verify that  $H$  is randomly hamiltonian. In order to prove this, it is necessary to show that any path  $u_1, u_2, \dots, u_k$  of  $H$  can be extended to a hamiltonian cycle of  $H$ . Since  $v$  is adjacent to  $u_1$ :  $v, u_1, u_2, \dots, u_k$  is a path of  $G$  and can be extended to a hamiltonian cycle  $v, u_1, u_2, \dots, u_k, u_{k+1}, \dots, u_{p-1}, v$  of  $G$ . However,  $v$  is also adjacent to  $u_2$ ; thus  $v, u_2, u_3, \dots, u_{p-1}$  is also a path of  $G$  and can be extended to a hamiltonian cycle of  $G$ . This implies that  $u_{p-1} u_1$  is an edge of  $G$  so that  $u_1, u_2, \dots, u_{p-1}, u_1$  is a hamiltonian cycle of  $H$ . Hence  $H$  is randomly hamiltonian, completing the proof.

The preceding theorem now indicates that the only graphs with  $p$  vertices which are randomly hamiltonian from some vertex  $v$  are  $C_p, K_p, K(p/2, p/2), C_{p-1} + v$ , and  $K((p-1)/2, (p-1)/2) + v$ . As one final observation, we state the following.

COROLLARY. *The number of vertices in a graph  $G$  with  $p$  vertices from which  $G$  is randomly hamiltonian is either 0, 1, or  $p$ .*

## REFERENCES

- [1] F. BÄBLER, *Über eine spezielle Klasse Euler'scher Graphen*, Comment. Math. Helv. 27 (1953), 81–100.
- [2] G. CHARTRAND and H. V. KRONK, *Randomly Traceable Graphs*, J. Soc. Indust. Appl. Math. (to appear).
- [3] F. HARARY, *On Arbitrarily Traceable Graphs and Directed Graphs*, Scripta Math. 23 (1957), 37–41.
- [4] O. ORE, *A Problem Regarding the Tracing of Graphs*, Elem. Math. 6 (1951), 49–53.

*Western Michigan University  
SUNY at Binghamton*

Received January 24, 1968