Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 44 (1969)

Artikel: Note on Poisson's Treatment of the Euler-Maclaurin Formula.
Autor: Ostrowski, Alexander
DOI: https://doi.org/10.5169/seals-33765

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 07.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Note on Poissson's Treatment of the Euler–Maclaurin Formula[1])

(To Hugo Hadwiger for his 60-th birthday)

By ALEXANDER OSTROWSKI (Basel)

1. In the Euler-Maclaurin Formula

$$\sum_{\mu=p}^{q}{}' f(\mu) = \int_p^q f(t)\,dt + \sum_{\nu=1}^{n} \frac{B_{2\nu}}{(2\nu)!}\left(f^{(2\nu-1)}(q) - f^{(2\nu-1)}(p)\right) + R_n \qquad (n \geqq 1), \tag{1}$$

an expression for the remainder term was first given by POISSON [6] 1827. This is

$$R_n = \frac{1}{(2n+1)!}\int_p^q \overline{B_{2n+1}}(t) f^{(2n+1)}(t)\,dt, \tag{2}$$

where $\overline{B_{2n+1}}(t)$ is given by

$$B_{2n+1}(t) = \frac{2}{(2\pi)^{2n+1}} U_{2n+1}(t), \qquad U_s(t) = \sum_{\nu=1}^{\infty} \frac{\sin 2\nu\pi t}{\nu^s} \qquad (s > 1). \tag{3}$$

To obtain from here an estimate for $U_{2n+1}(t)$, POISSON used the obvious bound

$$|U_{2n+1}(t)| \leqq \sum_{\nu=1}^{\infty} \frac{1}{\nu^{2n+1}}. \tag{4}$$

2. It has apparently not as yet been noticed, that the inequality (4) can be replaced with

$$|U_{2n+1}(t)| < 1 \qquad (0 \leqq t \leqq 1, \quad n = 1, 2, \ldots).^{2)} \tag{5}$$

---

[1]) The work on this paper was partly done under the Contract DA J A 37-67-C-0628 of the Institute of Mathematics, University of Basel with the US Department of the Army, European Research Office.

[2]) This follows immediately from the relation

$$\left|\frac{B_{2n+1}(t)}{(2n+1)!}\right| < \frac{2}{(2\pi)^{2n+1}} \qquad (0 \leqq t \leqq 1,\ n \geqq 1).$$

As to this formula, it is for $n > 1$ equivalent with a formula deduced by LEHMER on p. 538 of his paper, LEHMER [2] (see the formula without number, following immediately after LEHMER's formula (18) in his paper). However, this relation is also true for $n = 1$, as is immediately verified, using for instance the value of $\operatorname*{Max}_{\langle 0,1\rangle}\left|\frac{B_3(t)}{6}\right| = .00801875$ (which is obtained from LEHMER's value of his M_3), since this this is $< \frac{1}{4\pi^3} = .0806$.

However, the inequality (5) can be further improved. It is easy to see that we have in any case

$$\max_{\langle 0,\frac{1}{2}\rangle} |U_s(t)| > 1 - \frac{1}{3^s} \quad (s > 1) \tag{6}$$

(see section 4).

As to the upper limit of $|U_s(t)|$, we will prove that we have

$$|U_s(t)| < 1 - \frac{\frac{1}{2}}{3^s} \quad (s \geqq 7). \tag{7}$$

On the other hand we have

$$\max_{\langle 0,\frac{1}{2}\rangle} |U_5(t)| = 1 - \frac{.496}{3^5}, \tag{8}$$

$$\max_{\langle 0,\frac{1}{2}\rangle} |U_3(t)| = 1 - \frac{.147}{3^3}. \tag{9}$$

3. The inequality (7) will be proved for all real s by a rather elementary method, making no use of the results obtained in the last decades on the zeros of the Bernoullian Polynomials of even order. (NÖRLUND [4], LENSE [3], LEHMER [2], INKERI [1], OSTROWSKI [5].) As to the values (8) and (9), they are obtained using the connection between the Bernoullian Polynomials $B_5(t)$ and $B_3(t)$ with $U_5(t)$ and $U_3(t)$.

4. We obtain at once from (3)

$$U_s(\tfrac{1}{4}) = \sum_{\nu=1}^{\infty} \frac{\sin \nu\pi/2}{\nu^s} = \frac{1}{1^s} - \frac{1}{3^s} + \left(\frac{1}{5^s} - \frac{1}{7^s}\right) + \cdots > 1 - \frac{1}{3^s} \quad (s > 1),$$

and this proves (6).

5. In order to prove (7) we introduce

$$\varphi(t) = \sin \pi t + \frac{\sin 2\pi t}{2^s} + \frac{\sin 3\pi t}{3^s} \tag{10}$$

and put

$$U_s(t/2) = \varphi(t) + R(t), \quad R(t) = \sum_{\nu=4}^{\infty} \frac{\sin \nu\pi t}{\nu^s}. \tag{11}$$

Then we have

$$|R(t)| \leqq \frac{1}{4^s} + \sum_{\nu=5}^{\infty} \frac{1}{\nu^s} < \frac{1}{4^s} + \int_4^{\infty} \frac{dx}{x^s}$$

$$= \frac{1}{4^s}\left(1 + \frac{4}{s-1}\right) \leqq \frac{\frac{5}{3}}{4^s} = \frac{5}{3}\cdot\frac{1}{3^s}\cdot\left(\frac{3}{4}\right)^s \leqq \frac{5}{3}\cdot\left(\frac{3}{4}\right)^7 \frac{1}{3^s},$$

$$|R(t)| \leqq \frac{.2225}{3^s} \quad (s \geqq 7). \tag{12}$$

6. We are now going to discuss $\operatorname*{Max}_{\langle -1,1\rangle} |\varphi(t)| = \operatorname*{Max}_{\langle 0,1\rangle} |\varphi(t)|$ and to prove that

$$|\varphi(t)| \leqq 1 - \frac{.7232}{3^s} \qquad (|t| \leqq 1,\ s \geqq 7). \tag{13}$$

Let

$$u = \frac{\pi}{2} - \pi t, \quad 0 \leqq |u| \leqq \frac{\pi}{2}, \quad \sigma = \sin u, \quad \gamma = \cos u. \tag{14}$$

Then we have

$$\varphi(t) = \gamma K, \quad K = 1 - \frac{1}{3^s} + \frac{2\sigma}{2^s} + \frac{4\sigma^2}{3^s}, \tag{15}$$

$$f(u) := -\frac{d}{du}\varphi(t) = \sin u - \frac{3\sin 3u}{3^s} - \frac{2\cos 2u}{2^s}, \tag{16}$$

$$F_2(\sigma) := 3^{s-1} f(u) = 4\sigma^3 + 2\left(\frac{3}{2}\right)^{s-1}\sigma^2 + (3^{s-1} - 3)\,\sigma - \left(\frac{3}{2}\right)^{s-1}. \tag{17}$$

As $F_2(1) > (\frac{3}{2})^{s-1} > 0$ and by Descartes' rule, $F_2(\sigma)$ has exactly one positive zero, σ_0, which is <1. On the other hand, as $F_2'(\sigma) = 12\sigma^2 + 4(\frac{3}{2})^{s-1}\sigma + 3^{s-1} - 3$, we have for $|\sigma| \leqq 1$ and $s \geqq 7$:

$$|F_2'(\sigma)| \geqq 3^{s-1} - 3 - 12 - 4\left(\frac{3}{2}\right)^{s-1} = 3^{s-1}\left(1 - \frac{1}{2^{s-3}}\right) - 15 \geqq 729 - \frac{15}{16} - 15 > 0,$$

and we see that $F_2(\sigma)$ cannot have a zero in $\langle -1, 0\rangle$.

7. Putting $\sigma = \dfrac{x}{2^{s-1}}$ into $F_2(\sigma)$ we obtain

$$F_1(x) := 6^{s-1} f(u) = \frac{x^3}{4^{s-2}} + 2\left(\frac{3}{4}\right)^{s-1} x^2 + (3^{s-1} - 3)\,x - 3^{s-1},$$

$$F_1(1) = \frac{1}{4^{s-2}} + 2\left(\frac{3}{4}\right)^{s-1} - 3 < 0.$$

We see that $F_1(x)$ has exactly one positive zero, ξ, which is >1. Writing $x = 1 + y$, $F_1(x) = 3F(y)$ we obtain

$$F(y) = \frac{16}{3}\frac{y^3}{4^s} + \left[\frac{16}{4^s} + \frac{8}{9}\left(\frac{3}{4}\right)^s\right] y^2$$
$$+ \left[\frac{16}{4^s} + \frac{16}{9}\left(\frac{3}{4}\right)^s + \frac{1}{9}3^s - 1\right] y - \left[1 - \frac{16}{3}\frac{1}{4^s} - \frac{8}{9}\left(\frac{3}{4}\right)^s\right].$$

For positive y it follows

$$F(y) > (\tfrac{1}{9}3^s - 1)\,y - (1 - \tfrac{8}{9}(\tfrac{3}{4})^s)$$

and therefore
$$F\left(\frac{2}{2^s}\right) > \frac{2}{9}\left(\frac{3}{2}\right)^s - 1 + \frac{8}{9}\left(\frac{3}{4}\right)^s - \frac{2}{2^s} > 0 \quad (s \geqq 7).$$

8. We see that $\xi = 1 + \frac{2\theta}{2^s}$,
$$\sigma_0 = \frac{2}{2^s} + \theta\frac{4}{4^s}, \quad 0 < \theta < 1, \quad (s \geqq 7), \tag{18}$$
$$\sigma_0 = \frac{2}{2^s}\left(1 + \frac{2\theta}{2^s}\right) < \frac{2}{2^s}\left(1 + \frac{1}{2^6}\right) = \frac{2}{2^s}\left(1 + \frac{1}{64}\right) \quad (s \geqq 7). \tag{19}$$

Since $\varphi(t)$ vanishes for $t=0$ and $t=1$, the maximum of $\varphi(t)$ is attained at a point where $f(u)$ vanishes, that is to which corresponds $\sin u = \sigma_0$. The corresponding values of $\gamma = \cos u$ is then
$$\sqrt{1-\sigma_0^2} < 1 - \sigma_0^2/2 < 1 - \frac{2}{4^s}.$$

9. We now have from (15)
$$|\varphi(t)| < \left(1 - \frac{2}{4^s}\right)K,$$
$$K = 1 - \frac{1}{3^s} + K_1, \quad K_1 = \frac{2\sigma_0}{2^s} + \frac{4\sigma_0^2}{3^s} = \frac{2\sigma_0}{2^s}\left(1 + \frac{2\sigma_0}{(\frac{3}{2})^s}\right)$$
$$< \frac{4}{4^s}\left(1 + \frac{1}{64}\right)\left(1 + \frac{4 + \frac{1}{8}}{3^7}\right) < \left(1 + \frac{1}{56}\right)\frac{4}{4^s} < \frac{\frac{57}{14}}{4^s},$$
$$K < 1 - \frac{1}{3^s} + \frac{\frac{57}{14}}{4^s},$$
$$|\varphi(t)| < \left(1 - \frac{2}{4^s}\right)\left(1 - \frac{1}{3^s} + \frac{\frac{57}{14}}{4^s}\right) = 1 - \frac{1}{3^s} + \frac{\frac{29}{14}}{4^s} + \frac{2}{12^s} - \frac{8\frac{1}{7}}{16^s}$$
$$< 1 - \frac{1}{3^s} + \frac{1}{3^s}\left(\frac{29}{14}\left(\frac{3}{4}\right)^s + \frac{2}{4^s}\right) \leqq 1 - \frac{1}{3^s} + \frac{1}{3^s}\left(\frac{29}{14}\left(\frac{3}{4}\right)^7 + \frac{2}{4^7}\right),$$
$$|\varphi(t)| < 1 - \frac{1}{3^s} + \frac{.2766}{3^s} = 1 - \frac{.7234}{3^s},$$
and by (12) and (13)
$$|U_s(t)| < 1 - \frac{.7234}{3^s} + \frac{.2225}{3^s} = 1 - \frac{.5009}{3^s},$$
which proves (7).

It is of some interest to observe that (7) and our proof of (7) remain valid if $U_s(t) = \sum_{\nu=1}^{\infty} \frac{\sin 2\nu\pi t}{\nu^s}$ is replaced with $\sum_{\nu=1}^{N} \frac{\sin 2\nu\pi t}{\nu^s}$ for any $N \geqq 3$.

9. To verify (8) and (9) we use the fact that (3) is in $\langle 0, 1\rangle$ a polynomial, the so-called Bernoullian Polynomial $B_{2n+1}(t)$. We have in particular $B_3(t)=t^3-\frac{3}{2}t^2+\frac{1}{2}t$, $B_5(t)=t^5-\frac{5}{2}t^4+\frac{5}{3}t^3-\frac{1}{6}t$. The maximum of $|B_3(t)|$ in $\langle 0, 1\rangle$ is attained at $t=.211325$ and gives $|B_3(0.211325)|=.048113$. Since $\frac{2\pi^3}{3}=20.67085$, we obtain $\operatorname*{Max}_{\langle 0,1\rangle}|U_s(t)|=0.99454=1-\frac{.147}{27}$, which gives (9).

The maximum of $|B_5(t)|$ in $\langle 0, 1\rangle$ is attained at $t=.240335$ and the corresponding value of $|B_5(t)|$ is .0244582. Since $\frac{2\pi^5}{15}=40{,}8026$, we obtain

$$\operatorname*{Max}_{\langle 0,1\rangle}|U_5(t)| = .997958 = 1-\frac{.496}{243},$$

which proves (8).

REFERENCES

[1] K. Inkeri, *The real roots of Bernoulli polynomials*, Ann. Univ. Turkuensis [Series A], Nr 37 (Turku 1959).

[2] D. H. Lehmer, On the Maxima and Minima of Bernoulli polynomials, Am. Math. Monthly *47* (1940), 533–538.

[3] J. Lense, *Über die Nullstellen der Bernoullischen Polynome*, Wiener Monatshefte f. Mathematik *41* (1934), 188–190.

[4] N. E. Nörlund, *Mémoire sur les Polynômes de Bernoulli*, Acta Math. *43* (1920), 121–196.

[5] A. M. Ostrowski, *On the zeros of Bernoulli Polynomials of Even Order*, L'Enseignement Mathématique *VI* (1960), 27–47.

[6] S. D. Poisson, *Sur le calcul numérique des intégrales definies*, Mémoires Acad. Paris *VI* (1827), 571–602.

State University of New York at Buffalo
September 1968

Received Sept. 26, 1968