

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 44 (1969)

Artikel: A Note on a Paper of J.T. Marti.
Autor: Caradus, S.R.
DOI: <https://doi.org/10.5169/seals-33772>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A Note on a Paper of J. T. MARTI.

by S. R. CARADUS

Recently, J. T. MARTI showed [1] that, under a certain commutativity condition on the operators involved, it was possible to obtain an operational calculus for pairs of closed linear operators. The purpose of this note is to observe that MARTI's conclusions are valid under more general conditions which, moreover, permit extension of the theory to n -tuples of operators.

Let X be a Banach space and T_i closed linear operators with domains D_i and ranges in X , $i = 1, 2$. We shall also write D_{12} for the domain of $T_1 T_2$ i.e. $D_{12} = \{x \in D_2 : T_2 x \in D_1\}$. Similarly we define D_{21} . Following [1], we define

$$\mathcal{C}(X) = \{(T_1, T_2) : D_1 \subseteq D_2; T_2 D_1 \subseteq D_1; T_1 T_2 x = T_2 T_1 x \text{ for all } x \in D_{21}\}.$$

Then for $(T_1, T_2) \in \mathcal{C}(X)$, suppose $\sigma_e(T_i) \subseteq V_i$, $i = 1, 2$, where V_i is a Cauchy domain and let $H(V)$ denote the complex algebra of functions $\mathbf{C}^2 \rightarrow \mathbf{C}$, locally holomorphic in $V = V_1 \times V_2$, vanishing outside V . The main result of [1], (Theorem 2), asserts the existence of a proper homomorphism Φ of $H(V)$ into $B(X)$, mapping $(\lambda - \lambda_i)^{-1} | V$ onto $(\lambda - T_i)^{-1}$ for each $\lambda \notin V_i$ such that Φ is given by a Cauchy integral. The essential step is to show that the assumption $(T_1, T_2) \in \mathcal{C}(X)$ implies that the resolvent operators $R_i = (\lambda_i - T_i)^{-1}$ commute, $\lambda_i \in \rho(T_i)$, $i = 1, 2$.

PROPOSITION. *The resolvent operators R_1 and R_2 commute if and only if*

$$(i) \quad D_1 \cap D_{12} = D_2 \cap D_{21}$$

and

$$(ii) \quad T_1 T_2 x = T_2 T_1 x \text{ for } x \in D_{12} \cap D_{21}$$

Proof. Suppose R_1 and R_2 commute. Then it is easy to verify that, for any x , $R_1 R_2 x \in D_{21}$. For D_i is the range of R_i so that $R_1 R_2 x \in D_1$ and hence $(\lambda_1 - T_1) R_1 R_2 x$ is defined and equal to $R_2 x$ which lies in D_2 . Thus $R_1 R_2 x \in D[T_2(\lambda_1 - T_1)] = D_2 \cap D_{21}$. Similarly $R_2 R_1 x \in D_{12}$ so that since $R_1 R_2 x = R_2 R_1 x$, we have $R_1 R_2 x \in D_{12} \cap D_{21}$.

Consider now the equation

$$0 = (R_1 R_2 - R_2 R_1)x = R_1 R_2[(\lambda_1 - T_1)(\lambda_2 - T_2) - (\lambda_2 - T_2)(\lambda_1 - T_1)]R_2 R_1 x. \quad (1)$$

Since $R_2 R_1 x \in D_{12} \cap D_{21}$, (1) is valid for all x and can be rewritten

$$0 = R_1 R_2[T_2 T_1 - T_1 T_2]R_2 R_1 x.$$

This implies

$$[T_2 T_1 - T_1 T_2]R_2 R_1 x = 0 \text{ for all } x \in X.$$

But every $y \in D_{12} \cap D_{21}$ can be expressed as $R_2 R_1 x$. For let $x = (\lambda_1 - T_1)(\lambda_2 - T_2)y$. Hence condition (ii) is satisfied.

To prove (i), let $x \in D_1 \cap D_{12}$ and define $x' = (\lambda_1 - T_1)(\lambda_2 - T_2)x$. Clearly x' is well defined since $D[(\lambda_1 - T_1)(\lambda_2 - T_2)] = D_1 \cap D_{12}$. Hence $x = R_2 R_1 x' = R_1 R_2 x'$. But $R_1 R_2 x' \in D_2 \cap D_{21}$. Hence $D_1 \cap D_{12} \subseteq D_2 \cap D_{21}$. The reverse inclusion is obtained similarly.

Now suppose that conditions (i) and (ii) hold. Let $x \in X$. Then there exists $x_2 \in D_2$ such that $(\lambda_2 - T_2)x_2 = x$ and $x_1 \in D_1$ such that $(\lambda_1 - T_1)x_1 = x_2$. Hence $x = (\lambda_2 - T_2) \times (\lambda_1 - T_1)x_1$ and evidently $x_1 \in D_{21} \cap D_2$. By assumption (i), $(\lambda_1 - T_1)(\lambda_2 - T_2)x_1$ is defined and by assumption (ii), it must equal x . Hence $x_1 = R_1 R_2 x = R_2 R_1 x$.

COROLLARY 1. *Theorem 2 of [1] is valid whenever T_1 and T_2 satisfy conditions (i) and (ii).*

Proof. An examination of the proof of Theorem 2 shows that the commutativity assumption involved in the definition of $\mathcal{C}(X)$ was used only to establish the commutativity of R_1 and R_2 .

COROLLARY 2. *If $(T_1, T_2) \in \mathcal{C}(X)$, then $T_1 D_1 \subseteq D_2$.*

Proof. By [1] Lemma 1, R_1 and R_2 commute. Hence, in particular, by the above proposition, $D_1 \cap D_{12} = D_2 \cap D_{21}$. But $D_1 \cap D_{12} = \{x \in D_1 \cap D_2 : T_2 x \in D_1\} = D_1 \cap D_2 = D_1$. Hence $D_1 = D_2 \cap D_{21}$ and since $D_{21} \subseteq D_1 \subseteq D_2$, we must have $D_{21} = D_1$ i.e. $D_1 = \{x \in D_1 : T_1 x \in D_2\}$ i.e. $T_1 D_1 \subseteq D_2$.

COROLLARY 3. *Theorem 2 of [1] can now be extended in the obvious way to an operational calculus of n -tuples (T_1, T_2, \dots, T_n) of closed operators such that pairs (T_i, T_j) satisfy conditions (i) and (ii). The other results of [1] also have obvious extensions.*

REFERENCE

- [1] J. T. MARTI, *Operational Calculus for Two Commuting Closed Operators*, Comment. Math. Helv. 43 (1968), 87–97.

*Institute of Advanced Studies,
Australian National University,
Canberra*

Received July 19, 1968