

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 45 (1970)

Erratum: Addendum to "On Meromorphic Solutions of Algebraic Differential Equations".
Autor: Bank, Steven

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Addendum to „On Meromorphic Solutions of Algebraic Differential Equations”

by STEVEN BANK

In this note, we make two additional remarks concerning the paper [1].

1. An easy modification of the proof in [1] can be used to establish the following important generalization of the main theorem in [1]:

THEOREM: *Let $\Lambda(z, y, dy/dz) = \sum Q_{kj}(z) y^k (dy/dz)^j$ be a polynomial in y and dy/dz whose coefficients are functions of z which are subject to the following restriction: If p is the total degree of Λ in y and dy/dz , let $Q_{kj}(z)$ be an entire function of finite order if $k+j < p$, while if $k+j=p$, let $Q_{kj}(z)$ be a polynomial. Then a meromorphic solution of the differential equation $\Lambda(z, y, dy/dz)=0$ cannot be written as the quotient of two entire functions f/g , where f is of infinite order and g is of finite order. In particular, any entire solution, and more generally, any meromorphic solution whose sequence of poles has a finite exponent of convergence, must be of finite order of growth.*

2. Professor A. A. Gol'dberg has kindly informed the author that the question raised in [1], concerning the growth of meromorphic solutions of first order equations in the case when all coefficients $Q_{kj}(z)$ are polynomials, was solved in his paper [2], where it was shown that when all the coefficients are polynomials, then all meromorphic solutions must be of finite order. However, the author would like to point out that for the more extensive class of equations treated in Part 1 of this “Addendum”, the results obtained in [2] cannot be used to establish the theorem given in Part 1, since some coefficients are allowed to have essential singularities at ∞ , and, in fact, it is easy to see that such equations can possess meromorphic solutions of infinite order (e.g. $(\sin(e^z))^{-1}$ is a solution of the equation, $y^4 - y^2 - e^{-2z} (dy/dz)^2 = 0$).

REFERENCES

- [1] S. BANK, *On meromorphic solutions of algebraic differential equations*, Comm. Math. Helv., 44 (1969), 401–409.
- [2] A. A. GOL'DBERG, *On one-valued integrals of differential equations of the first order*, Ukrain. Mat. Ž., 8 (1956), 254–261 (Russian).

*University of Illinois
Urbana, Illinois.*

Received May 30, 1970.