

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 46 (1971)

Artikel: The Spectra of Hyponormal Integral Operators
Autor: Clancey, K.F. / Putnam, C.R.
DOI: <https://doi.org/10.5169/seals-35534>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Spectra of Hyponormal Integral Operators¹⁾

K. F. CLANCEY and C. R. PUTNAM

1. Recall that a bounded operator $T = H + iJ$ on a Hilbert space \mathfrak{H} is said to be hyponormal if

$$T^*T - TT^* = D \geq 0, \text{ that is, } HJ - JH = -iC, C = \frac{1}{2}D \geq 0. \quad (1.1)$$

It is known that such operators behave to some extent like normal operators; in particular, $\text{sp}(H)$ and $\text{sp}(J)$ are just the (real) projections of $\text{sp}(T)$ onto the real and imaginary axes; see Putnam [5b], p. 46.

Let H have the spectral resolution

$$H = \int \lambda dE_\lambda, \quad (1.2)$$

and let $E(\Delta)$ be the projection operator associated with an open interval Δ . For any bounded operator T (hyponormal or not), let $T_\Delta = E(\Delta)TE(\Delta)$, regarded as an operator on $E(\Delta)\mathfrak{H}$ and with spectrum $\text{sp}(T_\Delta)$. Since $H_\Delta J_\Delta - J_\Delta H_\Delta = -iC_\Delta$, it is seen that T_Δ is hyponormal on $E(\Delta)\mathfrak{H}$ whenever T is hyponormal on \mathfrak{H} . It was shown in [5d] that if T is hyponormal, then

$$\text{sp}(T_\Delta) \subset \text{sp}(T). \quad (1.3)$$

In case the self-commutator D of T in (1.1) is compact, the relation (1.3) was proved by Clancey [2a].

A refinement of (1.3) was proved in [5f] to the following

$$\text{sp}(T_\Delta) \cap \{z: \text{Re}(z) \in \Delta\} = \text{sp}(T) \cap \{z: \text{Re}(z) \in \Delta\}, \quad (1.4)$$

Δ being any open interval. In view of the projection properties mentioned above, the real part of $\text{sp}(T_\Delta)$ lies in the closure of Δ . It was noted in [5f] that, as a consequence of (1.4),

$$\text{Im}[\text{sp}(T) \cap \{z: \text{Re}(z) = s\}] = \bigcap_{\Delta} \text{sp}(E(\Delta)JE(\Delta)), \quad s \in \Delta, \quad (1.5)$$

the intersection being over all open intervals Δ containing s . This relation will be used below to determine the spectra of certain singular integral operators.

Suppose that

$$a(x), b(x) \in L^\infty(E), a(x) \text{ real, } b(x) \neq 0 \text{ a.e. on } E, \quad (1.6)$$

¹⁾ This work was supported by National Science Foundation research grants.

where E is a bounded set of positive measure on the real line. Let $T_0 = H_0 + iJ_0$ denote the bounded operator on $L^2(E)$ defined by

$$(H_0 f)(x) = xf(x) \quad \text{and} \quad (J_0 f)(x) = - \left[a(x)f(x) + \frac{b(x)}{i\pi} \int_E \frac{\bar{b}(t)}{t-x} f(t) dt \right], \quad (1.7)$$

where the integral is interpreted as a Cauchy principal value. It is easily verified that

$$H_0 J_0 - J_0 H_0 = -iC_0, \quad C_0 f = \pi^{-1}(f, b) b, \quad (1.8)$$

so that $C_0 \geq 0$ and hence T_0 is hyponormal. It is seen that the range of C_0 is spanned by the vector $b \in L^2(E)$ and that $H_0 = x$ has simple spectrum and that the vectors $\{H_0^n b\}$, $n=0, 1, 2, \dots$ span $L^2(E)$.

Conversely, if $T = H + iJ$ is any hyponormal operator on H satisfying

$$T^*T - TT^* = D \geq 0 \quad \text{and } D \text{ has rank one} \quad (1.9)$$

and

$$D = (\cdot, z) z \text{ and } \{H^n z\}, \quad n = 0, 1, 2, \dots, \text{ span } H, \quad (1.10)$$

then T is unitarily equivalent to a singular integral operator $T_0 = H_0 + iJ_0$ defined by (1.7). This result was first proved by Xa Dao-xeng [7]; a simpler proof using a result in [5a] was given by Rosenblum [6], p. 326.

It may be noted that the operator T_0 above is irreducible by virtue of the condition that $b(x) \neq 0$ a.e. on E . To see this, note that if $\Omega \neq 0$ reduces T_0 , then Ω reduces both H_0 and J_0 . If $f \in \Omega$, $f \neq 0$ (that is, $f(x) \neq 0$ a.e.) and if $(f, b) \neq 0$, then $(C_0 f)(x) = \pi^{-1}(f, b) b(x) \neq 0$ a.e. on E , and hence $\{(H_0^n C_0 f)(x)\}$, $n=0, 1, 2, \dots$, span the space $L^2(E)$, that is, $\Omega = L^2(E)$. If $(f, b) = 0$, then, since $f \neq 0$, $C_0 H_0^N f \neq 0$ for some positive integer N . Otherwise, by Weierstrass' theorem, $f(x) b(x) \equiv 0$ a.e. and hence, $f(x) = 0$ a.e., a contradiction. Thus, if $g = H_0^N f \neq 0$, one can proceed as above to show that $\Omega = L^2(E)$.

THEOREM 1. *Let $T_0 = H_0 + iJ_0$ be the hyponormal operator on $L^2(E)$ defined by (1.6) and (1.7). Then $\text{sp}(T_0)$ is the set of numbers $z = s + it$ (s, t real) for which*

$$\text{meas}_1 \{x \in E \cap \Delta : -a(x) - |b(x)|^2 - \varepsilon < t < -a(x) + |b(x)|^2 + \varepsilon\} > 0 \quad (1.11)$$

holds for every $\varepsilon > 0$ and for every open interval Δ containing s .

THEOREM 2. *Let T_0 be defined as in Theorem 1. Then for almost all points $x \in E$, there exists some vertical segment $\{x + iy : a_x \leq y \leq b_x\}$, where $a_x < b_x$, belonging to the spectrum of T_0 . In particular, $\text{sp}(T_0)$ cannot be totally disconnected.*

Theorem 1 generalizes results of Clancey [2a], Theorem 1 and Putnam [5c]. Its proof will be given in section 2. In a formulation involving a “determining set” or “determining function”, Theorem 1 is contained in Clancey [2b] and Pincus [3c]. All of these proofs, including the one of the present paper, use results of either Pincus [3a] or Rosenblum [6] together with the relation (1.4) (or (1.5)) established in [5f]. It may also be noted that in [3c], the operator D of (1.9) is assumed only to be of trace class, rather than of rank one, and that \mathfrak{H} is the least subspace reducing T and containing the range of D .

A hyponormal operator T is said to be completely hyponormal on \mathfrak{H} if there is no non-trivial subspace of \mathfrak{H} which reduces T and on which T is normal. A set S of the complex plane is said to have positive density if for every open disk N ,

$$\text{meas}_2(S \cap N) > 0 \quad \text{whenever } S \cap N \neq \emptyset. \quad (1.12)$$

It was shown in [5d] that if T is completely hyponormal then its spectrum has positive density. The converse question of whether every compact set S is the spectrum of some completely hyponormal operator is unsettled, although some partial results have been obtained; see [5g], also Theorem 3 below and the remarks in section 4.

For any set S , let S^- denote its closure and $\text{int}(S)$ its interior. There will be proved the following

THEOREM 3. *If S is any compact set for which*

$$S = (\text{int}(S))^- \quad (1.13)$$

(so that, in particular, S has positive density), then there exists a singular integral operator $T_0 = H_0 + iJ_0$ defined by (1.6) and (1.7) for which

$$\text{sp}(T_0) = S. \quad (1.14)$$

2. Proof of Theorem 1. It follows from Pincus [3a], p. 375, that $t \in \text{sp}(J_0)$, where J_0 is defined by (1.7), if and only if

$$\text{meas}_1 \{x \in E: -a(x) - |b(x)|^2 - \varepsilon < t < -a(x) + |b(x)|^2 + \varepsilon\} > 0$$

for every $\varepsilon > 0$. (In this connection, see also Rosenblum [6], p. 323; also the remarks in Pincus and Rovnyak [4], p. 620.) If the multiplication operator $H = H_0 = x$ of (1.7) has the spectral resolution (1.2) then for any open interval Δ (for which $E \cap \Delta \neq \emptyset$), $E(\Delta) J_0 E(\Delta)$ is simply the integral operator J_0 restricted to $E \cap \Delta$. It follows that the condition $t \in \text{sp}(E(\Delta) J_0 E(\Delta))$ reduces to (1.11), and Theorem 1 now follows from (1.5).

Proof of Theorem 2. Since $b(x) \neq 0$ a.e. on E , then

$$E = \bigcup_{n=1}^{\infty} E_n, \text{ a.e., where } E_n = \{x \in E : |b(x)|^2 > 1/n\} \text{ for } n = 1, 2, \dots.$$

Hence, $E_1 \subset E_2 \subset \dots$ and $\text{meas}_1(E - E_n) \rightarrow 0$ as $n \rightarrow \infty$. Choose N so large that $\text{meas}_1(E_n) > 0$ for $n \geq N$. Thus, at almost all $x \in E_n$, where $n \geq N$, E_n has metric density 1. For such an x , let $L = \text{ess lim sup } a(t)$, where $t \rightarrow x$ and t is restricted to E_n . Then, in every open interval containing x and for every $\varepsilon > 0$, there exists a subset of E of positive measure for which $|a(x) - L| < \varepsilon$ and $|b(x)|^2 > 1/N$. It follows from the criterion of (1.11) that the segment $x + iy$, where $L - 1/N \leq y \leq L + 1/N$, belongs to the spectrum of T_0 .

3. Proof of Theorem 3. For any Borel set α of the line, let $S(\alpha)$ denote the set $S(\alpha) = S \cap \{z : \text{Re}(z) \in \alpha\}$. For $k = 1, 2, \dots$, let Π_k denote a grid of squares in the complex plane with sides parallel to the axes and of length 2^{-k} . We assume that the squares contain their lower and left sides and that $z = 0$ is a lower left corner of some square in each grid. Since S is compact then the projection on the x -axis of S is contained in some interval $[c, d]$. Now choose a disjoint family $\{K_p\}$, $p = 1, 2, \dots$, of Cantor sets of positive measure in $[c, d]$ so that

$$\text{meas}_1 \left(\bigcup_{p=1}^q K_p \right) \rightarrow d - c \quad \text{as } q \rightarrow \infty. \quad (3.1)$$

Denote by R_1, \dots, R_{n_1} the elements of Π_1 satisfying

$$R_j \subset \text{int}(S) \equiv \Omega_1, \quad j = 1, \dots, n_1, \quad (3.2)$$

and let R'_1, \dots, R'_{n_1} be respective smaller concentric closed squares of side 2^{-2} . Then for $j = 1, \dots, n_1$, let K_{p_j} be the first K_p satisfying

$$\text{meas}_2(S(K_p) \cap R_j) > 0 \quad \text{and} \quad p_j > p_{j-1}. \quad (3.3)$$

Set $A_j = S(K_p) \cap R'_j$ and let D_j be the projection on the x -axis of A_j . Clearly, the set $\Omega_2 = \Omega_1 - \bigcup_{j=1}^{n_1} A_j$ is open. Denote by R_j , for $j = n_1 + 1, \dots, n_1 + n_2$, the squares in Π_2 satisfying

$$R_j \subset \Omega_2, \quad j = n_1 + 1, \dots, n_1 + n_2. \quad (3.4)$$

Again, form concentric squares $R'_{n_1+1}, \dots, R'_{n_1+n_2}$ of side 2^{-3} and, for $j = n_1 + 1, \dots, n_1 + n_2$, let K_{p_j} be the first K_p satisfying (3.3). Repeat the process of forming A_j and D_j for $j = n_1 + 1, \dots, n_1 + n_2$ and set $\Omega_3 = \Omega_2 - \bigcup_{j=1}^{n_1+n_2} A_j$. If this process is continued for each q and grid Π_q one obtains a family of closed sets $\{A_j\}$, $j = 1, 2, \dots$, satisfying

$$\text{closure} \left(\bigcup_{j=1}^{\infty} A_j \right) = S. \quad (3.5)$$

Now define functions $a(x)$ and $b(x)$ on $\bigcup D_j$ by setting

$$\left. \begin{aligned} a(x) &= (\text{value of } y\text{-coordinate of the center of } R'_j) \text{ on } D_j, \\ b(x) &= (\text{one-half the length of the side of } R'_j)^{1/2} \text{ on } D_j. \end{aligned} \right\} \quad (3.6)$$

Then if T_0 is the singular integral operator given by (1.7) and (3.6) acting on $L^2(\bigcup D_j)$ it follows from Theorem 1 that relation (1.14) holds.

4. Remarks. It was shown in [5g] that there exist irreducible hyponormal operators satisfying (1.9) and having totally disconnected spectra. (An example was also given in [5e].) In view of the last part of Theorem 1, such an operator $T=H+iJ$ cannot be of the type $T_0=H_0+iJ_0$ defined by (1.6) and (1.7). That is, by the result of Xa Dao-xeng, since T satisfies (1.9), then relation (1.10) fails to hold.

It was shown in Theorem 3 that any compact set equal to the closure of its interior is the spectrum of some singular integral operator $T_0=H_0+iJ_0$ defined by (1.6) and (1.7). Of course, the spectrum of a general such operator need not be of this type; indeed, if $a(x)=0$ and if $b(x)$ is the characteristic function of a Cantor set E of positive measure, then (cf. Theorem 1) the spectrum of T_0 is the set $E \times [-1, 1]$.

It is interesting to note that although the spectrum of T_0 cannot be totally disconnected, nevertheless, it may be a Mergelyan Swiss cheese. (Recall that this is a set $X=D-\bigcup_{n=1}^{\infty} D_n$ where D is the closed unit disk and the D_n are open disjoint disks in D with radii r_n satisfying $\sum r_n < \infty$, and for which X is nowhere dense; see Zalcman [8], p. 69.) The proof of this assertion depends upon a result of W. K. Allard (see Brennan [1], p. 13) that almost every cross-section of a Swiss cheese is the union of a finite number of disjoint closed intervals; for details, see Clancey [2b].

REFERENCES

- [1] BRENNAN, J. E., *Point evaluations and invariant subspaces*, Thesis, Brown University (1968).
- [2a] CLANCEY, K., *Seminormal operators with compact self-commutators*, Proc. Amer. Math. Soc. 26 (1970), 447-454.
- [b] ——, *On the determining set in the theory of commutators*, preprint (1971).
- [3a] PINCUS, J. D., *Commutators, generalized eigenfunction expansions and singular integral equations*, Trans. Amer. Math. Soc. 113 (1966), 358-377.
- [b] ——, *Commutators and systems of singular integral equations, I*, Acta Math. 121 (1968), 219-249.
- [c] ——, *The spectrum of seminormal operators*, preprint (1971).
- [4] PINCUS, J. D. and ROVNYAK, J., *A spectral theory for some unbounded self-adjoint singular integral operators*, Amer. J. Math. 91 (1969), 619-636.
- [5a] PUTNAM, C. R., *Commutators, absolutely continuous spectra, and singular integral operators*, Amer. J. Math. 86 (1965), 310-316.
- [b] ——, *Commutation properties of Hilbert space operators and related topics*, Ergebnisse der Math. 36 (1967), Springer.
- [c] ——, *The spectra of seminormal singular integral operators*, Canad. J. Math. 22 (1970), 134-150.
- [d] ——, *An inequality for the area of hyponormal spectra*, Math. Z. 116 (1970), 323-330.
- [e] ——, *The spectra of subnormal operators*, Proc. Amer. Math. Soc. 28 (1971), 473-477.

- [f] ——, *A similarity between hyponormal and normal spectra*, Ill. J. Math. (to appear).
- [g] ——, *The spectra of completely hyponormal operators*, Amer. J. Math. (to appear).
- [6] ROSENBLUM, M., *A spectral theory for self-adjoint singular integral operators*, Amer. J. Math. 88 (1966), 314-328.
- [7] DAO-XENG, XA, *On non-normal operators*, Chinese Math. 3 (1963), 232-246.
- [8] ZALCMAN, L., *Analytic capacity and rational approximation*, Lecture Notes in Mathematics 50 (1968), Springer.

*University of California, Los Angeles
Purdue University*

Received March 26, 1971