

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 46 (1971)  
  
**Artikel:** On the Absolute Continuity of a Surface Representation  
**Autor:** Reimann, Hans Martin  
**DOI:** <https://doi.org/10.5169/seals-35504>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 20.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## On the Absolute Continuity of a Surface Representation

by HANS MARTIN REIMANN

This note contains an example of a 2-dimensional surface in 3-space, which is represented by an absolutely continuous (in the sense of Tonelli) homeomorphism  $f$ . Although the surface has finite Lebesgue area and  $f$  is a mapping “of bounded distortion” with  $L^2$  – integrable partial derivatives, there exists a 2-dimensional zero set which is mapped onto a set of positive 2-dimensional Hausdorff measure.

A real valued continuous function  $f$  defined in a bounded domain  $G^k$  in  $k$ -dimensional Euclidean space  $E^k$  is absolutely continuous in the sense of Tonelli if:

- (i) Given any closed interval  $I^k \subset G^k$ ,  $I^k = \{(x_1, \dots, x_k) | a_i \leq x_i \leq b_i, i = 1, \dots, k\}$   $f$  is absolutely continuous as a function of  $x_i$  on a.e. line parallel to the  $x_i$  axis;  $i = 1, \dots, k$ ;
- (ii) The partial derivatives which exist a.e. are integrable in  $G^k$ . For mappings  $f = (f_1, \dots, f_n): G^k \rightarrow E^n$  we write  $f \in ACL^p$  ( $p > 1$ ), if all coordinate functions  $f_i$ ,  $i = 1, \dots, n$ , are absolutely continuous in the sense of Tonelli and furthermore the partial derivatives are integrable to the power  $p$ .

Cesari [1952] proved that mapping  $s f \in ACL^p$ ,  $p > 2$ ,  $f: G^2 \rightarrow E^2$  have the following property: Every subset of  $G^2$  of zero (2-dim.) measure is mapped onto a set of zero measure. We will refer to this property by saying that  $f$  satisfies condition  $N$  with respect to 2-dimensional Lebesgue measure  $m_2: N(m_2)$ . In the same paper Cesari presented examples of mappings  $f \in ACL^2$ ,  $f: G^2 \rightarrow E^2$ , which do not satisfy condition  $N(m_2)$  and give rise to further phenomena. Some of Cesari's examples are based on conformal representations as the one below.

Cesari's result carries over to higher dimensions: Calderon [1951] has shown that mappings  $f \in ACL^p$ ,  $p > k$ ,  $f: G^k \rightarrow E^k$  are generalized Lipschitzian in the sense of Rado-Reichelderfer [1955]. From their results it then follows that  $f$  satisfies condition  $N(m_k)$ . This result still holds if  $f \in ACL^p$ ,  $p > k$ , is a mapping  $f: G^k \rightarrow E^n$ ,  $n > k$ . Condition  $N$  is then satisfied with respect to  $k$ -dimensional Hausdorff measure  $H_k$ .

If  $f \in ACL^k$  is a homeomorphism,  $f: G^k \rightarrow E^k$ , one can also conclude that  $f$  satisfies  $N(m_k)$ . This is well known for  $k = 2$  (for a proof see e.g. Lehto-Virtanen [1965] p.158). A proof for the case  $k > 2$  has been given by Reshetnjak [1966].

A mapping  $f \in ACL^k$ ,  $f: G^k \rightarrow E^k$  is said to be of bounded distortion if there exists a constant  $C \geq 1$  such that

$$|df|^k \leq C Jf$$

holds a.e. in  $G^k$ . Here  $Jf(x)$  is the (signed) Jacobian and  $|df(x)|$  is the norm of the linear transformation  $df(x)$ , which is given by the partial derivatives of  $f$  at  $x$ . For mappings  $f \in ACL^k$ ,  $f: G^k \rightarrow E^n$ ,  $n > k$ , we interpret this condition as  $|df|^k \leq C \|Jf\|$

a.e. in  $G^k$  with

$$\|Jf\| = \left( \frac{1}{k!} \sum \left[ \frac{\partial(f_{\alpha_1}, \dots, f_{\alpha_k})}{\partial(x_1, \dots, x_k)} \right]^2 \right)^{1/2},$$

where the sum in this expression extends over all multiindices  $\alpha = (\alpha_1, \dots, \alpha_k)$ ,  $1 \leq \alpha_i \leq n$ . (Intuitively  $\|Jf\|$  denotes the "surface element".) To guarantee that  $f: G^2 \rightarrow E^3$ ,  $f \in ACL^2$ , is of bounded distortion it is sufficient to verify that a.e. in  $G^2$

$$\sum_{i,j} \left( \frac{\partial f_i}{\partial x_j} \right)^2 \leq C' \|Jf\|$$

for some constant  $C'$ .

From Reshetnjak's work [1967] it is known that mappings  $f \in ACL^k$ ,  $f: G^k \rightarrow E^k$ , which are of bounded distortion, satisfy  $N(m_k)$ . The homeomorphisms of bounded distortion are the quasiconformal mappings (see e.g. Gehring [1962]). The investigation of extremal length properties of quasiconformal mappings leads to the following question: Do homeomorphisms  $f: G^k \rightarrow E^n$ ,  $n > k$ ,  $f \in ACL^k$ , which are of bounded distortion, satisfy condition  $N(H_k)$ ? The following example provides a negative answer to this question.

Let  $J$  be an Osgood curve, i.e. a closed Jordan curve in the plane with positive 2-dimensional measure.  $J$  separates the plane into a bounded and an unbounded component. We map the unit square  $Q = \{(x, y) \mid 0 < x < 1, 0 < y < 1\}$  conformally onto the bounded component  $J^0$ . By the Carathéodory extension theorem this mapping  $h$  can be extended continuously and one to one to a mapping  $h_c$  of the closed square  $\bar{Q}$  onto  $J^0 \cup J$ . Furthermore we can choose  $h$  in such a way as to have  $A = \{(x, y) \mid x = 0, 0 < y < 1\}$  mapped onto a set of positive 2-dimensional measure.

We define now the continuous mapping  $g = (u, v): R \rightarrow J^0 \cup J$  by setting  $R = \{(x, y) \mid 0 \leq |x| < 1, 0 < y < 1\}$  and

$$g(x, y) = \begin{cases} h_c(x, y) & \text{for } (x, y) \in Q \cup A \\ h(-x, y) & \text{otherwise} \end{cases}$$

Next we construct an auxiliary function  $w: R \rightarrow E^1$  in terms of the bounded positive function  $a(x, y) = \min\{1, |h'(x, y)|\}: Q \rightarrow E^1$ , where  $h = (u, v)$ ,  $|h'|^2 = |u_x v_y - u_y v_x| = u_x^2 + v_x^2 = u_y^2 + v_y^2 > 0$ . We define

$$w(x, y) = \begin{cases} \inf_{\gamma} \int_{\gamma} a(x, y) ds & \text{for } (x, y) \in Q \\ 0 & \text{for } (x, y) \in R \setminus Q \end{cases}$$

where the infimum is taken over all rectifiable curves  $\gamma \subset Q$  connecting  $(x, y)$  with  $A$ .  $w(x, y)$  is positive for all  $(x, y) \in Q$  since  $a(x, y)$  is positive and continuous in  $Q$ .

**THEOREM.** The mapping  $f=(u, v, w):R \rightarrow E^3$  constructed above has all the properties:

- a)  $f \in ACL^2$
- b)  $f$  is a homeomorphism
- c)  $f$  is of bounded distortion
- d)  $f$  maps the set  $A$  (with  $H_2(A)=0$ ) onto a set  $B$  with  $H_2(B)>0$ .

a)  $w$  satisfies a uniform Lipschitz condition with constant 1, hence  $w \in ACL^2$ .  $g=(u, v)$  is conformal in  $Q$  and maps  $Q$  onto a bounded domain. Therefore

$$\int_R |g'|^2 dx dy = 2 \int_Q |g'|^2 dx dy < \infty,$$

which means that the partial derivatives of  $u$  and  $v$  are square integrable. In order to show that  $g \in ACL^2$  it is sufficient to prove that for a.e.  $y$ ,  $0 < y < 1$ ,  $g(x, y)$  is absolutely continuous as a function of  $x$ . We choose  $y$  in such a way that

$V(y) = \int_{-1}^1 |g'(x, y)| dx < \infty$ . For these values the function  $g(x, y)$  is absolutely continuous in  $x$ , since it has an integral representation

$$g(x, y) = g(x, 0) + \int_0^x g'(t, y) dt$$

and the total variation  $V(y)$  is finite.

b) Because  $w(x, y) \neq 0$  for  $(x, y) \in Q$ ,  $f$  is a homeomorphism.

c)  $F$  satisfies the distortion condition  $|df|^2 \leq C \|Jf\|$  a.e. in  $R$ . For  $(x, y) \in R \setminus \bar{Q}$  this is clearly true for any constant  $C \geq 1$ . In the case  $(x, y) \in Q$  we obtain the following estimates:

$$\|Jf\| \geq |u_x v_y - u_y v_x| = |g'|^2$$

and

$$|w_x| \leq \left| \lim_{h \rightarrow 0} h^{-1} \int_x^{x+h} a(t, y) dt \right| \leq a(x, y) \leq |g'(x, y)|$$

From this we conclude

$$|(u_x, v_x, w_x)|^2 \leq |g'|^2 + a^2 \leq 2 |g'|^2.$$

An analogous relation holds for the derivatives with respect to  $y$  and therefore  $|df|^2 \leq C \|Jf\|$  for any  $C \geq 4$ . This clearly is not the best estimate. We remark that by replacing the function  $a(x, y)$  in the definition for  $w(x, y)$  by  $c \cdot a(x, y)$ ,  $c$  constant, we obtain  $C \rightarrow 1$  for  $c \rightarrow 0$ .

d)  $f$  does not satisfy condition  $N(H_2)$

The set  $A = \{(x, y) \mid x=0, 0 < y < 1\}$  has zero 2-dimensional measure ( $H_2(A)=0$ ) and  $f$  maps  $A$  onto a set  $B$  with  $H_2(B) > 0$ . (Observe that  $H_2(B) = m_2(B)$ , since  $B$  lies in the plane  $w=0$ .)

We add a few remarks:

1)  $f$  does not satisfy condition  $N$  with respect to 2-dimensional integralgeometric measure  $I_2$ . Using the characterization of  $I_2$  given by Federer [1947] p. 145, this statement can easily be verified.

2) Since  $f \in ACL^2$ , the Lebesgue area of  $f$  is given by  $L(f) = \int_R \|Jf\| \, dx \, dy$ .  $f$  therefore is an example of a homeomorphism with the property that  $L(f) \neq H_2(f(R))$ . A similar example of such a mapping has been constructed by Breckenridge [1970].

3)  $g: R \rightarrow E^2$  is another example of a mapping of the type described by Cesari:  $g \in ACL^2$  does not satisfy condition  $N(m_2)$ .

## REFERENCES

- BRECKENRIDGE, J. [1970] *Significant sets in surface area theory* (to appear).  
 CALDERON, A. [1951] *On the differentiability of absolutely continuous functions*, Riv. di Mat. Parma 2 p. 203–214.  
 CESARI, L. [1942] *Sulle trasformazioni continue*, Annali di Mat. pura ed appl. IV 21 p. 157–188.  
 GEHRING, F. W. [1962] *Rings and quasiconformal mappings in space*, Trans. Amer. Math. Soc. 103 p. 353–393.  
 LEHTO, O. und VIRTANEN, K. I. [1965] *Quasikonforme Abbildungen*, Springer Verlag.  
 RADO T. and REICHELDERFER, P. V. [1955] *Continuous transformations in analysis*, Springer Verlag.  
 RESHETNJAK, Y. G. [1966] *Some geometric properties of functions and mappings with generalized derivatives*, Sib. Mat. J. 7 p. 704–732 (English translation).  
 RESHETNJAK, Y. G. [1967] *Space mappings with bounded distortion*, Sib. Mat. J. 8 p. 466–487 (English translation).

*University of Michigan, Ann Arbor.*

Received April 27, 1970