

# Generators for Certain Ideals in Regular Rings of Dimension Three

Autor(en): **Murthy, M. Pavanan**

Objekttyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **47 (1972)**

PDF erstellt am: **27.04.2024**

Persistenter Link: <https://doi.org/10.5169/seals-36358>

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# Generators for Certain Ideals in Regular Rings of Dimension Three

by M. PAVANAN MURTHY<sup>1)</sup>

## Introduction

We prove here the following

**THEOREM.** *Let  $A$  be a regular ring of dimension 3 with  $\tilde{K}_0 A = 0$ . Let  $\mathfrak{a}$  be an unmixed ideal of height 2. Suppose  $\mathfrak{a}$  is locally generated by  $r$  elements. Then  $\mathfrak{a}$  is generated by  $r+1$  elements.*

Applying this theorem to  $A = K[x_1, x_2, x_3]$ ,  $K$  a field we obtain for instance that if  $C$  is a curve in the affine three space  $A_3$ , which is locally a complete intersection (e.g.  $C$  non-singular), then the ideal of  $C$  is generated by three elements. We also show that this is best possible by giving an example of a non-singular curve in  $A_3$  which is not a complete intersection.

In the case  $A = K[x_1, x_2, x_3]$ ,  $K$  algebraically closed and  $\mathfrak{a}$  the ideal of a non-singular curve, S. Abhyankar has proved this by quite different methods (see his Montreal Lecture Notes [1]).

A basic tool in the proof is a lemma of Serre [6] which relates projective modules with generators of certain ideals of height 2. In fact for  $r \geq 3$ , our theorem easily follows from a corollary to Serre's lemma (see corollary to Lemma 1). For  $r=2$ , we have to make a separate argument using a remark of Bass [3] ( $P \oplus A = A^{2n} \Rightarrow P = P' \oplus A$ ).

We consider here only commutative noetherian rings and finitely generated modules. Most of the time we just use the ring  $A$ . For a module  $M$ ,  $\text{hd } M$  denotes its homological dimension.  $\dim A$  denotes the Krull dimension of  $A$ .

The following lemma is basic for what follows. We include a proof here for the sake of completeness.

**LEMMA 1** (Serre [6]). *Let  $A$  be a noetherian ring and  $M$  a left  $A$ -module of homological dimension  $\leq 1$ . Let  $\text{Ext}_A^1(M, A)$  be generated by one element. Then there is an exact sequence*

$$0 \rightarrow A \rightarrow P \rightarrow M \rightarrow 0$$

*with  $P$  projective.*

---

<sup>1)</sup> I am thankful to the Forschungsinstitut, ETH, Zurich, for support when this note was being written.

*Proof.* Let  $\alpha$  generate  $\text{Ext}_A^1(M, A)$  and let  $\alpha$  correspond to the extension  $(\alpha)$ :

$$0 \rightarrow A \rightarrow P \rightarrow M \rightarrow 0$$

Then homing  $(\alpha)$  with  $A$ , we get the exact sequence

$$\text{Hom}(P, A) \rightarrow \text{Hom}(A, A) \xrightarrow{f} \text{Ext}_A^1(M, A) \rightarrow \text{Ext}_A^1(P, A) \rightarrow 0.$$

By extension theory,  $f(1) = \alpha$ . Since  $\text{Ext}_A^1(M, A)$  is generated by  $\alpha$ , we have  $\text{Ext}_A^1(P, A) = 0$ . Since  $\text{hd } M \leq 1$ , by  $(\alpha)$ , it follows that  $\text{hd } P \leq 1$ . Since  $A$  is noetherian,  $\text{hd } P \leq 1$  and  $\text{Ext}_A^1(P, A) = 0$  together imply that  $P$  is projective.

**COROLLARY.** *Let  $A$  be a noetherian ring and  $M$  an  $A$ -module of homological dimension  $\leq 1$ . Let  $\text{Ext}_A(M, A)$  be generated by  $r$  elements. Then there is an exact sequence*

$$0 \rightarrow A^r \rightarrow P \rightarrow M \rightarrow 0$$

with  $P$  projective.

*Proof.* We prove the corollary by induction on  $r$ . For  $r = 1$ , this is precisely Serre's lemma. Assume that the corollary is true for  $r - 1$ . Let  $\alpha_1, \dots, \alpha_r$  generate  $\text{Ext}_A^1(M, A)$ . Let  $\alpha_1$  correspond to the extension  $(\alpha_1)$ :

$$0 \rightarrow A \rightarrow L \xrightarrow{h} M \rightarrow 0$$

Then we get the exact sequence

$$\text{Hom}(L, A) \rightarrow \text{Hom}(A, A) \xrightarrow{f} \text{Ext}_A^1(M, A) \xrightarrow{g} \text{Ext}_A^1(L, A) \rightarrow 0,$$

where  $f(1) = \alpha_1$ . Hence  $\text{Ext}_A^1(L, A)$  is generated by  $r - 1$  elements  $g(\alpha_2), \dots, g(\alpha_r)$ . The exact sequence  $(\alpha_1)$  and the hypothesis that  $\text{hd } M \leq 1$  show that  $\text{hd } L \leq 1$ . Hence by induction hypothesis there is an exact sequence  $I$ :

$$0 \rightarrow A^{r-1} \rightarrow P \xrightarrow{k} L \rightarrow 0$$

with  $P$  projective. We also have the exact sequence

$$0 \rightarrow \text{Ker}(h \circ k) \rightarrow P \xrightarrow{h \circ k} M \rightarrow 0.$$

By the exact sequences  $(\alpha_1)$  and  $I$  it easily follows that  $\text{Ker}(h \circ k) \approx A^r$ . The proof of the corollary is complete.

For a ring  $A$ , we denote  $\text{Max}(A)$  its maximal ideal spectrum and by  $\dim \text{Max}(A)$ , the dimension of  $\text{Max}(A)$ . If  $A$  is an integral domain with quotient field  $K$  and  $M$  an  $A$ -module, we recall that  $\text{rank } M = \dim_K(K \otimes_A M)$ .

**LEMMA 2.** *Let  $A$  be a noetherian domain. Let  $M$  be an  $A$ -module of  $\text{hd } M \leq 1$ . Let  $M$  be an  $A$ -module of rank  $n$ . Let  $\alpha$  be the annihilator of  $\text{Ext}_A^1(M, A)$ . Suppose that*

$\dim \text{Max}(A/\mathfrak{a}) \leq d$ . If  $M$  is locally generated by  $r$  elements, then  $\text{Ext}_A^1(M, A)$  is generated by  $r + d - n$  elements.

*Proof.* By Swan [7], we need only prove that  $\text{Ext}_A^1(M, A)$  is locally generated by  $r - n$  elements. So, we may assume  $A$  is local. Since  $\text{hd } M \leq 1$   $M$  is generated by  $r$  elements and  $\text{rank } M = n$ , we get an exact sequence (since projectives are free over local rings)

$$0 \rightarrow A^{r-n} \rightarrow A^r \rightarrow M \rightarrow 0.$$

Homming this with  $A$ , we see that  $\text{Hom}(A^{r-n}, A) \rightarrow \text{Ext}_A^1(M, A) \rightarrow 0$  is exact. This shows that  $\text{Ext}_A^1(M, A)$  is generated by  $r - n$  elements.

**COROLLARY.** *Let  $A$  be a regular domain of dimension 3. Let  $\mathfrak{a}$  be an unmixed ideal of height 2. If  $\mathfrak{a}$  is locally generated by  $r$  elements, then  $\text{Ext}_A^1(\mathfrak{a}, A)$  is generated by  $r$  elements.*

*Proof.* Using the well known fact that  $\text{depth } M + \text{hd } M = \dim A$  for a regular local ring  $A$ , one easily sees that  $\mathfrak{a}$  is unmixed of height 2 implies  $\text{hd } \mathfrak{a} \leq 1$ . Since  $\text{Ext}_A^1(\mathfrak{a}, A) \approx \text{Ext}_A^2(A/\mathfrak{a}, A)$ , it follows that annihilator  $\mathfrak{b}$  of  $\text{Ext}_A^1(\mathfrak{a}, A)$  contains  $\mathfrak{a}$ . Hence  $\dim A/\mathfrak{b} \leq \dim A/\mathfrak{a} \leq 1$ , since  $\mathfrak{a}$  is unmixed of height 2 and  $\dim A = 3$ . Now the corollary follows from Lemma 2.

Let  $A$  be a ring and  $P$  an  $A$ -module. We recall that  $s \in P$  is *unimodular* if  $s$  generates a free direct summand of  $P$ , isomorphic to  $A$ . For  $x \in \text{Max}(A)$ , we denote by  $s(x)$  the image of  $s$  under the canonical map  $P \rightarrow P/xP$ .

**LEMMA 3.** *Let  $A$  be a noetherian ring of dimension  $\leq 1$  and  $P$  a projective  $A$ -module of rank 2. If  $s_1, s_2, s_3$  generate  $P$ , then there exist  $\lambda_2, \lambda_3 \in A$  such that  $s_1 + \lambda_2 s_2 + \lambda_3 s_3$  is unimodular.*

*Proof.* Let  $\mathfrak{P}_1, \dots, \mathfrak{P}_r$  be the minimal prime ideals of  $A$ . Choose  $\mathfrak{M}_1, \dots, \mathfrak{M}_r \in \text{Max}(A)$  such that  $\mathfrak{M}_i \supset \mathfrak{P}_i$ . Since  $P$  is of rank 2,  $P/(\prod \mathfrak{M}_i)P$  is  $A/\prod \mathfrak{M}_i$  free of rank 2. Using Chinese Remainder Theorem we can find easily  $a, b \in A$  such that if we set  $s'_1 = s_1 + as_3$ ,  $s'_2 = s_2 + bs_3$ , then  $s'_1(\mathfrak{M}_i), s'_2(\mathfrak{M}_i)$  are linearly independent over  $A/\mathfrak{M}_i$ ,  $1 \leq i \leq r$ . Thus we may assume  $s_1, s_2$  are linearly independent at  $\mathfrak{M}_i$ ,  $1 \leq i \leq r$ . Then the set  $T = \{\mathfrak{M} \in \text{Max}(A) \mid s_1(\mathfrak{M}), s_2(\mathfrak{M}) \text{ are linearly dependent}\}$  is a closed set [5, p. 6] which does not contain any irreducible component of  $\text{Max}(A)$ . Since  $\dim A \leq 1$ , it follows that  $T$  is finite. Let  $U = \{\mathfrak{M} \in \text{Max}(A) \mid s_1(\mathfrak{M}) = 0\}$ . Then  $U \subset T$ . Since  $P$  is of constant rank 2 and  $s_1, s_2, s_3$  generate  $P$  it follows that  $s_2(\mathfrak{M}) \neq 0$ ,  $\mathfrak{M} \in U$ . Choose  $f \in A$  such that  $f(\mathfrak{M}) \neq 0$ ,  $\mathfrak{M} \in U$  and  $f(\mathfrak{M}) = 0$ ,  $\mathfrak{M} \in T - U$ . Clearly  $s_1 + fs_2$  is unimodular.

**THEOREM.** *Let  $A$  be a regular integral domain of dimension 3 with  $\tilde{K}_0 A = 0$ . Let  $\mathfrak{a}$  be an ideal unmixed of height 2. Suppose  $\mathfrak{a}$  is locally generated by  $r$  elements. Then  $\mathfrak{a}$  is generated by  $r + 1$  elements.*

We recall that  $\tilde{K}_0 A = 0$  is equivalent to saying that for any finitely generated projective  $A$ -module  $P$ ,  $P \oplus A^m \approx A^n$  for some  $m, n$ .

*Proof of the theorem.* By the corollary to Lemma 2, we see that  $\text{Ext}_A^1(\mathfrak{a}, A)$  is generated by  $r$  elements. Consequently by the corollary to Lemma 1 (since  $\text{hd } \mathfrak{a} \leq 1$ ), we get an exact sequence

$$0 \rightarrow A^r \rightarrow P \rightarrow \mathfrak{a} \rightarrow 0.$$

with  $P$  a projective  $A$ -module of rank  $r+1$ . If  $r \geq 3$ , then  $\text{rank } P \geq 4$ . Since  $\tilde{K}_0 A = 0$  and  $\dim A = 3$ , it follows from Bass' cancellation theorem [2, p. 184, (3.5)] that  $P$  is free. Hence  $\mathfrak{a}$  is generated by  $r+1$  elements. Thus the theorem is proved in case  $r \geq 3$ .

Now we consider the case  $r=2$ . In this case  $P$  is a projective module of rank 3. Again by [2, p. 184, (3.5)],  $P \oplus A \approx A^4$ . Hence by [3],  $P$  admits a free direct summand of rank 1:  $P = P' \oplus A\alpha$ . We have the exact sequence

$$0 \rightarrow A^2 \rightarrow P' \oplus A\alpha \xrightarrow{f} \mathfrak{a} \rightarrow 0. \quad (*)$$

Since  $A$  is a regular ring and  $\mathfrak{a}$  is an ideal of height 2, locally generated by two elements, it follows that  $\mathfrak{a}_{\mathfrak{M}}$  is generated by an  $A_{\mathfrak{M}}$ -sequence of length 2 for any maximal ideal  $\mathfrak{M} \supset \mathfrak{a}$ . Hence  $\mathfrak{a}/\mathfrak{a}^2$  is a locally free  $A/\mathfrak{a}$ -module of rank 2. Tensoring the exact sequence  $(*)$  by  $A/\mathfrak{a}$ , we get the exact sequence

$$A^2 \rightarrow \bar{P}' \oplus \bar{A}\bar{\alpha} \xrightarrow{\bar{f}} \mathfrak{a}/\mathfrak{a}^2 \rightarrow 0$$

where  $\bar{M} = M/\mathfrak{a}M$  for an  $A$ -module  $M$  and for  $x \in M$ ,  $\bar{x}$  denotes residue class of  $x$  modulo  $\mathfrak{a}M$ . Since  $P'$  is stably free, so is the  $\bar{A}$ -module  $\bar{P}'$ . Since dimension of  $\bar{A} \leq 1$ , it follows by [2, p. 170 §2], that  $\bar{P}' \approx \bar{A}^2$ . Let  $\alpha_1, \alpha_2 \in P'$  be such that  $\bar{\alpha}_1, \bar{\alpha}_2$  generate  $\bar{P}'$ . Then  $\bar{f}(\alpha), \bar{f}(\bar{\alpha}_1), \bar{f}(\bar{\alpha}_2)$  generate  $\mathfrak{a}/\mathfrak{a}^2$ . Since  $\mathfrak{a}/\mathfrak{a}^2$  is a projective  $A/\mathfrak{a}$ -module of rank 2, by Lemma 3, there exist  $\bar{a}_1, \bar{a}_2 \in \bar{A}$  such that  $\bar{f}(\bar{a} + \bar{a}_1\bar{\alpha}_1 + \bar{a}_2\bar{\alpha}_2)$  is unimodular in  $\mathfrak{a}/\mathfrak{a}^2$ . Since  $\alpha$  generates a free direct summand of rank 1 with supplement  $P'$ , it follows that  $\alpha + a_1\alpha_1 + a_2\alpha_2$  also generates a free direct summand of rank 1.

The upshot of the above discussion is that we may assume by changing  $\alpha$  to  $\alpha + a_1\alpha_1 + a_2\alpha_2$ , that the class of  $f(\alpha)$  in  $\mathfrak{a}/\mathfrak{a}^2$  is unimodular. Set  $f(\alpha) = a$ . We claim that  $\mathfrak{a}/Aa$  is projective ideal of rank 1 in  $A/aA$ . To show this we observe that since  $\mathfrak{a}/\mathfrak{a} \approx \bar{A}\bar{\alpha} \oplus D$  for some  $D$ , it follows that for any maximal ideal  $\mathfrak{M} \supset \mathfrak{a}$ ,  $a$  can be chosen as one of the two generators for  $\mathfrak{a}_{\mathfrak{M}}$ . Since any two generators of  $\mathfrak{a}_{\mathfrak{M}}$  form an  $A_{\mathfrak{M}}$ -sequence, it follows that  $\mathfrak{a}/Aa$  is locally generated by one element which is even a non-zero-divisor. Thus  $\mathfrak{a}/Aa$  is a projective  $A/aA$ -module of rank 1.

By the exact sequence  $(*)$ , we get (since  $f(\alpha) = a$ ) the exact sequence

$$0 \rightarrow A^2 \rightarrow P' \rightarrow \mathfrak{a}/Aa \rightarrow 0.$$

Tensoring this sequence with  $A/aA$ , we have the exact sequence

$$\frac{A^2}{aA^2} \xrightarrow{g} \frac{P'}{aP'} \rightarrow \mathfrak{a}/Aa \rightarrow 0.$$

Since  $\mathfrak{a}/Aa$  is projective, we have

$$\frac{P'}{aP'} \approx \text{Img} \oplus \mathfrak{a}/Aa$$

Taking  $\overset{2}{\wedge}$  both sides and observing that  $\overset{2}{\wedge} P' \approx A$  (since  $P'$  is stably free) and so  $\overset{2}{\wedge} (P'/aP') \approx A/Aa$ , it follows that

$$\mathfrak{a}/Aa \otimes \text{Img} \approx A/Aa \quad \text{i.e.} \quad \mathfrak{a}/aA \approx (\text{Img})^{-1}.$$

Since  $\text{Img}$  is generated by two elements, it easily follows that  $\mathfrak{a}/aA$  is generated by two elements. Hence  $\mathfrak{a}$  is generated by three elements. The proof of the theorem is completely established.

Since by [2, p. 639, (3.5)]  $\tilde{K}_0(K[x_1, \dots, x_n]) = 0$ , where  $K$  is a field, we have

**COROLLARY.** *Let  $\mathfrak{a} \subset K[x_1, x_2, x_3]$  be an unmixed ideal of height 2. Suppose  $\mathfrak{a}$  is locally generated by  $r$  elements. Then  $\mathfrak{a}$  is generated by  $r+1$  elements.*

A particular case of the Corollary which may be of interest to us

**COROLLARY 2.** *Let  $C$  be a closed affine curve in  $A_3$  which is locally a complete intersection (e.g.  $C$  non-singular), then the ideal  $I(C)$  of  $C$  is generated by three elements.*

*Remark.* (i) One can multiply examples of three dimensional regular rings with trivial  $\tilde{K}_0$  for instance, by using the following well known facts

a)  $\tilde{K}_0 A = 0$ , if  $A$  is a principal ideal domain or a local ring.

b) (Grothendieck), If  $A$  is regular,  $K_0 A \rightarrow K_0 A[x]$  is an isomorphism.

c) If  $A$  is regular, then  $K_0 A \rightarrow K_0 S^{-1} A$  is surjective,  $S$  being multiplicatively closed set.

(ii) (Fossum and Claborn) Let  $K$  be a field of characteristic  $\neq 2$  with  $\sqrt{-1} \in K$  or  $K = \mathbb{R}$  and  $A = K[x_0, x_1, x_2, x_3]$ ,  $\sum x_i^2 = 1$ . Then  $\tilde{K}_0 A = 0$ .

**AN EXAMPLE.** The theorem above is best possible in the sense that there do exist non-singular affine curves  $C$  in  $A_3$  whose prime ideals are *not* generated by two elements. For example, let  $C$  be a complete non-singular curve of genus 2. Let  $\Omega$  denote a divisor in its canonical class. Let  $P \in C$  be such that  $\Omega$  is not linearly equivalent to  $2P$  (such points exist since otherwise the Jacobian variety of  $C$  will be a 2-torsion group!). Consider  $C' = C - \{P\}$ . Then  $C'$  is an affine non-singular curve which can be embedded as a closed set in  $A_3$ . This we can do for instance by considering the complete linear system<sup>2)</sup>  $|5P|$  or by a well known result which says that any non-singular affine curve can be embedded as a closed set in  $A_3$ . Let  $\mathfrak{P}$  be the ideal of  $C'$  in  $A = K[x_1, x_2, x_3]$  ( $K$  algebraically closed). We claim that  $\mathfrak{P}$  is not generated

<sup>2)</sup> I am thankful M. S. Narasimhan for pointing this to me.

by two elements. For if  $\mathfrak{P}$  were generated by two elements, then  $\text{Ext}_A^2(A/\mathfrak{P}, A) \approx A/\mathfrak{P}$  (this one can be seen for example by Koszul-resolution for  $A/\mathfrak{P}$ ). Since by [4],  $\text{Ext}_A^2(A/\mathfrak{P}, A)$  is the module of sections of the canonical line bundle  $\Omega_{C'}$ , it follows that  $\Omega_{C'}$  is trivial. This implies that a canonical divisor  $\Omega \sim nP$ . Since  $\deg \Omega = 2$ , we have  $\Omega \sim 2P$ . Contradiction.

We do not know if the hypothesis  $\tilde{K}_0(A) = 0$  is essential in our theorem.

## REFERENCES

- [1] ABHYANKAR, S., *Algebraic space curves*, University of Montreal Lecture Notes.
- [2] BASS, HYMAN, *Algebraic K-theory*, (Benjamin, New York, 1968).
- [3] — —, *Modules which support non-singular forms*, Journal of Algebra, 13 (2) (1969), 264–252.
- [4] GROTHENDIECK, A., *Théorème de dualité pour les faisceaux algébriques cohérents*, Séminaire Bourbaki, 2e ed., vol. 9 (1956–57).
- [5] SERRE, J. P., *Modules projectifs et espace fibrés à fibre vectorielles*, Séminaire Dubriel-Pisot, 1957/58, no. 23.
- [6] — —, *Sur les modules projectifs*, Séminaire Dubriel-Pisot 1960/1961, no. 2.
- [7] SWAN, R. G., *The number of generators of a module*, Math. Z. 102 (1967) 318–322.

*Forschungsinstitut für Mathematik  
der ETH, Zürich*

Received May 20, 1971.

*Added in Proof:* In the case when  $r \geq 3$ , our result is an easy consequence of Swan [7]. Also, it is not difficult to see that in the statement of the theorem one can drop the hypothesis that  $a$  be of height 2.