

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 47 (1972)

Artikel: Homological Methods and the Third Dimension Subgroup
Autor: Bachmann, F. / Grünenfelder, L.
DOI: <https://doi.org/10.5169/seals-36383>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Homological Methods and the Third Dimension Subgroup

by F. BACHMANN and L. GRÜNENFELDER¹⁾

In this paper we use a certain commutative diagram to derive some well-known and some new results in group theory. Our approach is homological in the sense that we make use of various exact sequences arising in homology of groups.

Let us recall some notation (see [1]): Associated to the lower central series $\{G_n \mid G_1 = G\}$ of a group G is the Lie-ring LG whose underlying abelian group is given by

$$\bigoplus_{i \geq 1} G_i/G_{i+1}.$$

The bracket operation on LG is induced by the commutator in G . There is a natural surjection of graded rings

$$ULG \xrightarrow{\varphi_G} grZG,$$

where ULG denotes the universal envelope of LG and $grZG$ is given as follows: Let JG be the augmentation ideal in ZG ; then $grZG$ is the graded ring associated to the JG -adic filtration of ZG .

Our map φ_G is induced by the Lie-algebra map

$$LG \xrightarrow{i_G} grZG$$

which in degree n is defined by the inclusion of G_n into the n th dimension subgroup $D_n(G) = \{x \in G \mid x - 1 \in JG^n\}$.

In the present investigation we consider φ_G in degree 2. Let $U_n LG$ and $Q_n(G) = JG^n/JG^{n+1}$ be the n th components in ULG and $grZG$ respectively. Our main results are:

THEOREM I. *For any group G there is a commutative diagram (of abelian*

¹⁾ We thank Prof. B. Eckmann for having given us the opportunity to work at the Forschungsinstitut für Mathematik, ETH.

groups) with exact rows

$$\begin{array}{ccccccc} 0 & \rightarrow & G_2/G_3 & \xrightarrow{i_G} & Q_2(G) & \xrightarrow{p} & Q_2(G_{ab}) \rightarrow 0 \\ & & \parallel & & \uparrow \varphi_G & & \uparrow \varphi_{G_{ab}} \\ 0 & \rightarrow & G_2/G_3 & \xrightarrow{j_G} & U_2 LG & \rightarrow & U_2(G_{ab}) \rightarrow 0. \end{array}$$

Here, j_G is the canonical map from LG to ULG , p is the obvious projection, and $G_{ab} = G/G_2$.

COROLLARY 1. $D_3(G) = G_3$ for any group G .

COROLLARY 2. $U_2 LG \xrightarrow{\varphi_G} Q_2(G)$ is bijective for any group G .

THEOREM II. If G is finitely generated the sequence

$$0 \rightarrow G_2/G_3 \xrightarrow{i_G} Q_2(G) \xrightarrow{p} Q_2(G_{ab}) \rightarrow 0$$

is split exact.

§ 1. Diagrams

1. **LEMMA.** For any group G and all $n \geq 2$ there are exact sequences of abelian groups

$$(A_n): H_2(G, \mathbf{Z}) \rightarrow H_2(G/G_n, \mathbf{Z}) \xrightarrow{\beta} G_n/G_{n+1} \rightarrow 0$$

$$(B_n): H_2(G, \mathbf{Z}) \rightarrow H_1(G, JG/JG^n) \rightarrow Q_n(G) \rightarrow 0$$

$$(C_n): 0 \rightarrow G_n/G_{n+1} \xrightarrow{j_G} U_n(LG) \rightarrow U_n(L(G/G_n)) \rightarrow 0$$

Moreover, (C_n) is split exact in the following two cases:

- (a) $n=2$ and G finitely generated
- (b) $n>2$ and G arbitrary.

Proof. (A_n) forms a part of the “5-term-sequence” associated to the group epimorphism $G \rightarrow G/G_n$ (see e.g. [6]).

(B_n) forms a part of the homology sequence corresponding to the G -module sequence

$$0 \rightarrow JG^n \rightarrow JG \rightarrow JG/JG^n \rightarrow 0.$$

We used the identification $H_1(G, JG) \simeq H_2(G, \mathbf{Z})$.

To prove (C_n) we remark that we have an exact sequence of abelian groups

$$0 \rightarrow R \rightarrow LG \rightarrow L(G/G_n) \rightarrow 0,$$

where $R = \bigoplus_{i \geq n} G_i/G_{i+1}$. The kernel V of the algebra morphism

$$ULG \rightarrow UL(G/G_n)$$

is then given by the two-sided ideal in ULG generated by $j_G(R)$. As we are working over \mathbf{Z} , j_G is an injective map ([2]). Therefore the n th component of V coincides with G_n/G_{n+1} . The splitting of (C_n) will be proved in section 6.

2. By naturality the sequence (B_2) yields the following commutative diagram (with α_2 bijective and α_3 surjective):

$$\begin{array}{ccccccc} H_2(G, \mathbf{Z}) & \rightarrow & H_1(G, Q_1(G)) & \rightarrow & Q_2(G) & \rightarrow & 0 \\ \alpha_1 \downarrow & & \alpha_2 \downarrow & & \alpha_3 \downarrow & & \\ H_2(G_{ab}, \mathbf{Z}) & \xrightarrow{\gamma} & H_1(G_{ab}, Q_1(G_{ab})) & \rightarrow & Q_2(G_{ab}) & \rightarrow & 0 \end{array}$$

By a result of Stambach ([6]), γ is a monomorphism. Hence, by applying the ker-coker-sequence and (A_2) , we obtain an isomorphism

$$\partial: \text{Ker } \alpha_3 \cong G_2/G_3.$$

3. Consider now the square

$$\begin{array}{ccc} \text{Ker } \alpha_3 = K & \xrightarrow{\kappa} & Q_2(G) \\ \partial^{-1} \downarrow & * & \downarrow \varphi_G \\ G_2/G_3 & \xrightarrow{j_G} & U_2LG \end{array},$$

where κ is the kernel map. We will prove (see section 5) that $(*)$ commutes. It follows

$$\kappa \cdot \partial^{-1} = \varphi_G \cdot j_G = i_G: G_2/G_3 \rightarrow Q_2(G).$$

Therefore we have

THEOREM I. *For any group G there is a commutative diagram with exact rows*

$$\begin{array}{ccccccc} 0 \rightarrow G_2/G_3 & \xrightarrow{i_G} & Q_2(G) & \rightarrow & Q_2(G_{ab}) & \rightarrow & 0 \\ & \parallel & \varphi_G \uparrow & & \uparrow \varphi_{G_{ab}} & & \\ 0 \rightarrow G_2/G_3 & \xrightarrow{j_G} & U_2LG & \rightarrow & U_2(G_{ab}) & \rightarrow & 0 \end{array}.$$

COROLLARY 1. $D_3(G) = G_3$ for any group G .

COROLLARY 2. $U_2LG \xrightarrow{\varphi_G} Q_2(G)$ is bijective for any group G .

Indeed, Corollary 1 is an immediate consequence of the injectivity of i_G . (See [3], [5] for other proofs of this fact). For the proof of Corollary 2 we observe that $\varphi_{G_{ab}}$ is bijective ([4]) and apply the five lemma.

4. We turn now to the splitting property of the exact sequences in Theorem I. For finite groups, Sandling ([5]) established this fact using the Jennings-Hall-basis of the abelian group JG^2 . The isomorphism of Corollary 2 shows that now it is enough to have a splitting map

$$s: U_2LG \rightarrow G_2/G_3.$$

Such a map exists by lemma 1. Hence we have shown

THEOREM II. *If G is finitely generated the sequence*

$$0 \rightarrow G_2/G_3 \xrightarrow{i_G} Q_2(G) \rightarrow Q_2(G_{ab}) \rightarrow 0$$

is split exact.

§ 2. Proofs

5. To establish the commutativity of the diagram (*) we have to compute the homomorphism $\partial: K \simeq G_2/G_3$ explicitly. If S is the kernel of the canonical surjection $ZG \rightarrow ZG_{ab}$, i.e. S is the ideal $\langle xy - yx \mid x, y \in G \rangle$ in ZG , then

$$K = S + JG^3/JG^3.$$

We use the inhomogeneous bar solution $B(G_{ab})$ to compute ∂ . It is clear from the ker-coker-sequence, that

$$\partial(\overline{xy - yx}) = \beta\bar{z},$$

where \bar{z} is the element of $H_2(G_{ab}, \mathbf{Z})$ represented by the 2-cycle

$$z = 1 \otimes ([\bar{x} \mid \bar{y}] - [\bar{y} \mid \bar{x}])$$

in $\mathbf{Z} \otimes_{G_{ab}} B_2$. The morphism $\beta: H_2(G_{ab}, \mathbf{Z}) \rightarrow G_2/G_3$ in the sequence (A_2) is the composition

$$H_2(G_{ab}, \mathbf{Z}) \xrightarrow{\partial_1} H_1(G_{ab}, JG_{ab}) \xrightarrow{\partial_2} \mathbf{Z} \otimes_{G_{ab}} (G_2)_{ab} \simeq G_2/G_3,$$

where ∂_1 and ∂_2 are the connecting morphisms in the homology sequences of the short

exact sequences

$$JG_{ab} \twoheadrightarrow ZG_{ab} \twoheadrightarrow Z \quad \text{and} \quad (G_2)_{ab} \twoheadrightarrow ZG_{ab} \otimes_G JG \twoheadrightarrow JG_{ab}$$

respectively. For the computation of $\beta\bar{z}$ we remark that $\partial_1\bar{z}$ is represented by the 1-cycle

$$z' = (\bar{x} - 1) \otimes [\bar{y}] - (\bar{y} - 1) \otimes [\bar{x}]$$

in $JG_{ab} \otimes_{G_{ab}} B_1$ and $\partial_2\partial_1\bar{z}$ is represented by the 0-cycle

$$z'' = \overline{[x, y]}$$

in $(G_2)_{ab}$. For the typical element $xy - yx$ of K we therefore get

$$\partial(\overline{xy - yx}) = \beta\bar{z} = \overline{[x, y]}.$$

It is now immediate that the diagram (*) commutes.

6. It remains to show the splitting of the sequence (C_2) . Assume G finitely generated. $G/G_2 = G_{ab}$, being finitely generated, is a finite direct sum of cyclic groups. Let $\bar{x}_1, \dots, \bar{x}_n$ be the generators of its cyclic components. The commutator identity

$$[x^n, y] = [x, y] [[x, y], x^{n-1}] [x^{n-1}, y]$$

in G shows that there is a homomorphism

$$\sigma: (G_{ab} \otimes G_{ab}) \oplus G_2/G_3 \rightarrow G_2/G_3,$$

well defined by

$$\sigma(\bar{x}_i \otimes \bar{x}_j) = \begin{cases} \overline{[x_i, x_j]}, & \text{if } i \leq j \\ 0, & \text{if } i > j \end{cases}$$

on $G_{ab} \otimes G_{ab}$ and by the identity on G_2/G_3 . As

$$\sigma(\bar{x}_i \otimes \bar{x}_j - \bar{x}_j \otimes \bar{x}_i - \overline{[x_i, x_j]}) = 0$$

it factors through U_2LG . The induced homomorphism

$$s: U_2LG \rightarrow G_2/G_3$$

is clearly left inverse to $j_G: G_2/G_3 \rightarrow U_2LG$.

A different proof of Theorem II has been given in a forthcoming paper by G. Losey.

To complete the proof of the lemma in section 1 we assume now \bar{G} arbitrary and

$n > 2$. It is easy to see that

$$L(G/G_n) = \bigoplus_{i=1}^{n-1} G_i/G_{i+1}.$$

Consider the following commutative square (where T is the tensor algebra functor and all arrows are the obvious projections):

$$\begin{array}{ccc} T(LG) & \xrightarrow{\alpha} & T(L(G/G_n)) \\ p \downarrow & & \downarrow q \\ U(LG) & \xrightarrow{\beta} & U(L(G/G_n)) \end{array}$$

The above presentation of $L(G/G_n)$ shows that α has a splitting σ . Clearly we have

$$\sigma(\text{Ker } q) \subset \text{Ker } p.$$

Hence σ lifts to a splitting of β .

REFERENCES

- [1] BACHMANN, F. and GRÜNENFELDER, L., *Ueber Lie-Ringe von Gruppen und ihre universellen Enveloppen*, Comment. Math. Helv. 47 (1972), fasc. 3.
- [2] LAZARD, M., *Sur les algèbres enveloppantes universelles de certaines algèbres de Lie*, Publ. Sci. Univ. Alger (A), I (1954).
- [3] PASSI, I. B. S., *Dimension subgroups*, J. Algebra 9 (1968).
- [4] ———, *Polynomial functors*, Proc. Camb. Phil. Soc. 66 (1969).
- [5] SANDLING, R., *Dimension subgroups over arbitrary coefficient rings*, J. Algebra 21 (1972).
- [6] STAMMBACH, U., *Anwendungen der Homologietheorie der Gruppen auf Zentralreihen und auf Invarianten von Präsentierungen*, Math. Z. 94 (1966).

*Forschungsinstitut für Mathematik
ETH Zürich*

Received August 1972