Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 48 (1973)

Artikel: A Vanishing Theorem in Homological Algebra

Autor: Anderson, Donald W. / Davis, Donald W.

DOI: https://doi.org/10.5169/seals-37157

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A Vanishing Theorem in Homological Algebra

DONALD W. ANDERSON and DONALD W. DAVIS

1. Introduction

In this paper we shall study properties of modules M over the mod 2 Steenrod algebra \mathscr{A} which are useful when using the Adams spectral sequence [1]. Our main result is a vanishing theorem for $\operatorname{Ext}_{\mathscr{A}}(M)$ in terms of $H(M; P_t^s)$, where P_t^s are certain elements of \mathscr{A} whose square is zero so that they act as differentials on M.

THEOREM 1.1. If M is an \mathscr{A} -module and $P_{t_0}^{s_0}$ is the lowest degree P_t^s with s < t such that $H(M; P_t^s) \neq 0$, then $\operatorname{Ext}^{s,t}(M, \mathbb{Z}_2) = 0$ for ds > t + c, where $d = \deg(P_{t_0}^{s_0})$ and c/(d-1) is approximately $t_0 - 2$.

All modules in this paper are understood to be positively graded.

Vanishing theorems of this type were first proved by Adams [2], the case of Theorem 1.1 when d=3. In 1969 Anderson and Mahowald independently proved the case d=6 – that the Adams spectral sequence of an \mathcal{A}_1 -free module vanishes above slope $s/(t-s)=\frac{1}{5}$. This was related to Mahowald's computation of the image of J [8], for if \bar{S} is the sphere spectrum with its zero degree homotopy group killed and P is infinite real projective space, there is a map $P \to \bar{S}$ [8], whose mapping cone has \mathbb{Z}_2 -cohomology free over \mathcal{A}_1 . In this paper we shall employ the techniques which Anderson used to prove this important case.

The effective use of P_t^s -homologies was initiated by Adams and Margolis [4]. We shall use our methods to obtain a proof of the main theorem of [4] which avoids detailed computations using the Steenrod algebra.

THEOREM 1.2. If M is an A-module such that $H(M; P_t^s) = 0$ for all s < t, then M is a free A-module.

The main tool is an exact sequence relating $\operatorname{Ext}_A(\mathbb{Z}_2, \mathbb{Z}_2)$ and $\operatorname{Ext}_B(\mathbb{Z}_2, \mathbb{Z}_2)$, when A is obtained from B by addition of one generator. This is particularly useful if a certain element in $\operatorname{Ext}_A^1(\mathbb{Z}_2, \mathbb{Z}_2)$ can be shown to be nilpotent. We can accomplish this for certain subalgebras of the Steenrod algebra by using a variation on the bar resolution. The Steenrod algebra is then constructed one generator at a time, using the exact sequence to prove a vanishing theorem for the cohomology of each subalgebra.

We wish to thank the referee for many helpful suggestions, and particularly for his very nice proof of Proposition 3.4, which is the one which we give here.

2. The Bar Construction of a Hopf Algebra

If A is a graded algebra over k, and if M, N are two graded left A-modules, we can define bigraded groups $\operatorname{Ext}_A^{s,t}(M,N)$ as follows. Let $\mathbf{P} \to M$ be a projective resoluton of M as an A-module, and let $\operatorname{Hom}_A^s(P_n,N)$ be the A-homorphisms from P_n to N which decrease degree by s. Then $\operatorname{Ext}_A^{s,t}(M,N)$ is the t-dimensional homology of $\operatorname{Hom}_A^s(\mathbf{P},N)$.

Let A, A' be two graded modules over k, and let M, N (resp. M', N') be two modules over A (resp. A'). Then $M \otimes M'$ and $N \otimes N'$ are modules over $A \otimes A'$. If $P \to M$, $P' \to M'$ are projective resolutions, so is $P \otimes P' \to M \otimes M'$. This gives us a pairing

$$\operatorname{Hom}_{\mathbf{A}'}^{\mathbf{s}}(\mathbf{P}, N) \otimes \operatorname{Hom}_{\mathbf{A}'}^{\mathbf{s}'}(\mathbf{P}', N') \to \operatorname{Hom}_{\mathbf{A} \otimes \mathbf{A}'}^{\mathbf{s}+\mathbf{s}'}(\mathbf{P}' \otimes \mathbf{P}, N \otimes N'),$$

and consequently a pairing

$$\operatorname{Ext}_{A}^{s,t}(M,N)\otimes\operatorname{Ext}_{A}^{s',t'}(M',N')\to\operatorname{Ext}_{A\otimes A'}^{s+s',t+t'}(M\otimes M',N\otimes N').$$

If A is a co-commutative connected Hopf algebra over k in the sense of [9], the diagonal $\Delta: A \to A \otimes A$ is a map of algebras, and thus makes every $A \otimes A$ -module into an A-module. Since Ext is contravariant in the ring variable, Δ , together with the pairing above, induces a pairing

$$\operatorname{Ext}_{A}^{s,t}(M,N) \otimes \operatorname{Ext}_{A}^{s',t'}(M',N') \to \operatorname{Ext}_{A}^{s+s',t+t'}(M \otimes M',N \otimes N').$$

This pairing turns $\operatorname{Ext}_A^{**}(k, k)$ into an algebra, where A acts on k through the augmentation $\varepsilon: A \to k$. This product is the product used by Adams in [3].

Notice that if M is any A-module, M is naturally isomorphic to $k \otimes M$. Since $\varepsilon: A \to k$ is A-linear, so is $\varepsilon \otimes M: A \otimes M \to M$. Also, $\Delta: A \to A$ is A-linear, since it is an algebra map. Thus we can form an augmented simplicial A-module $\mathscr{H}_A(M)$ which in degree n is $A^{n+1} \otimes M$, with face and degeneracy maps given by $d_i = A^i \otimes \varepsilon \otimes A^{n-i} \otimes M$, $s_i = A^i \otimes \Delta \otimes A^{n-i} \otimes M$. The degeneracy $s_{-1} = \eta \otimes A^n \otimes M$ is a k-linear contracting homotopy, so we see that augmented simplicial A-module $\mathscr{H}_A(M) \to M$ is acyclic.

To show that the associated chain complex of $\mathcal{H}_A(M)$ is always a free resolution of M, it is necessary and sufficient to show that for any M, $A \otimes M$ is free as an A-module.

Let $_{\varepsilon}M$ be the A-module obtained from the k-module structure of M and the algebra map $\varepsilon: A \to k$. Then $A \otimes (_{\varepsilon}M)$ is certainly free, for A acts on it by the formula $a(a' \otimes m) = (aa') \otimes m$. We define two maps $\varphi: A \otimes_{\varepsilon}M \to A \otimes M$, $\gamma: A \otimes M \to A \otimes_{\varepsilon}M$. The map φ is the following composition:

$$A \otimes {}_{r}M \xrightarrow{A \otimes M} A \otimes A \otimes {}_{r}M \xrightarrow{A \otimes \alpha} A \otimes M$$

where α is the action of A on M. Both maps are A-linear, so φ is A-linear. The map γ is the following composition:

$$A \otimes M \xrightarrow{A \otimes M} A \otimes A \otimes M \xrightarrow{A \otimes \chi \otimes M} A \otimes A \otimes M \xrightarrow{A \otimes \alpha} A \otimes \alpha M$$

where $\chi: A \to A$ is the canonical antiautomorphism. The following proposition follows from the formula $\mu(\chi \otimes A)\Delta = \eta \varepsilon = \mu(A \otimes \chi)\Delta$ and the usual associative laws. The proof is elementary and is left to the reader.

PROPOSITION 2.1. φ and γ are inverse to one another. Thus γ is A-linear.

Notice that (2.1) implies that each term of $\mathcal{H}_A(M)$ is free, so the associated chain complex is a free resolution of M. We call $\mathcal{H}_A(M)$ the Hopf bar construction of A for M. One can use (2.1) to show that $\mathcal{H}_A(M)$ is isomorphic to the usual bar construction, though we shall not show this as we shall not need it.

Let $M^* = \operatorname{Hom}_k(M, k)$. Since there is a natural isomorphism $\operatorname{Hom}_A(A \otimes_{\varepsilon} M, k) \cong M^*$, we have, via φ , a natural isomorphism $\tau : \operatorname{Hom}_A(A \otimes M, k) \to M^*$. Since τ is natural, if $\lambda : M \to N$ is A-linear, $\tau \operatorname{Hom}_A(A \otimes \lambda, k) = \lambda^* \tau$.

PROPOSITION 2.2. Under the isomorphism τ above, $\varepsilon \otimes A \otimes M : A \otimes A \otimes M \to A \otimes M$ $\otimes M$ has (as its dual $(\chi \otimes M^*)\alpha^* : M^* \to A^* \otimes M^*$, where $\alpha : A \otimes M \to M$ is the action of A on M. Also, $\Delta \otimes M : A \otimes M \to A \otimes A \otimes M$ has as its dual $\eta^* \otimes M^* : A^* \otimes M^* \to M^*$. Thus $\operatorname{Hom}_A(\mathscr{H}_A(M), k)$ is isomorphic to a cosimplicial k-module which in codegree n is $(A^*)^n \otimes M^*$, and with coface and cogeneracy maps given by:

$$d^{i} = (A^{*})^{i-1} \otimes \varepsilon^{*} \otimes (A^{*})^{n-i+1} \otimes M^{*} \qquad i > 0$$

$$s^{i} = (A^{*})^{i-1} \otimes \Delta^{*} \otimes (A^{*})^{n-i-1} \qquad i > 0$$

$$s^{0} = \eta^{*} \otimes (A^{*})^{n} \otimes M^{*}$$

$$d^{0} = ((\chi(\Delta^{n+1})^{*}) \otimes (A^{*})^{n} \otimes M^{*}) T^{n+1} ((\mu^{*})^{n} \otimes \alpha^{*})$$

where $\Delta^{n+1}: A \to A^{n+1}$ is the iterated diagonal, and T^{n+1} is the signed interchange

$$T^{n+1}(a_1 \otimes a_1' \otimes \cdots \otimes a_{n+1} \otimes m) = \pm a_1 \otimes a_2 \otimes \cdots \otimes a_{n+1} \otimes a_1' \otimes \cdots \otimes a_n' \otimes m.$$

Given any simplicial abelian group G, Moore has shown how to define an associated chain complex \bar{G} with homology the same as that of G. Let $\bar{G}_n \subset G_n$ be $\operatorname{Ker}(d_1) \cap \ldots \cap \operatorname{Ker}(d_n)$, and let $d: \bar{G}_n \to \bar{G}_{n-1}$ be d_0 . Then $d^2 = 0$ on \bar{G} , and the obvious map $H_n(\bar{G}) \to H_n(G)$ is an isomorphism. Similarly, if G is a cosimplicial abelian group, let $\bar{G}_n = \bar{G}_n/I_m(d^1) + \cdots + I_m(d^n)$, and let $d: \bar{G}_n \to \bar{G}\eta_{+1}$ be d^0 . Then $H^n(\bar{G}) \to H^n(\bar{G})$ will be an isomorphism.

Let $G_n = \operatorname{Hom}_A(\mathcal{H}_A(k)_n, k)$. Then every element of G_n can be represented in the form $(\alpha_1|...|\alpha_n)$ where $\alpha_i \in A^*$ all i. In \overline{G}_n we set equal to 0 all $(\alpha_1|...|\alpha_n)$ with any $\alpha_i \in \operatorname{Im}(\varepsilon^*)$, which is to say $\deg(\alpha_i) = 0$. Notice that

$$d(\alpha_1|\ldots|\alpha_n) = \sum \pm \left(\chi(\alpha'_{1v_1}\ldots\alpha'_{nv_n})|\alpha''_{1v_1}|\ldots|\alpha''_{nv_n}\right) \text{ where } \mu^*(\alpha_i) = \sum \alpha'_{iv_i} \otimes \alpha''_{iv_i}.$$

Since the degreewise tensor product of $\mathcal{H}_A(M)$ with $\mathcal{H}_A(M')$ is $\mathcal{H}_{A\otimes A}(M\otimes M')$, we can use the Alexander-Whitney map to define a chain homotopy equivalence $\mathcal{H}_{A\otimes A}(M\otimes M')\to \mathcal{H}_A(M)\otimes \mathcal{H}_A(M')$. Recall that the Alexander-Whitney map is defined, for x,y of degree n, by $AW(x\otimes y)=\sum (d_i^{n-i+1}x)\otimes (d_0^iy)$. By direct calculation, it is easy to verify that the composite with the diagonal $\mathcal{H}_A(k)\to \mathcal{H}_{A\otimes A}(k)$ gives us a map in degree n which is the product of the maps $A^{n+1}\to A^{i+1}\otimes A^{n-i+1}$ given by $A^i\otimes A\otimes A^{n-i}$ for $0\leqslant i\leqslant n$. If we apply (2.1), we obtain the following.

PROPOSITION 2.3. The product in $\operatorname{Ext}_A(k, k)$, relative to the basis of cochains given before, is given by

$$(\alpha_1 | \dots | \alpha_i) (\alpha_{i+1} | \dots | \alpha_n) = \sum_{i=1}^n (\alpha_1 | \dots | \alpha_{i-1} | \alpha_i) (\alpha'_{i+1, \nu_{i+1}} \dots \alpha'_{n, \nu_n}) | \alpha''_{i+1, \nu_{i+1}} | \dots | \alpha''_{n, \nu_n})$$

where

$$\mu^*(\alpha_j) = \sum \alpha'_{j, \nu_j} \otimes \alpha''_{j, \nu_j}.$$

COROLLARY 2.4. If $\alpha_{i+1},...,\alpha_n$ are primitive in A^* , then for any $\alpha_1,...,\alpha_i$, $(\alpha_1|...|\alpha_i)$ $(\alpha_{i+1}|...|\alpha_n) = (\alpha_1|...|\alpha_n)$. In particular, if α is primitive, $(\alpha)^n = (\alpha|...|\alpha)$.

DEFINITION 2.5. [4] Let P_t^s be the Milnor basis element $Sq(0,...,0,2^s,0,...)$ with the 2^s in the tth position.

PROPOSITION 2.6. [5] The sub-Hopf-algebras of \mathscr{A} correspond bijectively to sequences of nonnegative integers $p_1, \ldots p_i$ (possibly equal to ∞) satisfying $p_i \ge \min(p_j, p_{i-j}-j)$ wherever i > j. The algebra $A(p_1, \ldots)$ corresponding to such a sequence has as \mathbb{Z}_2 -basis all $Sq(r_1, \ldots)$ with $r_i < 2P_i$.

We will find it convenient to use $\zeta_i = \chi(\xi_i)$ as the generators of \mathscr{A}^* . $\Delta(\zeta_i) = \sum_{j=0}^i \zeta_j \otimes \zeta_{i-j}^{2^j}$. The dual of $A(p_1, ...)$ is $\mathbb{Z}_2[\zeta_1, ...]/\zeta_i^{p_i} = 0$.

THEOREM 2.7. Suppose m and n are nonnegative integers and $A = A(p_1, ...)$ is a sub-Hopf-algebra with $p_i \le n$ for all $i \le m$, $p_{m+1} = m+n+2$, and $p_{2m+2} \ge n+1$. Then $\lceil \zeta_{m+1}^{2^{m+n+1}} \rceil$ is a nilpotent element of $\operatorname{Ext}_A(\mathbb{Z}_2, \mathbb{Z}_2)$.

 $\begin{bmatrix} \zeta_{m+1}^{2^{m+n+1}} \end{bmatrix} \text{ is a nilpotent element of } \operatorname{Ext}_{A}(\mathbb{Z}_{2}, \mathbb{Z}_{2}).$ $Proof. \ \Delta(\zeta_{m+1}^{2^{m+n+1}}) = \sum_{j=0}^{m+1} \zeta_{j}^{2^{m+n+1}} \otimes \zeta_{m+1-j}^{2^{m+n+1+j}} = \zeta_{m+1}^{2^{m+n+1}} \otimes 1 + 1 \otimes \zeta_{m+1}^{2^{m+n+1}} \text{ since } \zeta_{j}^{2^{n}} = 0 \text{ for } j \leq m. \text{ Thus in the dual of the Hopf bar resolution } d^{*}([\zeta_{m+1}^{2^{m+n+1}}]) = 0, \text{ and } d^{*}([\zeta_{m+1}^{2^{m+n+1}}]) = 0$

hence $[\zeta_{m+1}^{2^{m+n+1}}] \in \operatorname{Ext}_A^1(\mathbb{Z}_2, \mathbb{Z}_2)$. To show its nilpotence, we shall write $[\zeta_{m+1}^{2^{m+n+1}}]^{2^{m+2}} = [\zeta_{m+1}^{2^{m+n+1}}] \dots |\zeta_{m+1}^{2^{m+n+1}}]$ as a boundary. Let T be the element of $\overline{A^{*2^{m+2}-1}}$ in which ζ_{2m+2}^{2n} occurs in the first $2\eta^{+1}$ positions and ζ_{m+1}^{2m+n+1} occurs in the rest. Since $\Delta(\zeta_{2m+2}^{2n})$ = $\zeta_{2m+2}^{2n} \otimes 1 + 1 \otimes \zeta_{2m+2}^{2n} + \zeta_{m+1}^{2n} \otimes \zeta_{m+1}^{2m+n+1}$, $d^*(T)$ is the sum of all terms having $\chi(\zeta_{m+1}^{i2n})$ ($i \ge 1$) in the first position, ζ_{2m+2}^{2n} in some $2^{m+1} - i$ of the next 2^{m+1} positions, and $\zeta_{m+1}^{2^{m+n+1}}$ in all other positions. Let S be the sum of all $\binom{2^{m+2}-1}{2^{m+1}}$ terms having $\zeta_{2m+2}^{2^n}$ in 2^{m+1} positions and $\zeta_{m+1}^{2^{m+n+1}}$ in $2^{m+1}-1$ positions of an element of $\overline{A^*}^{2^{m+2}-1}$. Then $d^*(S) = [\zeta_{m+1}^{2^{m+n+1}}|...|\zeta_{m+1}^{2^{m+n+1}}]$. To see this note that $[\chi(\zeta_{m+1}^{2^{m+n+1}})|...|\zeta_{m+1}^{2^{m+n+1}}]$ occurs an odd number $((\zeta_{m+1}^{2^{m+2}-1}))$ of times in the sum, and

$$\begin{split} \chi\left(\zeta_{m+1}^{2^{m+n+1}}\right) &= \chi\left(\zeta_{m+1}\right)^{2^{m+n+1}} = \left(\zeta_{m+1} + p\left(\zeta_{1}, ..., \zeta_{m}\right)\right)^{2^{m+n+1}} \\ &= \zeta_{m+1}^{2^{m+n+1}} + p\left(\zeta_{1}^{2^{m+n+1}}, ..., \zeta_{m}^{2^{m+n+1}}\right) = \zeta_{m+1}^{2^{m+n+1}}. \end{split}$$

Any term having $\zeta_{m+1}^{2^{m+n+1}}$ in some fixed $2^{m+1}-1+i\left(0< i< 2^{m+1}\right)$ of the positions occurs in the image under d^* of each term having its $2^{m+1} - 1\zeta_{m+1}^{2^{m+n+1}}$, s in some of these positions; there are $\binom{2^{m+1}-1+i}{2^{m+1}-1}$ such terms, and this is an even number. This also shows $\begin{bmatrix} \xi_{m+n}^{2^{m+n+1}} \end{bmatrix}$ nilpotent, since $\xi_{m+1}^{2^{m+n+1}} = \chi(\zeta_{m+1}^{2^{m+n+1}}) = \zeta_{m+1}^{2^{m+n+1}}$.

3. Techniques that Will be Used to Prove the main Theorems

DEFINITION 3.1. If A is an algebra over k, M an A-module, and $z \in \text{Ext}_{A}(k, k)$, we say z acts nilpotently on $\operatorname{Ext}_A(M, k)$ if every element of $\operatorname{Ext}_A(M, k)$ is annihilated by some power of z.

Our main technique will be to build up the algebras by using

THEOREM 3.2. Suppose B is a subalgebra of A such that A is a free B-module on two generators, 1, x, of degrees 0, n. Then there is a long exact sequence for any Amodule M

$$\rightarrow \operatorname{Ext}_{A}^{s,t+n}(M,k) \rightarrow \operatorname{Ext}_{B}^{s,t+n}(M,k) \rightarrow \operatorname{Ext}_{A}^{s,t}(M,k) \stackrel{[\widehat{x}]}{\rightarrow} \operatorname{Ext}_{A}^{s+1,t+n}(M,k) \rightarrow ,$$

where $[\hat{x}] \in \text{Ext}_A^{1,n}(k,k)$ corresponds to the extension

$$0 \to k \to \operatorname{Hom}_{B}(A, k) \to k \to 0.$$

Moreover, if $\lceil \hat{x} \rceil$ acts nilpotently on $\operatorname{Ext}_{A}(M, k)$, then

- (i) If M is a free B-module, then M is a free A-module.
- (ii) If $\operatorname{Ext}_{R}^{s,t}(M,k)=0$ for ds>t+c with $n \leq d$, then $\operatorname{Ext}_{A}^{s,t}(M,k)=0$ for ds>t++c+n.

Proof. The exact sequence follows from the exact sequence

$$\to \operatorname{Ext}_{A}^{s,t+n}(M,k) \to \operatorname{Ext}_{A}^{s,t+n}(M,\operatorname{Hom}_{B}(A,k)) \to \operatorname{Ext}_{A}^{s,t}(M,k)$$
$$\to \operatorname{Ext}_{A}^{s+1,t+n}(M,k) \to$$

and the change-of-rings isomorphism [6; Proposition 4.1.4] $\operatorname{Ext}_A(M, \operatorname{Hom}_B(A, k)) \approx \operatorname{Ext}_B(M, k)$. The differential is due to [3; Theorem 2.6.1.] If $y \in \operatorname{Ext}_A^{s,t}(M, k)$, then for some $m \ge 0$ $[\hat{x}]^m y$ is a nonzero element of $\operatorname{ker}(\operatorname{Ext}_A^{s+m,t+nm}(M,k) \xrightarrow{[\hat{x}]} \operatorname{Ext}_A^{s+m+1,t+(m+1)n}(M,k))$ and hence pulls back to an element of $\operatorname{Ext}_B^{s+m,t+n(m+1)}(M,k)$.

- (i) If M is B-free, $\operatorname{Ext}_{B}^{s+m, t+nm}(M, k) = 0$ for s+m>0. Thus y cannot exist unless s=0. Thus M is A-free.
- (ii) If the element induced by y is to exist, we must have $t + n(m+1) + c \ge d(s+m)$. Therefore $t + c + n \ge ds + m(d-n) \ge ds$.

Theorem 3.2 will be applied both to the nilpotent elements found in Theorem 2.7 and to elements which will be seen to act nilpotently due to the following result.

PROPOSITION 3.3. If $\operatorname{Ext}_{A}^{s,t}(M,k)=0$ for ds>t+c and $x\in\operatorname{Ext}_{A}^{1,n}(k,k)$ with n< d, then x acts nilpotently on $\operatorname{Ext}_{A}(M,k)$.

Proof. If
$$y \in \text{Ext}_A^{s,t}(M,k)$$
 and $m > (t-ds+c)/(d-n)$, then $x^m y = 0$.

The hypothesis of Proposition 3.3 will often be verified using Proposition 3.4 below, which has as corollary that vanishing theorems of the type which we are seeking hold for modules over exterior algebras on generators of distinct degrees.

PROPOSITION 3.4. Assume that A is free as a left module over the subalgebra B, with a B-base consisting of 1 and other elements a_{α} of degree at least d. Suppose also that M is a B-free module over A and that $M_t=0$ for t < c. Then $\operatorname{Ext}_A^{s,t}(M,k)=0$ for t < ds + c.

Proof. By induction over s. Form the exact sequence

$$0\to K\to A\otimes_B M\to M\to 0\,.$$

Let $\{m_{\mu}\}$ be a *B*-base for *M*; then $A \otimes_B M$ has an *A*-base $\{1 \otimes m_n\}$ and a *B*-base $\{1 \otimes m_{\mu}, a_{\alpha} \otimes m_{\mu}\}$. So *K* has a *B*-base $\{a_{\alpha} \otimes m_n - 1 \otimes a_{\alpha} m_{\mu}\}$. Thus $K_t = 0$ for t < d + c. The inductive hypothesis shows that $\operatorname{Ext}_A^{s-1,t}(V,k) = 0$ for t < d(s-1) + d + c. Since $\operatorname{Ext}_A^{s-1,t}(K,k)$ maps onto $\operatorname{Ext}_A^{s,t}(M,k)$, the latter is zero for t < ds + c.

We shall apply Proposition 3.4 in cases when B is a sub-Hopf-Algebra of A, so that the first hypothesis is fulfilled by [9; Theorem 4.4].

If E is an exterior algebra on generators x_i of distinct degree, and M is an E-module, then the x_i act as differentials on M.

COROLLARY 3.5. If E and M are as above and

$$d = \min \left\{ \deg x_i : H(M; x_i) \neq 0 \right\},\,$$

then $\operatorname{Ext}_{E}^{s,t}(M,k)=0$ for ds>t.

Proof. By [4; Theorem 2.1] both M and E are free over $E[\{x_i: \deg x_i < d\}]$, so the corollary follows from Proposition 3.4.

We shall prove vanishing theorems over finite sub-Hopf-algebras and then conclude vanishing theorems over the entire Steenrod algebra by applying

PROPOSITION 3.6. Suppose A is a sub-Hopf-algebra of \mathscr{A} and $\overline{\mathscr{A}/\mathscr{A}}$ begins in degree n. Suppose $x_1, ..., x_m$ are elements of A such that whenever $H(M; x_i) = 0$, i = 1, ..., m, $\operatorname{Ext}_A^{st}(M, \mathbb{Z}_2) = 0$ for ds > t + c with $d \le n$. If M is an \mathscr{A} -module such that $H(M; x_i) = 0$, i = 1, ..., m, then $\operatorname{Ext}_{\mathscr{A}}^{st}(M, \mathbb{Z}_2) = 0$ for ds > t + c.

Proof. Let $W = \mathcal{A}/\mathcal{A}\bar{A}$. The exact sequences

$$0 \to \bar{W} \otimes \bar{W}^{i-1} \otimes M \to W \otimes \bar{W}^{i-1} \otimes M \xrightarrow{f} \bar{W}^{i-1} \otimes M \to 0$$

(diagonal \mathcal{A} -action, $f(w \otimes p) = \varepsilon(w)p$), yield exact sequences

$$\begin{split} & \to \operatorname{Ext}_{\mathscr{A}}^{s-i,\,t}\big(\bar{W}^i \otimes M,\,\mathbb{Z}_2\big) \to \operatorname{Ext}_{\mathscr{A}}^{s-i+1,\,t}\big(\bar{W}^{i-1} \otimes M,\,\mathbb{Z}_2\big) \\ & \to \operatorname{Ext}_A^{s-i+1,\,t}\big(\bar{W}^{i-1} \otimes M,\,\mathbb{Z}_2\big) \to \,, \end{split}$$

where the Ext_A -group has been obtained because $\operatorname{Ext}_{\mathscr{A}}(w \otimes N, \mathbb{Z}_2) \approx \operatorname{Ext}_A(N, \mathbb{Z}_2)$. Thus an element of $\operatorname{Ext}_{\mathscr{A}}^{s,t}(M, \mathbb{Z}_2)$ must come from an element of $\operatorname{Ext}_A^{s-i,t}(\bar{W}^i \otimes M, \mathbb{Z}_2)$ for some *i*. Since *M* is x_i -acyclic, the same is true of $\bar{W}^i \otimes M$; this is easily seen even if x_i is not primitive. Since $\bar{W}^i \otimes N$ begins in degree $\operatorname{in} \operatorname{Ext}_A^{s-i,t}(\bar{W}^i \otimes M, \mathbb{Z}_2) = 0$ for $d(s-i) > t - \operatorname{in} + c$ and hence makes no contribution to $\operatorname{Ext}_{\mathscr{A}}^{s,t}(M, \mathbb{Z}_2)$ when ds > t + c.

4. Freeness and Vanishing Theorems

In this section we shall prove the freeness theorem 1.2 and a vanishing theorem slightly weaker than Theorem 1.1. The proofs are done similarly, proving them first over an exterior algebra, and then building up the algebra one generator at a time, in an order such that new generators act nilpotently, until we have proved it over a sufficiently large finite subalgebra.

Let $A(O^i, p_{i+1}, p_{i+2},...)$ denote the algebra corresponding by Proposition 2.6 to a sequence beginning with i zeros. Let $B_{i,j,n} = A(O^i, j, n-i, n-i-1,...,1)$ if $i+j \le n+1$.

LEMMA 4.1. Suppose M is an \mathcal{A} -module and suppose we are given integers i, j, n

such that $i + j \le n$. Also, suppose that if j < i + 1, then $H(M; p_t^s) = 0$ for all P_t^s such that s < t, $s + t \le i + j + 1$, $t \ge i + 1$. Then

- (i) If M is free over $B_{i,j,n}$, then M is free over $B_{i,j+1,n}$. (ii) If $\operatorname{Ext}_{B_{i,j,n}}^{st}(M,\mathbb{Z}_2)=0$ for ds>t+c with $d\geqslant (2^{i+1}-1)2^j$, then $\operatorname{Ext}_{B_{i,j+1,n}}^{st}\times$ $\times (M, \mathbb{Z}_2) = 0$ for $ds > t + c + (2^{i+1} - 1)2^j$.

Proof. If $j \ge i+1$, then Theorem 2.7 shows $\begin{bmatrix} \xi_{i+1}^{2^j} \end{bmatrix}$ is nilpotent in $H^*(B_{i,j+1,n})$ and the lemma follows from Theorem 3.2.

If j < i+1, then (since $[P_{t_1}^{s_1}, P_{t_2}^{s_2}] = 0$ whenever $s_1, s_2 < t_1, t_2$) $B_{i, j+1, i+j}$ is an exterior algebra on generators of distinct degree with respect to which M is acyclic. By [4; Theorem 2.1] M is free over $B_{i,j+1,i+j}$. Hence by Proposition 3.4 $\operatorname{Ext}_{B_{i,j+1,n}}^{st}$ $(M, \mathbb{Z}_2) = 0$ for $t < 2^j (2^{i+2} - 1)s$. Thus by Proposition 3.3 $[\xi_{i+1}^{2^j}]$ acts nilpotently on $\operatorname{Ext}_{B_{i,j+1,n}}(M,\mathbb{Z}_2)$. The lemma now follows from Theorem 3.2.

Proof of Theorem 1.2. We shall show that if $A_n = A(n+1, n, ..., 1)$, then M is a free A_n -module. Hence by Proposition 3.4, $\operatorname{Ext}_{\mathscr{A}}^{s,t}(M,\mathbb{Z}_2)=0$ for $2^{n+1}s>t$. Since this is true for all n, $\operatorname{Ext}_{\mathscr{A}}^{s,t}(M,\mathbb{Z}_2)=0$ for s>0. Thus M is \mathscr{A} -free.

If n = 2m, then [4; Theorem 2.1] implies M free over the exterior algebra $B_{m, m+1, 2m}$. Repeated application of Lemma 4.1 shows "M free over $B_{i, 2m+1-i, 2m} = B_{i-1, 0, 2m}$ implies M free over $B_{i-1,2m+1-(i-1),2m}$." Thus by induction M is free over $B_{0,2m+1,2m} = A_{2m}$.

If n=2m-1, M is free over the exterior algebra $B_{m-1, m, 2m-1}$ and by Lemma 4.1 it is free over $B_{m-1,m+1,2m-1}$. By the same induction as above it is free over $B_{0,2m,2m-1}=A_{2m-1}$.

We can now use exactly the same method to prove a vanishing theorem similar to Theorem 1.1 but with slightly larger value for the y-intercept c/(d-1). In the next section we will consider several methods of decreasing the y-intercept. The result proved here will be necessary to prove the stronger result.

THEOREM 4.2. If M is an \mathscr{A} -module and $P_{t_0}^{s_0}$ is the lowest degree P_t^s with s < t such that $H(M; P_t^s) \neq 0$, then $\operatorname{Ext}_{\mathscr{A}}^{st}(M, \mathbb{Z}_2) = 0$ for ds > t + c, where $d = \deg P_{t_0}^{s_0} = 2^{s_0}(2^{t_0} - 1)$ and c is defined as follows. If $s_0 + t_0 = 2m + 1$, then $c = (m-1)2^{2m+2} + 2^{m+1} + m + 2$. If $s_0 + t_0 = 2m$, then $c = (m - 3/2)2^{2m+1} + 2^{m+1} + m + 1$. Thus c/(d-1) is approximately $s_0 + t_0$.

Proof. Suppose $s_0 + t_0 = 2m + 1$. By Corollary 3.5, $\operatorname{Ext}_{B_{m,m+1,2m}}^{st}(M, \mathbb{Z}_2) = 0$ for ds > t. Repeated application of Lemma 4.1 shows "If $\operatorname{Ext}_{B_{t,2m+1-t,2m}}^{st}(M,\mathbb{Z}_{2}) = 0$ for ds > t + c, then $\text{Ext}_{B_{i-1}, 2m+1-(i-1), 2m}^{st}$ $(M, \mathbb{Z}_2) = 0$ for $ds > t + c + (2^i - 1)$ $(2^{2m+2-i}-1)$." Thus by induction $\operatorname{Ext}_{B_{0,2m+1,2m}}^{st}(M,\mathbb{Z}_{2})=0$ for $ds>t+\sum_{i=1}^{m}(2^{i}-1)$ $(2^{2m+2-i}-1)$. The theorem now follows from Proposition 3.6.

If $s_0 + t_0 = 2m$, then Corollary 3.5 shows $\operatorname{Ext}_{B_{m-1, m, 2m-1}}^{st}(M, \mathbb{Z}_2) = 0$ for ds > t. Lemma 4.1 shows $\text{Ext}_{B_{m-1, m+1, 2m-1}}^{st}(M, \mathbb{Z}_2) = 0$ for $ds > t + (2^m - 1)2^m$. Proceeding as above, we find $\operatorname{Ext}_{B_0, 2m, 2m-1}^{st}(M, \mathbb{Z}_2) = 0$ for $ds > t + (2^m - 1)2^m + \sum_{i=1}^{m-1} (2^i - 1)(2^{m2+1-i} - 1)$. The theorem follows from Proposition 3.6.

Although we make no claims about our y-intercept being minimal, the slope is certainly minimal. Indeed,

THEOREM 4.3. If A is an algebra over \mathbb{Z}_2 , and $x \in A_d$ satisfies $x^2 = 0$ and $H(M; x) \neq 0$, and if D > d and C is arbitrary, then there exists s, t such that Ds > t + C and $\operatorname{Ext}_A^{st}(M, \mathbb{Z}_2) \neq 0$.

Proof. If

$$0 \leftarrow M \leftarrow C_0 \leftarrow C_1 \leftarrow C_2 \leftarrow \cdots$$

$$\nwarrow \swarrow \swarrow \swarrow$$

$$K_0 \quad K_1$$

$$\swarrow \qquad \nwarrow$$

$$0 \quad 0 \quad 0$$

is a minimal A-resolution, then the x-homology groups satisfy

$$0 \neq H_i(M) \approx H_{i+d}(K_0) \approx \cdots \approx H_{i+sd}(K_{s-1}).$$

Thus C_s has an element in degree $\leq j+sd$. Thus for all s there is a $t\leq j+sd$ such that $\operatorname{Ext}_A^{st}(M,\mathbb{Z}_2)\neq j$. For large $s,j+sd\leq -c+sD$.

5. Lowering the Vanishing Line

In order to see the problems involved in lowering the vanishing line, let us consider an example. Suppose the first P_t^s with respect to which M has homology is P_3^1 of degree 14 (so M is a free \mathscr{A}_2 -module). Theorem 4.2 showed that $\operatorname{Ext}_{\mathscr{A}}^{st}(M, \mathbb{Z}_2) = 0$ for 14s > t+12+1+2+4+8 by building up the algebras in the order A(0, 2, 2, 1), $A(0, 3, 2, 1), A(1, 3, 2, 1), A(2, 3, 2, 1), A(3, 3, 2, 1), A(4, 3, 2, 1), \mathscr{A}$. Since M is free over A(3, 2, 1), the increments of c when adding the generators P_1^0 , P_1^1 , and P_1^2 may have seemed somewhat unnecessary. Indeed Proposition 3.4 shows that Extst_{A(3, 2, 2, 1)} $(M, \mathbb{Z}_2) = 0$ for 14s > t so that if we can build the algebras in the order A(3, 2, 2, 1), $A(3, 3, 2, 1), A(4, 3, 2, 1), \mathcal{A}$, we would show $\text{Ext}_{\mathcal{A}}^{st}(M, \mathbb{Z}_2) = 0$ for 14s > t + 20. The only problem is the addition of the generator P_2^2 , for $[\zeta_2^4]$ may not be nilpotent in $H^*(A(3, 3, 2, 1))$ despite the fact that it was nilpotent in $H^*(A(0, 3, 2, 1))$. Nor does the argument used before to show that elements act nilpotently apply, for P_2^2 need not lie in a subalgebra of A(3, 3, 2, 1) over which M is free and which contains all of A(3, 3, 2, 1) through $deg(P_2^2)$. However, we are saved by Theorem 4.2, for it gives us a vanishing line for $\operatorname{Ext}_{A(3,3,2,1)}(M,\mathbb{Z}_2)$ which with Proposition 3.3 enables us to conclude that $[\zeta_2^4]$ acts nilpotently. Thus we can prove Theorem 1.1, where the precise value of c is $(t_0-2)2^{t_0+s_0}+2^{s_0+1}$.

Proof. Let $n=t_0+s_0$. Let $B_i=A(n-1, n-2, ..., i+2, i+1, i+1, i, ..., 1)$. By Theorem 1.2 M is free over A(n-1, n-2, ..., 1), and by Proposition 3.4 $\operatorname{Ext}_{B_{s_0}}^{st}(M, \mathbb{Z}_2)=0$ for ds>t. Suppose $\operatorname{Ext}_{B_t}^{st}(M, \mathbb{Z}_2)=0$ for

$$ds > t + c_i (i \ge s_0)$$
 with $c_i = \sum_{j=s_0+1}^i 2^j (2^{n-j} - 1)$.

Since $2^{i+1}(2^{n-i-1}-1)2^{so}(2^{to}-1)$, Theorem 4.2 and Proposition 3.3 apply to show $[\zeta_{n-i-1}^{2^{i+1}}]$ acts nilpotently in $\operatorname{Ext}_{B_{i+1}}(M,\mathbb{Z}_2)$ and Theorem 3.2 then shows $\operatorname{Ext}_{B_{i+1}}^{st}(M,\mathbb{Z}_2)=0$ for $ds>t+c_i+2^{i+1}(2^{n-i-1}-1)$, extending the induction hypothesis. Hence $\operatorname{Ext}_{A(n,n-1,...,1)}^{st}(M,\mathbb{Z}_2)=\operatorname{Ext}_{B_{n-1}}^{st}(M,\mathbb{Z}_2)=0$ for $ds>t+c_{n-1}$ where

$$c_{n-1} = \sum_{j=s_0+1}^{n-1} (2^n - 2^j) = (n - s_0 - 1) 2^n - 2^{s_0+1} (1 + \dots + 2^{n-s_0-2})$$

= $(t_0 - 2) 2^n + 2^{s_0+1}$.

Since $2^n > 2^{so}(2^{to} - 1)$, Proposition 3.6 applies to give the desired result for Ext (M, Z_2) .

For small values of t we can get better results by sacrificing slope for intercept. For example, suppose the first nonacyclic P_t^s is P_5^0 of degree 31. M is free over A(4, 3, 2, 1). Hence by Proposition 3.4 $\operatorname{Ext}_{A(5, 4, 3, 2, 1)}^{st}(M, \mathbb{Z}_2) = 0$ for 16s > t and $\operatorname{Ext}_{A(4, 4, 3, 2, 1)}^{st}(M, \mathbb{Z}_2) = 0$ for 24s > t. Thus by Theorem 2.7 and Theorem 3.2 applied to $\begin{bmatrix} \xi_1^{16} \end{bmatrix}$ and Proposition 3.6, $\operatorname{Ext}_{A(4, 3, 3, 2, 1)}^{st}(M, \mathbb{Z}_2) = 0$ if 16s > t or if 24s > t + 16. Similarly Proposition 3.4 shows $\operatorname{Ext}_{A(4, 3, 3, 2, 1)}^{st}(M, \mathbb{Z}_2) = 0$ for 28s > t and hence two applications of Theorem 3.2 to build it up to A(5, 4, 3, 2, 1) followed by 3.6 shows $\operatorname{Ext}_{A}^{st}(M, \mathbb{Z}_2) = 0$ for 28s > t + 40. Thus we see that for such modules M $\operatorname{Ext}_{A}^{st}(M, \mathbb{Z}_2) = 0$ for $t > \max(16s, 24s - 16, 28s - 40, 30s - 68)$.

REFERENCES

- [1] ADAMS, J. F., On the structure and applications of the Steenrod algebra, Comment. Helv. 32 (1958), 180-214.
- [2] —, A periodicity theorem in homological algebra, Proc. Cambridge Phil. Soc., 62 (1966).
- [3] —, On the nonexistence of elements of Hopf invariant one, Ann. of Math., 72 (1960), 20-104.
- [4] Adams, J. F. and Margolis, H. R., Modules over the Steenrod Algebra, Topology 10 (1971), 271–282.
- [5] —, Sub-Hopf-Algebras of the Steenrod Algebra, preprint.
- [6] CARTAN, H. and EILENBERG, S., Homological algebra (Princeton, 1956).
- [7] LIULEVICIUS, A., The Cohomology of Massey-Peterson Algebras, Math. Z., 105 (1968), 226-256.
- [8] MAHOWALD, M. E., The Order of the Image of the J-Homomorphism, Bull. Amer. Math. Soc. 76 (1970), 1310-1313.
- [9] MILNOR, J. and Moore, J., On the Structure of Hopf Algebras, Ann. of Math. 81 (1965), 211-264.

Dept. of Mathematics, University of California, La Jolla, Calif. 92037, U.S.A.

Received October 27, 1972/May 15, 1973