**Zeitschrift:** Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

**Band:** 48 (1973)

Artikel: Quadratic Spaces with Few Isometries (Quadratic Forms and Linear

Topologies VI)

Autor: Gross, Herbert / Ogg, Erwin

**DOI:** https://doi.org/10.5169/seals-37170

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

# Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 11.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# **Quadratic Spaces with Few Isometries**(Quadratic Forms and Linear Topologies VI)

HERBERT GROSS AND ERWIN OGG

Herrn Professor Dr. Alexander M. Ostrowski zum 80. Geburtstag gewidmet.

#### Introduction

What sort of metric automorphisms do always exist on infinite dimensional quadratic spaces? Clearly, we always have the symmetries about (nondegenerate) hyperplanes, the identity 1 of the space, -1, and of course finite products of these isometries; they form an invariant subgroup  $\Im$  in the full orthogonal group of the space. In the finite dimensional case  $\Im$  is already the full orthogonal group. In the infinite case however,  $\Im$  usually represents only a negligible part of the orthogonal group associated with the space. In this note we shall show that there are quadratic spaces of arbitrarily large dimension whose full orthogonal groups equal  $\Im$ . In  $\S1$  we shall describe how to define such spaces over prescribed (non denumerable) base fields.

The spaces E which we shall investigate below share the following property on subspaces F,

$$F \subset E \& \dim F \geqslant \aleph_0 \to \dim F^1 < \dim E$$
. (\*)

In particular, if such a space E is decomposed orthogonally,  $E = E_1 \oplus E_2$ , then one of the summands  $E_i$  necessarily is of finite dimension. Spaces with such few orthogonal splittings are an extreme counterpart to quadratic spaces admitting orthogonal bases. For subspaces F of spaces wich admit orthogonal bases we invariably have  $\dim E/F^{\perp} = \dim F$  which sharply contrasts (\*). We see in particular that  $\dim E \neq \aleph_0$  for all E satisfying (\*). The construction given in §1 yields spaces which actually satisfy the stronger property on subspaces F,

$$F \subset E \& \dim F \geqslant \aleph_0 \to \dim F^{\perp} \leqslant \aleph_0. \tag{**}$$

The notion which stands in the center of our discussion of spaces with small orthogonal group  $\mathfrak D$  in the sense indicated above  $(\mathfrak D=\mathfrak J)$  is that of a locally algebraic isometry (§2). An isometry T on E is called locally algebraic if T admits for every  $x\in E$  a polynomial  $f_x(T)$  (with coefficients in the base field of E) that annihilates x,  $f_x(T)x=0$ . If  $f_x$  does not depend on x we call T algebraic. Theorem 3 of §2 says that the spaces constructed in §1 admit locally algebraic isometries only; in other words, there are infinite dimensional (\*\*)-spaces E with property  $(\lambda)$ : 'Every

isometry on E is locally algebraic'. By means of somewhat complicated examples one can however show that (\*\*) does not, in general, imply ( $\lambda$ ) (the converse implication is seen not to be true either by Theorem 3 of §2). Spaces with property ( $\lambda$ ) and which, in addition, satisfy (\*) absolutely (i.e. which preserve (\*) under extensions of the base field) are seen to have trivial quotient  $\mathfrak{D}/\mathfrak{J}$  (Corollary 1 of Theorem 3 in §2).

In [3] it is shown that certain spaces constructed in §1 satisfy Witt's cancellation theorem: If  $E = E_1 \oplus E_2 = F_1 \oplus F_2$  are orthogonal decompositions of E with  $E_1$  and  $F_1$  isometric, then  $E_2$  and  $F_2$  must be isometric; a rare thing indeed to happen in the infinite dimensional case.

# Notations.

Generally speaking, forms  $\Phi: E \times E \to k$  are additive in each argument and satisfy  $\Phi(\lambda x, y) = \lambda \Phi(x, y), \ \Phi(x, \lambda y) = \Phi(x, y) \lambda^{\alpha}$  with respect to some fixed involution  $\alpha$  (= antiautomorphism of period 2) of the division ring k. We shall however always assume below that k is commutative. We shall furthermore assume  $\Phi$  to be  $\varepsilon$ -hermitean, i.e.  $\Phi(y, x) = \varepsilon \Phi(x, y)^{\alpha}$  with  $\varepsilon = +1$  (hermitean) or  $\varepsilon = -1$  (antihermitean). If  $\alpha$  is the identity, then k is necessarily commutative and we speak of symmetric and antisymmetric forms respectively. In any case, ' $x \perp y$ ', defined as usual to be ' $\Phi(x, y) = 0$ ', is a symmetric relation.  $E^{\perp}$  is called the *radical* of E (rad E). If rad E=(0) we call  $\Phi$  nondegenerate and – in analogy with algebras – the space  $(E, \Phi)$  semisimple.  $\Phi$  is said to be tracevalued if for every  $x \in E$  there is a  $\xi \in k$  such that  $\Phi(x, x) = \xi + \varepsilon \xi^{\alpha}$ . We shall always assume  $\Phi$  to be tracevalued, a non trivial requirement only when chark k=2([1] §4, No. 2). We shall make use of Witt's theorem in §2 below ([1] §4, No. 3): Let E be a space with a non degenerate form  $\Phi$  which is hermitean or anti-hermitean, and tracevalued if it is hermitean. Then any isometry (=vectorspace isomorphism that preserves  $\Phi$ ) between finite-dimensional subspaces can be extended to a isometric automorphism of E.

Let  $(E, \Phi)$  be an  $\varepsilon$ -hermitean k-vectorspace with respect to the involution  $\alpha$ . Assume that the division ring k' contains k and admits an extension (involution) of  $\alpha$  to k'. We know that the abelian group  $E' = k' \otimes_k E$  may be regarded as a vector-space over k and as a vectorspace over k'. The form  $\Phi': E' \times E' \to k'$ , defined by  $\Phi'(\sum \lambda_i \otimes x_i, \sum \mu_j \otimes y_i) = \sum \lambda_i \Phi(x_i, y_j) \mu_j^{\alpha}$  for  $\lambda_i, \mu_j \in k'$  is  $\varepsilon$ -hermitean. We say that  $\Phi$  satisfies (\*), or (\*\*), absolutely, if the form  $\Phi'$  possesses these properties for all extensions k' of k.  $(E', \Phi')$  is called the k'-ification of  $(E, \Phi)$  or the space obtained from  $(E, \Phi)$  by extending the ring of scalars.

A space  $(E, \Phi)$  is called anisotropic if it contains no isotropic elements, i. e. no vectors  $x \neq 0$  with  $\Phi(x, x) = 0$ .

Unless stated otherwise,  $(E, \Phi)$  will be assumed to be of infinite dimension.

# §1. The Existence of Spaces with Property \*\*

In this short section we shall describe the construction of infinite dimensional spaces  $(E, \Phi)$  where  $\Phi$  is an  $\varepsilon$ -hermitean form satisfying (\*\*) absolutely.

Let  $\alpha$  be an involution of the commutative field k,  $\operatorname{card} k > \aleph_0$ . Let  $X \subset k$  be a maximal subset of algebraically independent elements over the prime field  $k_0$  so that k is an algebraic extension of  $k_0(X)$ . Let  $\varepsilon = +1$  or -1. Since  $\alpha$  is of period 2, there is a subset  $Y \subset X$  with  $\operatorname{card} Y = \operatorname{card} X$  ( $= \operatorname{card} k$ ) and for every  $\eta \in Y$  either  $\varepsilon \eta^{\alpha} = \eta$  or  $\varepsilon \eta^{\alpha} \notin Y$ . Let then  $(e_i)_{i \in I}$  be a basis of a k-vectorspace with  $\operatorname{card} k \geqslant \operatorname{card} I > \aleph_0$ . We define an  $\varepsilon$ -hermitean form  $\Phi$  on  $E \times E$  as follows: Pick an ordering on I. For all  $i < \kappa$  in I set  $\Phi(e_i, e_k) = \varepsilon \Phi(e_k, e_i)^{\alpha} = \eta_{i\kappa} \in Y$  such that all elements  $\eta_{i\kappa}$  ( $i < \kappa$ ) are different. Furthermore  $\Phi(e_i, e_i) = \varepsilon \Phi(e_i, e_i)^{\alpha} \in k$  such that no  $\Phi(e_i, e_i)$  equals a  $\Phi(e_i, e_k)$  with  $i \neq \kappa$ . We assert that  $\Phi$  satisfies (\*\*).

**Proof.** Let U and V be subspaces of E with  $\dim V > \dim U = \aleph_0$ ,  $(u_i)_{i \in \mathbb{N}}$  and  $(v_i)_{i \in J}$  bases of U and V respectively.  $u_i = \sum \alpha_{ik} e_{\kappa}$ ,  $v_i = \sum \beta_{i\kappa} e_{\kappa}$  where the first sum extends over the finite set  $M_i = \{\kappa \in I \mid \alpha_{i\kappa} \neq 0\}$ , the second over the finite set  $N_i = \{\kappa \in I \mid \beta_{i\kappa} \neq 0\}$ . Set  $M = \bigcup_{\mathbb{N}} M_i$ ,  $N = \bigcup_{J} N_i$ . Thus card  $N > \operatorname{card} M = \aleph_0$ . Our assertion is proved if we can exhibit a pair  $u, v \in U \times V$  with  $\Phi(u, v) \neq 0$ . Such a pair is found as follows.

- (i) X contains a denumerable subset A such that  $\{\alpha_{i\kappa} \mid i \in \mathbb{N}, \kappa \in M_i\}$  is contained in the algebraic closure in k of the subfield  $k_0(A)$ .
  - (ii) There is a  $\varrho_0 \in N \setminus M$  such that

$$A \cap \{\Phi(e_{\nu}, e_{\rho_0}), \Phi(e_{\rho_0}, e_{\nu}) \mid \nu \in I \setminus \{\varrho_0\}\} = \emptyset.$$

Let  $\varrho_0 \in N_{\nu_0}$ .

- (iii) X contains a finite subset B such that  $\{\beta_{\nu_0\mu} \mid \mu \in N_{\nu_0}\}$  is contained in the algebraic closure in k of  $k_0(B)$ . Since M is infinite, there is a  $\kappa_0 \in M$  such that  $\Phi(e_{\kappa_0}, e_{\varrho_0}), \Phi(e_{\varrho_0}, e_{\kappa_0}) \notin B$ . Let  $\kappa_0 \in M_{\iota_0}$ .
  - (iv) Notice that  $\kappa_0 \neq \varrho_0$ . If  $\kappa_0 < \varrho_0$  we let

$$C = \{ \Phi(e_{\kappa}, e_{o}) \mid (\kappa, \varrho) \in M_{\iota_{0}} \times N_{\nu_{0}} \setminus \{(\kappa_{0}, \varrho_{0})\} \};$$

if  $\varrho_0 < \kappa_0$  we let

$$C = \{ \Phi(e_{\varrho}, e_{\kappa}) \mid (\varrho, \kappa) \in N_{\nu_0} \times M_{\nu_0} \setminus \{ (\varrho_0, \kappa_0) \} \}.$$

Thus, if  $\kappa_0 < \varrho_0$  we see by (ii), (iii), (iv) that  $\eta_{\kappa_0 \varrho_0} = \Phi(e_{\kappa_0}, e_{\varrho_0}) \notin A \cup B \cup C$ ; similarly, if  $\varrho_0 < \kappa_0$  we have  $\eta_{\varrho_0 \kappa_0} = \Phi(e_{\varrho_0}, e_{\kappa_0}) \notin A \cup B \cup C$ . Thus, if  $k_1$  is the algebraic closure in k of  $k_0 (A \cup B \cup C)$  we see that  $\eta_{\kappa_0 \varrho_0} \notin k_1$  if  $\kappa_0 < \varrho_0$  and  $\eta_{\varrho_0 \kappa_0} \notin k_1$  when  $\varrho_0 < \kappa_0$ . In

the first case we consider

$$\Phi\left(u_{\iota_{0}},\,v_{\nu_{0}}\right) = \sum_{\left(\kappa,\,\rho\right)\neq\left(\kappa_{0},\,\rho_{0}\right)}\alpha_{\iota_{0}\;\kappa}\beta_{\nu_{0}\varrho}\;\phi\left(e_{\kappa},\,e_{\varrho}\right) + \alpha_{\iota_{0}\kappa_{0}}\beta_{\nu_{0}\varrho_{0}}\,\eta_{\kappa_{0}\varrho_{0}}.$$

If we had  $\Phi(u_{i_0}, v_{v_0}) = 0$  then we had a nontrivial linear equation for  $\eta_{\kappa_0 \varrho_0}$  with coefficients in  $k_1$ , so  $\eta_{\kappa_0 \varrho_0} \in k_1$ . If  $\varrho_0 < \kappa_0$  we conclude in the same manner that  $\Phi(v_{v_0}, u_{i_0}) \neq 0$ . Clearly our proof remains valid if we pass to the form  $\Phi'$  on the k'-ification  $E' = k' \otimes_k E$  of E with respect to some overfield k' of k (admitting an extension of  $\alpha$ ). This proves our assertions. We note our result as

THEOREM 1. For  $\varepsilon = +1$  and for  $\varepsilon = -1$  there exist  $\varepsilon$ -hermitean forms  $\Phi$  over any commutative field k with given involution and  $\operatorname{card} k > \aleph_0$  which satisfy (\*\*) absolutely; we may choose the dimension of  $\Phi$  to be  $\operatorname{card} k$ .

We had card  $k \ge \dim E$  for the spaces E in the above construction. We do not know if this is necessarily so for spaces with property (\*\*). It is easy to see that (\*\*) does imply  $(\operatorname{card} k) \aleph_0 \ge \dim E$ . Thus, at least in the special cases where  $\operatorname{card} k$  is a beth (e.g. when  $k = \mathbb{R}$  or  $\mathbb{C}$ ), (\*\*) does imply  $\operatorname{card} k \ge \dim E$ .

THEOREM 2. Let  $k = k_0(X)$  be a purely transcendental extension of  $k_0$  and  $\operatorname{card} X > \aleph_0$ . If – in the notation of the preceding construction –  $\Phi$  is chosen symmetric with  $\Phi(e_i, e_{\kappa}) = \xi_{i\kappa} \in X(i, \kappa \in I \text{ and } \operatorname{card} I > \aleph_0)$  such that  $\xi_{i\kappa} = \xi_{\nu\mu}$  if and only if  $\{i, \kappa\} = \{v, \mu\}$ , then  $\Phi(x, x)$  is a square in k only when x = 0.

This result is proved in [3]; it guarantees the existence of anisotropic forms with property (\*\*) over all fields of a certain type. In the special case where  $k_0$  is assumed orderable the part of theorem 2 ruling out isotropic vectors follows directly from Jacobi's diagonalization formula (for finite spaces). It is clear that after extending the base field  $\Phi$  may admit isotropic vectors. The fact that k is a purely transcendental extension of some  $k_0$  is not however crucial for the existence of an anisotropic  $\Phi$  over k satisfying (\*\*). We give an example of such a form over  $\mathbb{R}$  by specifying a subspace of an infinite separable Hilbertspace  $(H, \Phi)$  over the reals: Note that the collection of all sets M of linearly independent vectors  $x, y, \dots$  with  $\{\Phi(x, y) \mid x, y \in M\}$  algebraically independent over Q is inductively ordered by inclusion. Let  $M_0$  be a maximal element by Zorn's lemma. If card  $M_0 > \aleph_0$ , then the restriction of  $\Phi$  to the span of  $M_0$  satisfies (\*\*) as we have demonstrated above. Assume by way of contradiction that card  $M_0 \leq \aleph_0$ . Let  $(x_i)_{i \in J}$  be the elements of  $M_0$  in some ordering, and let  $A = \{\Phi(x_i, x_j) \mid i, j \in J\}$ . Introduce an orthonormal basis  $(e_i)_{i \in J}$  in the span X of the  $x_i$   $(i \in J)$ ,  $e_i = \sum \alpha_{ij} x_j$  with  $(\alpha_{ij})$  triangular. Then  $(\alpha_{ij})^{-1} = (\beta_{ij})$  is triangular and  $\alpha_{ij}$ ,  $\beta_{ij} \in \overline{\mathbb{Q}(A)}$  (real closure). Since card  $A \leq \aleph_0$  we can pick a family  $(t_i)_{i \in J}$ , the  $t_i$  in  $\mathbb{R}$  and algebraically independent dent over  $\overline{\mathbb{Q}(A)}$  with  $\sum_{i} t_{i}^{2} = t < \infty$ . The closure  $\bar{X}$  of X in H (in the normtopology of  $\Phi$ ) contains a vector  $\overline{x}$  with  $\Phi(x, e_i) = \lambda_i t_i$  for any choice of  $\lambda_i$  with, say,  $0 < \lambda_i < 1$ . We

have  $\Phi(x, x_i) = \sum \beta_{ij} \lambda_j t_j$ . It follows that the set  $\{\Phi(x, x_j) \mid j \in J\}$  is algebraically independent over  $(\mathbb{Q}A)$  for  $\lambda_i$  rational. If we can arrange for  $\Phi(x, x) = \sum (\lambda_i t_i)^2$  to be outside  $\overline{\mathbb{Q}(A \cup (t_i)_J)}$  we have the desired contradiction:  $M_0 \cup \{x\}$  contradicting the maximality of  $M_0$ . Now if J should be finite, then  $\overline{X} = X$  and we may, if necessary, pass from x to a vector x + y with  $y \in X^\perp$  and  $\Phi(y, y) = \alpha - \Phi(x, x)$  and suitably chosen  $\alpha$ . If card  $J = \aleph_0$ , then by varying the rational  $\lambda_i$  in the open unit interval we can arrange for  $\Phi(x, x)$  to be any real number of the open interval [0, t]. Clearly then, there is a choice with  $\Phi(x, x)$  outside the denumerable  $\overline{\mathbb{Q}(A \cup (t_i)_J)}$ . Q.E.D. We can do the same for hermitean forms over a complex Hilbert space. Thus

THEOREM 3. There exist (infinite) positive definite symmetric (hermitean) forms over  $\mathbb{R}(\mathbb{C})$  which satisfy (\*\*) absolutely.

Remark. We briefly indicate how to construct spaces which satisfy (\*\*) but not absolutely so. Let k be nondenumerably infinite. Let  $(f_i)_{i \in I}$ ,  $(g_i)_{i \in I}$  be bases of k-vectorspaces F and G respectively, card  $I = \operatorname{card} k$ . Choose subsets X and Y of k with  $X \cap Y = \emptyset$  and  $X \cup Y$  algebraically independent over the primefield  $k_0$  of k. Define a symmetric bilinear form  $\Phi$  on  $E = F \oplus G$  as follows:  $\Phi(f_i, f_k) = -\Phi(g_i, g_k) = \xi_{ik}$ .  $\Phi(f_i, g_k) = \Phi(f_k, g_i) = \eta_{ik}$  with  $\xi_{ik} \in X$ ,  $\eta_{ik} \in Y$  and  $\xi_{ik} = \xi_{v\mu}$  and  $\eta_{ik} = \eta_{v\mu}$  if and only if  $\{i, k\} = \{v, \mu\}$ . If k is assumed orderable, then the reader proves by the method illustrated above that  $E = F \oplus G$  satisfies (\*\*). However, over the extension  $k(\sqrt{-1})$  E decomposes orthogonally,  $E = H \oplus L$  with H spanned by all  $f_i + \sqrt{-1} \cdot g_i$  ( $i \in I$ ) and L spanned by all  $f_i - \sqrt{-1} \cdot g_i$  ( $i \in I$ ).

# §2. The Orthogonal Group

In this section we study the orthogonal group  $\mathfrak D$  associated with certain infinite dimensional spaces  $(E, \Phi)$  which satisfy (\*). Here  $\Phi$  will always be symmetric or anti-symmetric and tracevalued if it is symmetric.

Consider an isometry T such that there is an orthogonal decomposition  $E = E_0 \oplus E_1$  with  $\dim E_1 < \infty$  and  $T = \pm 1$  on  $E_0$ . Any isometry T with  $\operatorname{Ker}(T-1)$  or  $\operatorname{Ker}(T+1)$  of finite codimension in E admits such an orthogonal decomposition of E. The set  $\mathfrak J$  of all such isometries T is an invariant subgroup of the orthogonal group  $\mathfrak D$  associated with the space E; it contains the subgroup  $\mathfrak J_0$  of index  $\leq 2$  of all T which are the identity on almost all of E. For symmetric  $\Phi$  and  $\operatorname{char} k \neq 2$  [2] gives a detailed account of  $\mathfrak J_0$ ; in that case  $\mathfrak J_0$  is generated by all symmetries about semisimple hyperplanes. We shall show that for prescribed natural n > 1 there are infinite spaces  $(E, \Phi)$  with  $\mathfrak D/\mathfrak J_0$  isomorphic to a product of n copies of  $\mathbb Z_2$  (characteristic not 2).

It is natural to expect, that spaces with few orthogonal splittings in the sense of (\*\*) admit 'few' isometries. A confirmation of this expection is provided by the first two theorems.

THEOREM 1. If  $(E, \Phi)$  satisfies (\*), then every isometry on E is determined modulo a factor from  $\mathfrak{J}_0$  by its action on any subspace of denumerably infinite dimension.

THEOREM 2. If  $(E, \Phi)$  satisfies (\*), and this absolutely so when the base field is not algebraically closed, then every locally algebraic isometry belongs to the group  $\mathfrak{J}$  associated with  $(E, \Phi)$ .

Proof of Theorem 1. Assume first that E is semisimple. For  $\lambda \neq 0$  an element of the basefield k let  $X(\lambda)$  be the eigenspace  $\ker(T-\lambda 1)$  of the isometry T of E.  $X(\lambda) \perp X(\mu)$  if  $\lambda \mu \neq 1$ . Thus we cannot have  $\dim X(\lambda) = \dim E$  unless  $\lambda^2 = 1$  by (\*).  $\operatorname{Im}(\lambda T - 1) \subset X(\lambda)^{\perp}$  and  $\operatorname{Ker}(\lambda T - 1) = X(\lambda^{-1})$  so

$$\dim E/X(\lambda^{-1}) \leqslant \dim X(\lambda)^{\perp}. \tag{1}$$

Assume that for some subspace U of E we have  $T|_{U}=1_{U}$ ,  $\dim U=\aleph_{0}$ . Since T preserves  $\Phi$  we conclude that  $\operatorname{Im}(T-1)$  is contained in  $U^{\perp}$  and thus of dimension smaller than  $\dim E$ . Hence we must have  $\dim X(1)=\dim E$  and therefore  $\dim X(1)^{\perp}<\infty$  by (\*). Hence  $\dim E/X(1)<\infty$  by (1) and therefore  $\dim X(1)^{\perp}\leqslant\dim E/X(1)$  as E is semisimple. Together with (1)  $\dim X(1)^{\perp}=\dim E/X(1)<\infty$ . From this we conclude that there exists a subspace  $H\subset X(1)$  of finite codimension in E with  $E=H\oplus H^{\perp}$ . Since T is the identity on H we have  $T\in\mathfrak{J}_{0}$ . If E is not semisimple, then rad E is of finite dimension. Let  $E_{0}$  be a linear complement of rad E in E. We can find E0 in E1 such that E2 in E3. Since radicals are mapped onto themselves under isometries we must have E3 is determined modulo E4 by its action on E5 Ker E6. Hence the same holds for E7. Q.E.D.

**Proof of Theorem 2.** Case 1: there is a  $\lambda$  with dim  $X(\lambda) = \dim E$ . Hence  $\lambda^2 = 1$  and  $T \in \mathcal{J}$  by Theorem 1.

Case 2:  $\dim X(\lambda^{-1}) < \dim E$  for all  $\lambda \in k \setminus \{0\}$ . Thus  $\dim X(\lambda)^{\perp} = \dim E$  by (1) and so  $\dim X(\lambda) < \infty$  for all  $\lambda \in k \setminus \{0\}$  by (\*). For every member x of a Basis  $\mathscr{B}$  of E we let  $f_x$  be the annihilating polynomial.  $f_x$  splits into linear factors over the algebraic closure k' of  $k, f_x = \prod (Z - \lambda_i)$ . Every linear factor provides an eigenvalue  $\lambda_i \in k'$  of  $T' : E' = k' \otimes E \to E'$ . Since E' satisfies (\*) by the assumptions of the theorem we see that the number l of different  $\lambda_i$  must be less than  $\dim E$ . Hence there are only  $l < \dim E$  different annihilating polynomials  $f_x(x \in \mathscr{B})$ . We conclude that there is at least one  $f_x$  annihilating a subspace  $G \subset E$  of dimension  $\dim G = \dim E$ . Let  $f_x = \prod (Z - \lambda_i)$  be the splitting of this very polynomial. If some of the  $\lambda_i$  equal  $\pm 1$  we let  $G_0$  be the image of G under the map  $\prod_{\lambda_i = \pm 1} (T - \lambda_i 1)$ . We have  $\dim G_0 = \dim G$  in the present case. Let g be the product of the remaining linear factors  $(Z - \lambda)$ . Since  $\dim G_0 = \dim E$  and since g(T) annihilates  $G_0$  and hence also  $G'_0 = k' \otimes G_0$ , we conclude that the dimension of  $\ker(T - \lambda)$  must equal  $\dim E$  for at least one  $\lambda \neq \pm 1$ . This is a contradiction as  $G'_0$  satisfies property (\*).

COROLLARY. If  $(E, \Phi)$  is as in Theorem 2, then the set of all locally algebraic isometries on E is a group. It coincides with the set of all algebraic isometries on E and it is generated by all  $T \in \mathbb{D}$  with E/Ker(T-1) or E/Ker(T+1) finite dimensional; hence it is a normal subgroup of  $\mathbb{D}$ .

LEMMA. Assume that  $E_1, ..., E_n$  all satisfy (\*) and that  $\dim E_i > \dim E_{i+1}$  (i = 1, ..., n-1). If T is any endomorphism of the orthogonal sum  $E = E_1 \oplus \cdots \oplus E_n$  that preserves orthogonality then the  $E_i$  are left almost invariant under T:  $\dim (E_i + T(E_i))/E_i$  is finite for all i.

Proof. Let  $F_1 = E_2 \oplus \cdots \oplus E_n$ . Dim  $E_1 > \dim F_1$  so that there is a subspace  $V_1$  of  $E_1$  with  $T(V_1) \subset E_1$  and dim  $V_1 = \dim E_1$ . By the assumptions of the lemma dim  $T(V_1) = \dim E_1$ . Call  $K_1$  the projection of  $T(F_1)$  onto  $E_1$  (for the decomposition  $E = E_1 \oplus F_1$ ).  $T(V_1) \perp K_1$  hence  $K_1$  and  $(F_1 + T(F_1))/F_1$  are finite dimensional. Setting  $F_2 = E_3 \oplus \cdots \oplus E_n$  we have dim  $E_2 > \dim F_2$ . As  $F_1 + T(F_1)/F_1$  is finite dimensional we conclude that there exists  $V_2 \subset E_2$  with  $T(V_2) \subset E_2$  and dim  $V_2 = \dim E_2$ . It is now clear how the argument may be repeated in order to conclude that there exist spaces  $V_1 \subset E_i$  with  $T(V_i) \subset E_i$  and dim  $V_i = \dim E_i$ . Let then  $K_{ij}$  be the projection of  $T(E_i)$  on  $E_j$ .  $K_{ij} \perp V_j$  for all  $i \neq j$ . Since dim  $T(V_i) = \dim E_i$  by the choice of the  $V_i$  and by the assumptions of the lemma, we conclude that  $K_{ij}$  is finite dimensional for all pairs  $i \neq j$ . This is what the lemma asserts.

We now consider the orthogonal sum of finitely many spaces  $(E_i, \Phi_i)$  of the kind constructed in §1. For the sake of simplicity we choose  $\Phi_i$  symmetric: For i=1, 2, ..., n let  $(e_i^i)_{i \in J(i)}$  be a basis of  $E_i$ ,  $\Phi_i(e_i^i, e_v^i) = \xi_{iv}^i$  where  $\xi_{iv}^i = \xi_{\mu\kappa}^i$  if and only if  $\{i, v\} = \{\mu, \kappa\}$  and where, for every fixed i, the set  $X^i$  of all  $\xi_{iv}^i$   $\{i, v \in J(i)\}$  is algebraically independent over the prime field  $k_0$  of the basefield k. We shall *not* assume that the sets  $X^1, ..., X^n$  are disjoint. For these symmetric spaces we prove

THEOREM 3. Assume that  $\dim E_i > \dim E_{i+1} > \aleph_0$  (i=1,...,n-1). Then every isometry of the orthogonal sum  $E = E_1 \oplus \cdots \oplus E_n$  is locally algebraic.

*Proof.* For the sake of simplicity we omit the superscript 1 when mentioning  $e_i^1$  and  $x_{iv}^1$ ; furthermore let J(1)=J. Let us study the action of T on  $E_1$  for T an isometry of E:

$$Te_i = \sum_{J} \alpha_{i\mu} e_{\mu} + g_i$$
, where  $g_i \in E_2 \oplus \cdots \oplus E_n$ 

By the previous lemma,  $G = k(g_i)_{i \in J}$  is of finite dimension. Let  $Q \subset J$  be such that  $(g_i)_{i \in Q}$  is a basis of G. We introduce the finite sets  $M(\iota) = \{\mu \in J \mid \alpha_{\iota\mu} \neq 0\}$ . Let  $M = \bigcup_{\iota \in J} [M(\iota) \setminus \{\iota\}]$ . We show that M is finite. Assume by way of contradiction that M is infinite. There is a denumerably infinite subset  $S \subset J$  and a map  $\kappa$  that assigns to every  $\iota \in S$  a  $\kappa(\iota) \in J$  with  $\kappa(\iota) \in M(\iota) \setminus \{\iota\}$  and  $\kappa(\iota) \neq \kappa(\nu)$  for all  $\iota \neq \nu$  in S. There is a

subset A of  $X^1 \cup \cdots \cup X^n$  with card  $A \leq \aleph_0$  such that  $\alpha_{\iota\kappa} \in \overline{k_0(A)}$  (the algebraic closure of  $k_0(A)$  in k) for all  $\kappa \in M(\iota)$ ,  $\iota \in S$  and  $\Phi(g_\iota, g_\kappa) \in \overline{k_0(A)}$  for all  $\iota \in S$ ,  $\kappa \in Q$ . Let  $N = \bigcup_{\iota \in S} M(\iota)$ . card  $N = \aleph_0$ . There is a  $\iota \in J$  and for it a  $\mu_0 \in M_{\nu}$  such that  $\mu_0 \notin N \cup S$  and

$$\{\xi_{\iota\mu_0} \in X \mid \iota \in N\} \cap A = \emptyset \tag{2}$$

For  $Te_v = \sum_{\mu \in M} (v) \alpha_{v\mu} \cdot e_{\mu} + \sum_{\kappa \in Q} \beta_{v\kappa} g_{\kappa}$  we have

$$(Te_{i}, Te_{v}) = \xi_{iv} = \alpha_{i\kappa(i)}\alpha_{v\mu_{0}}\xi_{\kappa(i)\mu_{0}} + \sum_{\kappa\mu}\alpha_{i\kappa}\alpha_{v\mu}\xi_{\kappa\mu} + \sum_{\kappa\in Q}\beta_{\mu\kappa}\Phi(g_{i}, g_{\kappa})$$
(3)

The first sum in (3) extends over the set  $[M(\iota) \times M(\nu)] \setminus \{\kappa(\iota), \mu_0\}$ . There is a finite subset B of  $X^1 \cup \cdots \cup X^n$  such that  $\alpha_{\nu\mu} \in \overline{k_0(B)}$  for all  $\mu \in M(\nu)$  and  $\beta_{\nu\kappa} \in \overline{k_0(B)}$  for all  $\kappa \in Q$ . Since S is infinite, there is a  $\sigma \in S$  with  $\xi_{\kappa(\sigma)\mu_0} \notin B$ . As  $\kappa(\sigma) \neq \sigma$  by the choice of the map  $\kappa$  and since  $\mu_0 \neq \sigma$  we have  $\xi_{\sigma\nu} \neq \xi_{\kappa(\sigma)\mu_0}$ . Let  $C = A \cup B \cup \{\xi_{\sigma\nu}, \xi_{\kappa\mu} \mid (\kappa, \mu) \in [M(\sigma) \times M(\nu)] \setminus (\kappa(\sigma), \mu_0)\}$ . By (2) we have  $\xi_{\kappa(\sigma)\mu_0} \notin A$ , hence  $\xi_{\kappa(\sigma)\mu_0} \notin C$ . All quantities in equation (3) equated for  $\iota = \sigma$  are contained in  $\overline{k_0(C)}$  with the exception of  $\xi_{\kappa(\sigma)\mu_0}$ . The coefficient of  $\xi_{\kappa(\sigma)\mu_0}$  in (3) is not zero. Hence we should have  $\xi_{\kappa(\sigma)\mu_0} \in \overline{k_0(C)}$ ; so  $\xi_{\kappa(\sigma)\mu_0}$  is algebraically dependent over C which is a contradiction. We have thus shown that M is finite. G being finite dimensional, there is a subspace  $F_1$  of E, spanned by finitely many  $e_1^i$ ,  $\iota \in J(i)$ ,  $\iota = 1, \ldots, n$  such that  $Te_{\nu}^1 \in k(e_{\nu}^1) + F_1$  for all  $\nu \in J(1)$ . In the same manner we find for  $\iota = 2, \ldots, n$  finite dimensional spaces  $F_i$  such that  $Te_{\nu}^i \in k(e_{\nu}^i) + F_1$ . Set  $F = \sum_{i=1}^n F_i$ . We have  $Te_{\mu}^i \in k(e_{\mu}^i) + F$  for all  $\mu \in J(i)$  and all  $\iota = 1, \ldots, n$ . In particular  $T(F) \subset F$ . Since F is finite dimensional we conclude that T is locally algebraic on all basis vectors  $e_{\mu}$  and hence locally algebraic on each  $\kappa \in C$ . Q.E.D.

Let us look at the proof for one more moment. We have shown that there is a subspace F of E, spanned by finitely many of the basisvectors  $e_i^i$  such that  $Te \in k(e) + F$  for all basis vectors  $e = e_i^i$ . Hence F is the orthogonal sum of its projections onto the summands  $E_i$  in the decomposition  $E = E_1 + \cdots + E_n$ . These projections, say  $G_i$ , are semissimple (as are all spans of collections of basisvectors of our particular bases  $(e_i^i)_{i \in J(i)}, (i=1,\ldots,n)$ ). Therefore  $E_i = G_i \oplus (G_i^1 \cap E_i)$ . Since T(F) = F it follows that the spaces  $G_i^1 \cap E_i$  are left invariant under T. If we extend  $T^{-1}|_F$  to an isometry  $T_0$  on E by letting  $T_0$  act as the identity on  $F^1$  we have  $T_0 \in \mathfrak{J}_0(E)$  and  $T_0 \circ T$  leaves each summand  $E_i$  of E invariant. The restriction of  $T_0 \circ T$  to  $E_i$  is locally algebraic. Hence if char  $k \neq 2$  then we see by Theorem 2 that these restrictions are, up to a factor  $\pm 1$ , a product of finitely many symmetries. We have thus shown that we can find altogether finitely many symmetries S on E such that  $T_0 \circ T \circ \prod S$  acts on each  $E_i$  as  $\mathbf{1}_{E_i}$  or  $-\mathbf{1}_{E_i}$ . Since  $T_0 \in \mathfrak{J}_0(E)$  we obtain the

COROLLARY. Let  $E = E_1 \oplus \cdots \oplus E_n$  be as in Theorem 3 and char  $k \neq 2$ . The

quotient group  $\mathfrak{D}/\mathfrak{J}_0$  of the full orthogonal group of E modulo the invariant subgroup  $\mathfrak{J}_0$  is isomorphic to the direct product of n copies of  $\mathbb{Z}_2$ . In particular, if n=1, then  $\mathfrak{D}/\mathfrak{J}$  is trivial.

# **BIBLIOGRAPHY**

- [1] BOURBAKI, N., Algèbre, ch. 9 ASI 1272 Paris: Hermann (1959).
- [2] GROSS, H., On a Special Group of Isometries of an Infinite Dimensional Vectorspace, Math. Ann. 150 (1963), 285-292.
- [3] SCHNEIDER, U., Über Räume mit wenig orthogonalen Zerlegungen, Diplomarbeit, Math. Institut Univ. Zürich (1972).

Received April 27, 1973