

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 49 (1974)  
  
**Artikel:** Scattering Theory for Elliptic Operators of Arbitrary Order  
**Autor:** Schechter, Martin  
**DOI:** <https://doi.org/10.5169/seals-37981>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Scattering Theory for Elliptic Operators of Arbitrary Order

MARTIN SCHECHTER

## Abstract

We prove the main theorems of scattering theory for selfadjoint elliptic partial differential operators of arbitrary order. Under various hypotheses we show that the wave operators exist and are complete, that the intertwining relations hold, and that the invariance principle holds. One of our main hypotheses is that each lower order coefficient  $q(x)$  satisfies

$$(1+|x|)^\alpha \int_{|x-y|<a} |q(y)| dy \in L^p(E^n)$$

for some  $\alpha \geq 0$ ,  $a > 0$  and for  $p \leq \infty$  such that

$$\alpha > 1 - \frac{2n}{(n+1)p}$$

## 1. Introduction

In this paper we study scattering theory for operators of the form

$$P(D) + Q(x, D) \tag{1.1}$$

in  $E^n$ , where  $P(D)$  is an elliptic operator of order  $2m$  with constant real coefficients, and  $Q(x, D)$  is a symmetric operator of order  $\leq 2m-2$  with variable coefficients. The prototype of (1.1) is the operator

$$-\Delta + q(x), \tag{1.2}$$

which has been studied extensively by many authors. Our results are new even in this case.

One of our results (Theorem 2.7) shows that for  $P(D)$  rotationally invariant the main conclusions of scattering theory hold for (1.1) (and consequently for (1.2)) if each coefficient  $q(x)$  of  $Q(x, D)$  satisfies

$$\sup_x \int_{|x-y|<\delta} |q(y)| |x-y|^{2-n} dy \rightarrow 0 \quad \text{as } \delta \rightarrow 0, \tag{1.3}$$

$$\int_{|x-y|<1} |q(y)| dy \rightarrow 0 \quad \text{as } |x| \rightarrow \infty, \tag{1.4}$$

and there are numbers  $a > 0$ ,  $\alpha \geq 0$  and  $p \leq \infty$  such that

$$\alpha > 1 - \frac{2n}{(n+1)p} \quad (1.5)$$

and

$$(1+|x|)^\alpha \int_{|x-y|<a} |q(y)| dy \in L^p(E^n). \quad (1.6)$$

In particular the wave operators exist and are complete; the intertwining relations and the invariance principle hold (see Section 2). Another result (Theorem 2.6) comes to the same conclusion if (1.6) is replaced by

$$\sup_x \int |q(y)| (1+|x-y|)^{-\beta} dy < +\infty \quad (1.7)$$

holds for some  $\beta < \frac{1}{2}(n-1)$ . Still others (Theorems 2.3 and 2.4) give hypotheses depending on the operator  $P(D)$  and hold even for  $P(D)$  not rotationally invariant.

Kuroda [7] proved these results (and the discreteness of the continuum eigenvalues) under the hypothesis

$$(1+|x|)^{1+\varepsilon} q(x) \in L^\infty \quad (1.8)$$

for some  $\varepsilon > 0$ . He does not require that  $P(D)$  be rotationally invariant. Note that (1.8) implies (1.3)–(1.6). Beals [17] and Birman [18] were the first to study scattering theory for higher order operators. Although his results are weaker, Beals was able to treat operators of the form

$$P(x, D) + q(x), \quad (1.9)$$

where  $P(x, D)$  is a uniformly strongly elliptic operator with smooth coefficients. One of his assumptions is

$$\int_{|x-y|<1} |q(y)| |x-y|^{2m-n} dy \leq C (1+|x|)^{-a} \quad (1.10)$$

for some  $a > n$ . This implies our hypotheses (cf. Theorem 2.7). Birman obtains similar results for  $q(x)$  bounded and

$$(1+|x|)^\alpha q(x) \in L^p \quad (1.11)$$

with  $p=2$  and  $\alpha > \frac{1}{2}n$ . Again this implies (1.3)–(1.6).

The operator (1.2) has received considerable attention in recent years. We quote

only some of the more recent results. Most of them obtained eigenfunctions expansions as well. Ikebe [9] assumed  $q(x) = O(|x|^{-2-\varepsilon})$  as  $|x| \rightarrow \infty$ . Rejto [11] allowed the estimate  $q(x) = O(|x|^{-\varepsilon-4/3})$ , while Kato [13] showed that (1.8) suffices. Kuroda [6] assumed (1.3), (1.4) and (1.11) with

$$\alpha > 1 - p^{-1}. \quad (1.12)$$

Alsholm and Schmidt [10] assumed

$$\int |q(y)| |x-y|^{(1-n)/2} dy \quad (1.13)$$

bounded and tending to 0 as  $|x| \rightarrow \infty$ ,

$$\sup_x \int_{|x-y| < 1} |q(y)|^2 |x-y|^{-\delta} dy < \infty \quad (1.14)$$

for some  $\delta > n-4$ , and (1.11) holds for  $p=2$  and some  $\alpha > 1 - \frac{1}{2}n$ . For  $n=3$ , Kato and Kuroda [1] showed that  $q \in L^{3/2}$  is sufficient, while Simon [13], using a result of Kato [3], extended this to those  $q$  satisfying

$$\int \int |q(x) q(y)| |x-y|^{-2} d^3x d^3y < \infty. \quad (1.15)$$

(He calls this the Rollnik class.) Greifenegger, Jorgens, Weidmann and Winkler [12] assumed

$$\sup_x (1+|x|)^\tau \int_{|x-y| < 1} |q(y)|^2 |x-y|^{-\delta} dy < \infty$$

for some  $\tau > 2$  and  $\delta > n-4$ . It is simple to find potentials which are covered by the present paper but which are not included in any of the results quoted. Moreover, we recapture most of these results.

We follow the method introduced by Kato and Kuroda [1-7], which first proves an abstract theorem in Hilbert space and then applies this to the partial differential operator. The abstract theorems that they proved did not quite suit our needs, although the version given in [7, I] comes fairly close. In Section 3 we present a theorem which differs from that of [7, I] in several crucial respects, but retains most of the basic features. In Section 4 we show how this theorem can be applied to obtain our main results, which are stated in the next section. We would like to thank S. T. Kuroda for making a manuscript of [7] available to us.

In [19] and at a seminar at the Hebrew University given in May, 1973, Agmon



described his results on general eigenfunction expansions for symmetric operators of the form

$$P(D) + \sum_{|\mu| \leq 2m} b_\mu^0(x) D^\mu + \sum_{|\mu| < 2m} b_\mu^1(x) D^\mu.$$

He assumes that the  $b_\mu^0$  satisfy (1.8) and that  $b_\mu^1$  has compact support and is a compact operator from  $H^{2m-|\mu|,2}$  to  $L^2$ . After these lectures, we realized that the results of the present paper hold even when  $Q(x, D)$  is of order  $2m$  as long as its highest order coefficients are bounded and  $P(D) + Q(x, D)$  is bounded from below. All of the proofs apply unchanged.

M. Thompson (Comm. Pure Appl. Math. 25 (1972) 499–532) obtains eigenfunction expansions for higher order elliptic operators. Because his expansions are done in the classical sense, he is required to make assumptions on the operator which are more restrictive than those of Agmon.

## 2. The Main Results

Let  $x = (x_1, \dots, x_n)$  represent a generic point in  $E^n$ , and set  $|x|^2 = x_1^2 + \dots + x_n^2$ .  $D_j = \partial / \partial x_j$ ,  $1 \leq j \leq n$ . For any multi-index  $\mu = (\mu_1, \dots, \mu_n)$  of nonnegative integers, we set  $|\mu| = \mu_1 + \dots + \mu_n$  and  $D^\mu = D_1^{\mu_1} \dots D_n^{\mu_n}$ . We also write  $x^\mu = x_1^{\mu_1} \dots x_n^{\mu_n}$ , and consider a polynomial

$$P(\xi) = \sum_{|\mu| \leq 2m} a_\mu \xi^\mu \quad (2.1)$$

of degree  $2m$  with real coefficients, where  $\xi = (\xi_1, \dots, \xi_n)$ . The partial differential operator corresponding to  $P(\xi)$  is

$$P(D) = \sum_{|\mu| \leq 2m} a_\mu D^\mu. \quad (2.2)$$

Let  $C_0^\infty$  denote the set of complex valued functions in  $C_0^\infty(E^n)$  which have compact supports. Consider  $P(D)$  as an operator on  $L^2 = L^2(E^n)$  with domain  $C_0^\infty$ . It is known that it has a closure  $P_0$  which is self-adjoint (cf., e.g., [14, p. 62]). Our first assumption is

(I) There are positive constants  $a$  and  $K$  such that

$$P(\xi) \geq a|\xi|^{2m} - K, \quad \xi \in E^n. \quad (2.3)$$

Next we consider the variable coefficient operator

$$Q(x, D) = \sum_{j,k=1}^N \bar{P}_j(D) q_{jk}(x) P_k(D), \quad (2.4)$$

where the  $P_k(D)$  are constant coefficient operators of orders  $m_k < m$  and  $\bar{P}(\xi)$  denotes the polynomial whose coefficients are the complex conjugates of those of  $P(\xi)$ . We assume that the coefficients satisfy  $\overline{q_{jk}(x)} = q_{kj}(x)$ . Our hypotheses concerning the coefficients will be expressed in terms of the quantities

$$N_{\alpha, \delta}(h) = \sup_x \int_{|x-y| < \delta} |h(y)|^2 \omega_\alpha(x-y) dy,$$

where

$$\begin{aligned} \omega_\alpha(x) &= |x|^{\alpha-n} & \text{for } \alpha < n \\ &= 1 - \log|x| & \text{for } \alpha = n \\ &= 1 & \text{for } \alpha > n. \end{aligned}$$

We write  $N_\alpha(h) = N_{\alpha,1}(h)$ , and we shall say that  $h \in N_\alpha$  if  $N_\alpha(h) < \infty$ . Our second assumption is

(II) For each  $j$  and  $k$ ,  $q_{jk}(x) = g_{jk}(x) h_{jk}(x)$ , where  $g_{jk} \in N_{2m-2m_j}$  and  $h_{jk} \in N_{2m-2m_k}$ .

Also

$$\int_{|x-y| < 1} |h_{jk}(y)|^2 dy \rightarrow 0 \quad \text{as } |x| \rightarrow \infty. \quad (2.5)$$

Moreover, for those  $j$  satisfying  $2m \leq 2m_j + n$ , we assume

$$N_{2m-2m_j, \delta}(g_{jk}) \rightarrow 0 \quad \text{as } \delta \rightarrow 0, \quad \text{each } k. \quad (2.6)$$

We shall make use of

LEMMA 2.1. *If (I) and (II) hold, then there is a unique self-adjoint operator  $H$  on  $L^2$  such that  $C_0^\infty \subset D(|H|^{1/2})$  and*

$$(Hu, v) = (P(D)u + Q(x, D)u, v), \quad u, v \in C_0^\infty.$$

Next, let  $\Gamma$  be the set of those real  $\lambda$  for which there is a  $\xi \in E^n$  satisfying  $P(\xi) = \lambda$ ,  $\text{grad } P(\xi) = 0$ . It can be shown that  $\Gamma$  consists of a finite number of points (cf. Agmon [19]). We shall prove

LEMMA 2.2. *If  $\lambda$  is not in  $\Gamma$ , then the limit*

$$K_\lambda(x) = \lim_{\varepsilon \rightarrow 0} \int_{|P(\xi) - \lambda| < 1} \frac{\varepsilon e^{ix\xi} d\xi}{(P(\xi) - \lambda)^2 + \varepsilon^2} \quad (2.7)$$

*exists for each  $x \in E^n$  (here  $x\xi = x_1\xi_1 + \cdots + x_n\xi_n$ ). The convergence is uniform on bounded sets and  $K_\lambda(x)$  is infinitely differentiable.*

The proof of this lemma will be given in Section 5. Our final hypothesis will be given in terms of the function (2.7).

(III) For each  $j, k, s, t$  and each  $\lambda \notin \Gamma$

$$\sup_x \int (|g_{jk}(y)|^2 + |h_{st}(y)|^2) |P_j(D) P_t(D) K_\lambda(x-y)| dy < \infty, \quad (2.8)$$

and there are positive constants  $\alpha, C$  such that

$$\begin{aligned} \sup_x \int & (|g_{jk}(y)|^2 + |h_{st}(y)|^2) P_j(D) P_t(D) \\ & \times [K_\lambda(x-y) - K_{\lambda'}(x-y)] dy \leq C |\lambda - \lambda'|^\alpha. \end{aligned} \quad (2.9)$$

One of our main results is

**THEOREM 2.3.** *Under hypotheses (I)–(III) the operator  $H$  given in Lemma 2.1 has the following properties:*

1) *There is a closed set  $e \subset E^1$  of (Lebesgue) measure 0 such that  $H$  restricted to  $E[\sigma(P_0) - (\Gamma \cup e)] L^2$  is absolutely continuous (here  $E(I)$  denotes the spectral measure associated with  $H$ ).*

2) *The absolutely continuous part of  $H$  is unitarily equivalent to  $P_0$ .*

3) *The wave operators*

$$W = s - \lim_{t \rightarrow \pm \infty} e^{itH} e^{-itP_0} \quad (2.10)$$

*exist and are complete.*

4) *The invariance principle holds.*

5) *The intertwining relations*

$$HW_\pm = W_\pm P_0 \quad (2.11)$$

*hold.*

6) *The operator  $S = W_+^* W_-$  is unitary on  $L^2$  and commutes with  $P_0$ .*

Perhaps a few words of explanation are in order. The limit in (2.10) is the strong limit (taken when the operator is applied to each  $u \in L^2$ ). Completeness of the wave operators means that their ranges coincide. This is required for  $S$  to be unitary. The invariance principle means that

$$W_\pm = s - \lim_{t \rightarrow \pm \infty} e^{it\varphi(H)} e^{-it\varphi(P_0)} \quad (2.12)$$

holds for any function  $\varphi(\lambda)$  satisfying

$$\int_0^\infty \left| \int_I e^{-it\varphi(\lambda) - i\xi\lambda} d\lambda \right|^2 d\xi \rightarrow 0 \quad \text{as } t \rightarrow +\infty \quad (2.13)$$

for any compact interval  $I$  contained in  $\sigma(P_0) - e$ . Note that (2.13) holds trivially for  $\varphi(\lambda) = \lambda$ . Further explanations of the conclusions of Theorem 2.3 may be found in Section 3 and in Kato [4, Chapter X].

A slight variation of Theorem 2.3 is given by

**THEOREM 2.4.** *The conclusions of Theorem 2.3 hold if hypothesis (III) is replaced by*

(III') For each  $j, k, s, t$  and each  $\lambda \notin \Gamma$

$$\int \int |g_{jk}(x)|^2 (|g_{st}(y)|^2 + |h_{st}(y)|^2) |P_j(D) P_t(D) K_\lambda(x-y)|^2 dx dy < \infty \quad (2.14)$$

and

$$\int \int |g_{jk}(x)|^2 (|g_{st}(y)|^2 + |h_{st}(y)|^2) |P_j(D) P_t(D) [K_\lambda(x-y) - K_{\lambda'}(x-y)]| dx dy \leq C |\lambda - \lambda'|^\alpha \quad (2.15)$$

for some positive constants  $\alpha, C$ .

Next we restrict our attention to operators which are rotationally invariant. Assume that

$$P(\xi) = p(|\xi|^2), \quad (2.16)$$

where  $p(t)$  is a polynomial of degree  $m$  in one variable. In this case we can compute the functions (2.7) explicitly. In fact we have

**THEOREM 2.5.** *If (2.16) holds and  $\lambda \notin \Gamma$ , then*

$$K_\lambda(x) = \pi^2 (2\pi/|x|)^\gamma \sum_{p(t_k^2) = \lambda} t_k^\gamma J_\gamma(t_k |x|) / p'(t_k^2), \quad (2.17)$$

where  $\gamma = \frac{1}{2}n - 1$  and the  $t_k$  are the positive roots of  $p(t^2) = \lambda$  (here  $p'$  denotes the derivative of  $p$ ). Moreover, one has

$$|D^\mu K_\lambda(x)| \leq C_{\lambda, \mu} (1 + |x|)^{(1-n)/2} \quad (2.18)$$

for each  $\mu$  and

$$\frac{|D^\mu[K_\lambda(x) - K_{\lambda'}(x)]|}{(1+|x|)^{\alpha+(1-n)/2}} \leq C_{\lambda, \mu, \alpha} |\lambda - \lambda'|^\alpha \quad (2.19)$$

for each  $\alpha > 0$  and  $\mu$ .

Using the estimates (2.18) and (2.19), we have

**THEOREM 2.6.** *Suppose  $P(\xi)$  is rotationally invariant and that (I) and (II) hold. Assume also that for each  $j$  and  $k$  there is a  $\beta < \frac{1}{2}(n-1)$  such that*

$$\sup_x \int (|g_{jk}(y)|^2 + |h_{jk}(y)|^2) (1+|x-y|)^{-\beta} dy < \infty. \quad (2.20)$$

*Then all of the conclusions of Theorem 2.3 hold.*

**THEOREM 2.7.** *Suppose  $P(\xi)$  is rotationally invariant and that (I) and (II) hold. Assume also that for each  $j$  and  $k$  there are numbers  $\delta > 0$ ,  $\alpha \geq 0$  and  $p \leq \infty$  such that*

$$\alpha > 1 - \frac{2n}{(n+1)p}$$

and

$$(1+|x|)^\alpha \int_{|x-y| < \delta} (|g_{jk}(y)|^2 + |h_{jk}(y)|^2) dy \in L^p.$$

*Then all of the conclusions of Theorem 2.3 hold.*

### 3. An Abstract Theorem

We now give a theorem in Hilbert space which we use in proving our results in Section 2. It is a variation of the Kato-Kuroda approach (cf. [1-7]), and we found it most useful for the applications at hand. We believe that it is of interest in its own right. Our results are similar to Theorems 3.11-3.13 of [7, I], but there are important differences. In particular, we were unable to use the results of Kuroda to obtain the theorems of Section 2.

First we discuss notations and conventions. For a Hilbert space  $\mathcal{H}$ , the norm and scalar product will be denoted by  $\|\cdot\|_{\mathcal{H}}$  and  $(\cdot, \cdot)_{\mathcal{H}}$ , respectively.  $B(\mathcal{H}, \mathcal{K})$  will denote the set of bounded linear operators from  $\mathcal{H}$  to  $\mathcal{K}$ . We shall write  $B(\mathcal{H})$  for  $B(\mathcal{H}, \mathcal{H})$ . All operators will be densely defined. The domain and range of an operator will be denoted by  $D(A)$  and  $R(A)$ , respectively. Its resolvent and spectrum sets will be denoted by  $\varrho(A)$  and  $\sigma(A)$ , respectively. The closure of a closable operator will be

denoted by  $[A]^a$ . All subsets of the real line will be Borel sets. For each such set  $I$ ,  $\chi_I$  will denote its characteristic function. The Hilbert space of all strongly measurable  $\mathcal{H}$ -valued functions with square integrable norms over  $I$  will be denoted by  $L^2(I, \mathcal{H})$ .  $I \subset \subset J$  will mean that the closure  $\bar{I}$  of  $I$  is a compact subset of  $J$ .

A bilinear form  $a(u, v)$  with domain  $D(a)$  is a complex valued functional on  $D(a) \times D(a)$  which is linear in  $u$  and conjugate linear in  $v$ .  $D(a)$  is to be a subspace of a Hilbert space (cf. [22, Chapter XII]). An operator  $A$  is *associated* with a bilinear form  $a(u, v)$  if the statements  $u \in D(A)$  and  $Au = f$  are equivalent to  $u \in D(a)$  and  $a(u, v) = (f, v)$  for all  $v \in D(a)$ . Every self-adjoint operator is associated with a bilinear form.

Our assumptions are

1.  $H$  is a self-adjoint operator on a Hilbert space  $\mathcal{H}$ . Let  $\{E(\lambda)\}$  denote its spectral family and  $\{E(I)\}$  its spectral measure. For  $\zeta \in \rho(\mathcal{H})$ , set  $R(\zeta) = (\zeta - H)^{-1}$ . All norms and scalar products without subscripts will be those of  $H$ .

2. There are an open subset  $\wedge$  of the real line, a Hilbert space  $\mathcal{C}$  and a unitary operator  $F$  from  $E(\wedge) \mathcal{H}$  onto  $L^2(\wedge, \mathcal{C})$  such that

$$FE(I)F^{-1} = \chi_I, \quad I \subset \wedge. \quad (3.1)$$

3. There are a Hilbert space  $\mathcal{K}$  and closed operators  $A, B$  from  $H$  to  $K$  such that

$$\tilde{D} = D(|H|^{1/2}) \subset D(A) \cap D(B). \quad (3.2)$$

4.  $A$  is injective.

5. There is a  $\zeta_0 \in \rho(H)$  such that  $BR(\zeta)[R(\zeta_0)A^*]^a$  is a compact operator on  $\mathcal{K}$  for each  $\zeta \in \rho(H)$ .

6. There are continuous functions  $Q_{\pm}(\lambda)$  from  $\wedge$  to  $B(\mathcal{K})$  such that  $[BR(\zeta)A^*]^a$  converges in norm to  $Q_{\pm}(\lambda)$  as  $\zeta \rightarrow \lambda$  for each  $\lambda \in \wedge$ .

7. There is a function  $M(\lambda)$  on  $\wedge$  with values in  $B(\mathcal{K})$  which is integrable over any compact subset of  $\wedge$  and such that

$$\frac{d}{d\lambda}(E(\lambda)A^*u, A^*v) = (M(\lambda)u, v)_{\mathcal{K}} \quad \text{a.e. } u, v \in D(A^*) \quad (3.3)$$

8. There is a self-adjoint operator  $H_1$  on  $\mathcal{H}$  such that  $D(|H_1|^{1/2}) = \tilde{D}$  and its associated bilinear form satisfies

$$h_1(u, v) = h(u, v) + (Bu, Av)_{\mathcal{K}}, \quad u, v \in \tilde{D}, \quad (3.4)$$

where  $h(u, v)$  is the bilinear form associated with  $H$ . We let  $\{E_1(\lambda)\}$  and  $\{E_1(I)\}$  denote the spectral family and measure of  $H_1$ , respectively. For  $\zeta \in \rho(H_1)$  we put  $R_1(\zeta) = (\zeta - H_1)^{-1}$ .

The main theorem of this section is:

**THEOREM 3.1.** *Under Assumptions 1–8, there is a closed set  $e \subset \Lambda$  of Lebesgue measure 0 and unitary operators  $W_{\pm}$  from  $E(\Lambda) \mathcal{H}$  onto  $E_1(\Lambda - e) \mathcal{H}$  such that:*

- (a)  $H_1 W_{\pm} = W_{\pm} H$  on  $E(\Lambda) \mathcal{H}$
- (b)  $S = W_+^* W_-$  is unitary on  $E(\Lambda) \mathcal{H}$  and commutes with  $H$ .
- (c)  $W_{\pm} u = \lim_{t \rightarrow \pm \infty} e^{itH_1} e^{-itH} u, \quad u \in E(\Lambda) \mathcal{H}$
- (d) *The invariance principle holds.*

We give the proof by means of a series of lemmas. Lemmas 3.2–3.5 were proved in [7, I]. We make the following definitions.

$$\begin{aligned}
 Q(\zeta) &= [BR(\zeta) A^*]^a, \quad G(\zeta) = I + Q(\zeta). \\
 Q_1(\zeta) &= [BR_1(\zeta) A^*]^a, \quad G_1(\zeta) = I - Q_1(\zeta). \\
 \Pi^{\pm} &= \{\zeta \mid \operatorname{Im} \zeta \gtrless 0\} \\
 \Pi_I^{\pm} &= \Pi^{\pm} \cup I \\
 G_{\pm}(\zeta) &= G(\zeta) \quad \text{in } \Pi^{\pm} \\
 G_{\pm}(\lambda) &= I + Q_{\pm}(\lambda), \quad \lambda \in \Lambda. \\
 e_{\pm} &= \{\lambda \in \Lambda \mid G_{\pm}(\lambda) \text{ is not injective}\} \\
 e &= e_+ \cup e_- \\
 G_{1\pm}(\zeta) &= G_1(\zeta) \quad \text{in } \Pi^{\pm} \\
 G_{1\pm}(\lambda) &= G_{\pm}(\lambda)^{-1} \quad \text{in } \Lambda - e.
 \end{aligned}$$

**LEMMA 3.2.** *The set  $e$  is closed and of Lebesgue measure 0.*

**LEMMA 3.3.** *For  $\zeta \in \varrho(H) \cap \varrho(H_1)$*

$$BR(\zeta) = G(\zeta) BR_1(\zeta) \tag{3.5}$$

and

$$[R(\delta) A^*]^a = [R_1(\zeta) A^*]^a G(\zeta). \tag{3.6}$$

**LEMMA 3.4.**  *$G_{\pm}(\zeta)$  is continuous in  $\Pi_{\Lambda}^{\pm}$ , and  $G_{1\pm}(\zeta)$  is continuous in  $\Pi_{\Lambda - e}^{\pm}$ . Also*

$$G_{1\pm}(\zeta) = G_{\pm}(\zeta)^{-1} \quad \text{in } \Pi_{\Lambda - e}^{\pm}. \tag{3.7}$$

**LEMMA 3.5.** *If  $I$  is bounded, then  $[E(I) A^*]^a$  is in  $B(\mathcal{H}, \mathcal{H})$ .*

**LEMMA 3.6.** *Set*

$$T = T_{\lambda} = FE(\Lambda) A^*.$$

Then  $T$  maps  $D(A^*)$  into  $L^2(\wedge, \mathcal{C})$ . Also

$$\frac{d}{d\lambda} (E(\lambda) A^* u, A^* v) = (T_\lambda u, T_\lambda v)_\mathcal{C} \quad \text{a.e., } u, v \in D(A^*). \quad (3.8)$$

*Proof.* Let  $I$  be any subset of  $\wedge$ . Then

$$\begin{aligned} (E(I) A^* u, A^* v) &= (E(\wedge) E(I) E(\wedge) A^* u, E(\wedge) A^* v) \\ &= \int_{\wedge} (FE(I) E(\wedge) A^* u, FE(\wedge) A^* v)_\mathcal{C} d\lambda \\ &= \int_I \chi_I(\lambda) (FE(\wedge) A^* u, FE(\wedge) A^* v)_\mathcal{C} d\lambda \\ &= \int_I (Tu, Tv)_\mathcal{C} d\lambda. \end{aligned} \quad (3.9)$$

Since this is true for any subset  $I$ , we get (2.8).

LEMMA 3.7.  $R(T)$  is dense in  $L^2(\wedge, \mathcal{C})$ .

*Proof.* Let  $g(\lambda)$  be any function in  $L^2(\wedge, \mathcal{C})$ . By Assumption 2 there is a  $w$  in  $\mathcal{H}$  such that  $g = Fw$ . Since  $A$  is injective (Assumption 4),  $R(A^*)$  is dense in  $\mathcal{H}$ . Thus there is a sequence  $\{u_k\}$  of elements in  $D(A^*)$  such that  $A^* u_k \rightarrow w$  in  $\mathcal{H}$ . Thus  $E(\wedge) A^* u_k \rightarrow E(\wedge) w$  in  $\mathcal{H}$  and  $FE(\wedge) A^* u_k \rightarrow FE(\wedge) w$  in  $L^2(\wedge, \mathcal{C})$ .  $\square$

LEMMA 3.8. For each bounded subset  $I$  of  $\wedge$  and each bounded Borel measurable function  $f(\lambda)$ ,

$$[A f(H) E(I) A^*]^a = \int_I f(\lambda) M(\lambda) d\lambda. \quad (3.10)$$

*Proof.* We have

$$\begin{aligned} (f(H) E(I) A^* u, A^* v) &= \int_I f(\lambda) \frac{d}{d\lambda} (E(\lambda) A^* u, A^* v) d\lambda \\ &= \int_I f(\lambda) (M(\lambda) u, v)_\mathcal{X} d\lambda, \quad u, v \in D(A^*). \end{aligned}$$

Since  $D(A^*)$  is dense in  $\mathcal{X}$ , we get (2.10).

LEMMA 3.9.

$$(M(\lambda) u, v)_\mathcal{X} = (T_\lambda u, T_\lambda v)_\mathcal{C} \quad \text{a.e., } u, v \in D(A^*).$$



*Proof.* Apply Lemma 3.6 to Assumption 7.  $\square$

Set

$$\delta_\varepsilon(\mu) = \frac{\varepsilon}{\pi} (\mu^2 + \varepsilon^2)^{-1}$$

Thus

$$\delta_\varepsilon(H - \lambda) = \frac{\varepsilon}{\pi} R(\lambda + i\varepsilon) R(\lambda - i\varepsilon).$$

LEMMA 3.10. *For almost all  $\lambda \in \Lambda$ ,*

$$[A\delta_\varepsilon(H - \lambda) A^*]^a \rightarrow M(\lambda)$$

*in norm as  $\varepsilon \rightarrow 0$ .*

*Proof.* Let  $I$  be any compact interval in  $\Lambda$ . Then for  $\lambda \in I$

$$\begin{aligned} [A\delta_\varepsilon(H - \lambda) A^*]^a &= [A\delta_\varepsilon(H - \lambda) E(I) A^*]^a \\ &\quad + [A\delta_\varepsilon(H - \lambda) E(R - I) A^*]^a. \end{aligned}$$

The first term on the right equals

$$\frac{\varepsilon}{\pi} \int_I \frac{1}{(\mu - \lambda)^2 + \varepsilon^2} M(\mu) d\mu$$

by (3.10). This converges a.e. in norm to  $M(\lambda)$ . The second term equals

$$\frac{\varepsilon}{\pi} AR(\lambda + i\varepsilon) E(R - I) [R(\lambda - i\varepsilon) E(R - I) A^*]^a.$$

Since  $AR(\lambda + i\varepsilon)$  is bounded on  $E(R - I)$  for  $\lambda$  an interior point of  $I$ , this term tends to 0 in norm as  $\varepsilon \rightarrow 0$  for such  $\lambda$ .  $\square$

LEMMA 3.11. *For  $u, v \in D(A^*)$*

$$(\delta_\varepsilon(H_a - \lambda) A^*u, A^*v) = ([A\delta_\varepsilon(H - \lambda) A^*]^a G_{1\pm}(\zeta) u, G_{1\pm}(\zeta) v)_{\mathcal{H}} \quad (3.11)$$

*Proof.* The left hand side of (3.11) equals

$$\frac{\varepsilon}{\pi} (R_1(\zeta) A^*u, R_1(\zeta) A^*v),$$

where  $\zeta = \lambda + i\varepsilon$ . By (3.6) and (3.7) this equals

$$\frac{\varepsilon}{\pi} ([R(\zeta) A^*]^a G_{1\pm}(\zeta) u, [R(\zeta) A^*]^a G_{1\pm}(\zeta) v). \quad \square$$

LEMMA 3.12. *If  $I \subset \subset \wedge$ , then  $[\chi_I T]^a$  is in  $B(\mathcal{H}, L^2(\wedge, \mathcal{C}))$ .*

*Proof.* By Assumption 2 we have

$$\chi_I T = \chi_I F E(\wedge) A^* = F E(I) A^*.$$

Thus

$$[\chi_I T]^a = F[E(I) A^*]^a,$$

and the latter operator is bounded by Lemma 3.5.  $\square$

LEMMA 3.13. *There is a linear map  $\tilde{T}$  of  $\mathcal{H}$  into  $L^2_{\text{loc}}(\wedge, \mathcal{C})$  such that*

- 1)  $\tilde{T}$  is an extension of  $T$
  - 2)  $D(\tilde{T}) = \mathcal{H}$
  - 3)  $\chi_I \tilde{T} = F[E(I) A^*], \quad I \subset \subset \wedge$
  - 4)  $\chi_I \tilde{T} \in B(\mathcal{H}, L^2(I, \mathcal{C})), \quad I \subset \subset \wedge.$
- (3.12)

*Proof.* For any  $I \subset \subset \wedge$ , define  $\chi_I \tilde{T}$  by (3.12). That this defines  $\tilde{T}$  a.e. and does not depend on the choice of  $I$  follows from

$$\begin{aligned} \chi_J \chi_I \tilde{T} &= \chi_J F[E(I) A^*]^a = F[E(J) E(I) A^*]^a \\ &= \chi_I F[E(J) A^*]^a. \end{aligned}$$

The other properties follow from the definition and Lemma 3.12.  $\square$

LEMMA 3.14.

$$(M(\lambda) u, v)_{\mathcal{H}} = (\tilde{T}_{\lambda} u, \tilde{T}_{\lambda} v)_{\mathcal{C}} \quad \text{a.e., } u, v \in \mathcal{H}.$$

*Proof.* By Lemma 3.9, this holds a.e. for  $u, v \in D(A^*)$ . Thus for any  $I \subset \subset \wedge$ ,

$$\int_I (M(\lambda) u, v)_{\mathcal{H}} d\lambda = \int_I (\tilde{T}_{\lambda} u, \tilde{T}_{\lambda} v)_{\mathcal{C}} d\lambda \quad (3.13)$$

holds for all  $u, v \in D(A^*)$ . Since both sides of (3.13) are continuous on  $\mathcal{H} \times \mathcal{H}$  and  $D(A^*)$  is dense in  $\mathcal{H}$ , we see that (3.13) holds for  $u, v \in \mathcal{H}$ . Since this is true for any  $I \subset \subset \wedge$ , the lemma follows.  $\square$

LEMMA 3.15. *Set*

$$T_{\pm} = T_{\pm\lambda} = \tilde{T}_{\lambda} G_{1\pm}(\lambda).$$

*Then*

$$\frac{d}{d\lambda} (E_1(\lambda) A^* u, A^* v) = (T_{\pm\lambda} u, T_{\pm\lambda} v)_{\mathcal{H}} \quad \text{a.e., } u, v \in D(A^*). \quad (3.14)$$

*Proof.* Let  $\varepsilon \rightarrow 0$  in (3.11). The left hand side converges to the left side of (3.14) a.e. By Lemmas 3.4 and 3.10, the right hand side converges to

$$(M(\lambda) G_{1\pm}(\lambda) u, G_{1\pm}(\lambda) v)_{\mathcal{H}}$$

a.e. in  $\Lambda - e$ . We now apply Lemma 3.14.  $\square$

Since  $R(A^*)$  is dense in  $\mathcal{H}$ , elements of the form

$$u = \sum_{k=1}^l E_1(I_k) A^* u_k, \quad I_k \subset \subset \Lambda - e, u_k \in D(A^*) \quad (3.15)$$

are dense in  $E_1(\Lambda - e) \mathcal{H}$ . For such an element, set

$$u_{\pm}(\lambda) = \sum_{k=1}^l \chi_{I_k}(\lambda) T_{\pm\lambda} u_k. \quad (3.16)$$

Clearly, this is an element of  $L^2(\Lambda, \mathcal{H})$  with the square of its norm equal to

$$\begin{aligned} & \sum_{I_j \cap I_k} \int (T_{\pm\lambda} u_j, T_{\pm\lambda} u_k)_{\mathcal{H}} d\lambda \\ &= \sum (E_1(I_j) A^* u_j, E_1(I_k) A^* u_k) = \|u\|^2 \end{aligned}$$

by (3.14). Thus the mappings  $u \rightarrow u_{\pm}$  can be extended to isometries  $F_{\pm}$  of  $E_1(\Lambda - e) \mathcal{H}$  into  $L^2(\Lambda, \mathcal{H})$ .

LEMMA 3.16. *The operators  $F_{\pm}$  are surjective.*

*Proof.* Suppose  $h(\lambda)$  is in  $L^2(\Lambda, \mathcal{H})$  and

$$\int_{\Lambda} (F_{\pm} u, h)_{\mathcal{H}} d\lambda = 0, \quad u \in E_1(\Lambda - e) \mathcal{H}.$$

In particular,

$$\int_{\Lambda} (F_{\pm} E_1(I) A^* u, h)_{\mathcal{H}} d\lambda = 0$$

holds for any  $I \subset \subset \wedge - e$  and  $u \in D(A^*)$ . By (3.15) and (3.16), this means

$$\int_I (T_{\pm} u, h)_{\mathcal{H}} d\lambda = 0, \quad I \subset \subset \wedge - e, u \in D(A^*)$$

If  $I \subset \subset \wedge - e$ , then  $\chi_I T_{\pm}$  is a continuous map from  $\mathcal{H}$  to  $L^2(I, \mathcal{H})$  (Lemmas 3.4 and 3.13). Since  $D(A^*)$  is dense in  $\mathcal{H}$ , this implies

$$\int_I (T_{\pm} u, h)_{\mathcal{H}} d\lambda = 0, \quad I \subset \subset \wedge - e, \quad u \in \mathcal{H}.$$

Since this is true for any such interval  $I$ , we have

$$(T_{\pm \lambda} u, h(\lambda))_{\mathcal{H}} = 0 \quad \text{a.e., } u \in \mathcal{H},$$

or

$$(\tilde{T}_{\lambda} G_{1\pm}(\lambda) u, h(\lambda))_{\mathcal{H}} = 0 \quad \text{a.e., } u \in \mathcal{H}$$

Since  $G_{1\pm}(\lambda)$  is surjective for  $\lambda \in \wedge - e$ , this implies

$$(\tilde{T}_{\lambda} w, h(\lambda))_{\mathcal{H}} = 0 \quad \text{a.e., } w \in \mathcal{H}.$$

In particular, we have

$$(T_{\lambda} v, h(\lambda))_{\mathcal{H}} = 0 \quad \text{a.e., } v \in D(A^*).$$

Thus

$$\int_{\wedge} (T_{\lambda} v, h(\lambda))_{\mathcal{H}} d\lambda = 0, \quad v \in D(A^*).$$

Since  $R(T)$  is dense in  $L^2(\wedge, \mathcal{H})$  (Lemma 3.7), this implies that  $h(\lambda) = 0$  a.e.  $\square$

**LEMMA 3.17.** *Elements of the form (3.16) are dense in  $L^2(\wedge, \mathcal{H})$ .*

*Proof.* By Lemma 3.16 every element in  $L^2(\wedge, \mathcal{H})$  is the limit of such elements.

**LEMMA 3.18.**  *$F_{\pm} E_1(J) F_{\pm}^{-1} = \chi_J$  on  $L^2(\wedge, \mathcal{H})$  for each  $J \subset \wedge - e$ .*

*Proof.* By Lemma 3.17, it suffices to verify this on elements of the form

$$g(\lambda) = \chi_I(\lambda) T_{\pm \lambda} u, \quad I \subset \subset \wedge - e, \quad u \in D(A^*).$$

By (3.15) and (3.16),

$$F_{\pm}^{-1} g = E_1(I) A^* u.$$

Thus

$$\begin{aligned} F_{\pm} E_1(J) F_{\pm}^{-1} g &= F_{\pm} E_1(I \cap J) A^* u \\ &= \chi_{I \cap J}(\lambda) T_{\pm \lambda} u = \chi_J(\lambda) g(\lambda). \quad \square \end{aligned}$$

LEMMA 3.19. *Set*

$$\tilde{M}(\lambda) = G_{1\pm}(\lambda)^* M(\lambda) G_{1\pm}(\lambda).$$

*Then  $\tilde{M}(\lambda)$  is in  $B(\mathcal{H})$  for each  $\lambda \in \Lambda - e$  and it is integrable over each set  $I \subset \subset \Lambda - e$ . Moreover, for each bounded, Borel measurable function  $f(\lambda)$ ,*

$$[Af(H_1) \delta_{\varepsilon}(H_1 - \lambda) A^*]^a \rightarrow f(\lambda) \tilde{M}(\lambda)$$

*in norm a.e. as  $\varepsilon \rightarrow 0$ .*

*Proof.* The first statement is obvious. To prove the second, suppose  $I \subset \subset \Lambda - e$ . For  $\lambda$  in the interior of  $I$  we write

$$\begin{aligned} [Af(H_1) \delta_{\varepsilon}(H_1 - \lambda) A^*]^a &= [Af(H_1) \delta_{\varepsilon}(H_1 - \lambda) E(I) A^*]^a \\ &\quad + [Af(H_1) \delta_{\varepsilon}(H_1 - \lambda) E(R - I) A^*]^a. \end{aligned}$$

The first term equals

$$\frac{\varepsilon}{\pi} \int_I \frac{1}{(\mu - \lambda)^2 + \varepsilon^2} f(\mu) \tilde{M}(\mu) d\mu$$

by Lemma 3.15. This converges to  $f(\lambda) \tilde{M}(\lambda)$  in norm a.e. as  $\varepsilon \rightarrow 0$ . As in the proof of Lemma 3.10, the second term goes to 0 in norm.  $\square$

LEMMA 3.20.  $W_{\pm} E(I) = E_1(I) W_{\pm}$  on  $E(\Lambda) \mathcal{H}$  for each  $I \subset \Lambda - e$ , where  $W_{\pm} = F_{\pm}^* F$ .

*Proof.* By Lemma 3.18,

$$F_{\pm} E_1(I) = \chi_I F_{\pm}.$$

Thus

$$E_1(I) F_{\pm}^* = F_{\pm}^* \chi_I.$$

Hence by Assumption 2,

$$\begin{aligned} W_{\pm} E(I) &= F_{\pm}^* F E(I) = F_{\pm}^* \chi_I F \\ &= E_1(I) F_{\pm}^* F = E_1(I) W_{\pm}. \quad \square \end{aligned}$$

LEMMA 3.21. Let  $B_0$  be the set of complex valued Borel measurable functions with compact supports in  $\Lambda - e$ . Then for each  $f \in B_0$  and each  $u \in D(A^*)$

$$AR_1(\lambda + i\varepsilon)f(H_1)A^*u \rightarrow h(\lambda, f, u) \quad (3.17)$$

in  $L^2(R, \mathcal{K})$  as  $\varepsilon \rightarrow 0$ , and

$$\int_{-\infty}^{\infty} \|h(\lambda, f, u)\|_{\mathcal{K}}^2 d\lambda = 2\pi \int_0^{\infty} d\xi \left\| \int e^{-i\xi\mu} f(\mu) \tilde{M}(\mu) u d_{\mu} \right\|_{\mathcal{K}}^2. \quad (3.18)$$

*Proof.* Let  $I \subset \subset \Lambda - e$  contain the support of  $f$ . By lemmas 3.14 and 3.15

$$AR_1(\lambda + i\varepsilon)f(H_1)E_1(I)A^*u = \int_I \frac{f(\mu)}{\lambda + i\varepsilon - \mu} \tilde{M}(\mu) u d_{\mu}.$$

Since  $f(x) \tilde{M}(x) u \in L^2(R, \mathcal{K})$ , this converges in  $L^2(R, \mathcal{K})$  to an element  $h(\lambda, f, u)$ . The identity (3.18) follows from the theory of Fourier transforms.  $\square$

LEMMA 3.22. For  $g \in B_0$ ,  $u, v \in D(A^*)$ ,

$$\begin{aligned} \frac{d}{d\lambda} (E(\lambda) A^*u, g(H_1) A^*v) &= g(\lambda) (\tilde{M}(\lambda) G_+(\lambda) u, v)_{\mathcal{K}} \\ &+ \frac{1}{2\pi i} ([G_+(\lambda) - G_-(\lambda)] u, h(\lambda, g, v))_{\mathcal{K}}. \end{aligned} \quad (3.19)$$

*Proof.* Put  $\zeta = \lambda + i\varepsilon$ . Then by (3.6)

$$\begin{aligned} 2\pi i (\delta_{\varepsilon}(H - \lambda) A^*u, g(H_1) A^*v) &= ([R(\zeta) - R(\bar{\zeta})] A^*u, g(H_1) A^*v) \\ &= (\{[R_1(\zeta) A^*]^a G(\zeta) - [R_1(\bar{\zeta}) A^*]^a G(\bar{\zeta})\} u, g(H_1) A^*v) \\ &= 2\pi i ([\delta_{\varepsilon}(H_1 - \lambda) A^*]^a G(\zeta) u, g(H_1) A^*v) \\ &+ ([G(\zeta) - G(\bar{\zeta})] u, AR_1(\zeta) g(H_1) A^*v)_{\mathcal{K}}. \end{aligned}$$

Take the limit as  $\varepsilon \rightarrow 0$  and use Lemma 3.21.  $\square$

LEMMA 3.23. For  $f, g \in B_0$  and  $u, v \in D(A^*)$ , we have

$$\begin{aligned} ([W_+ - I] f(H) A^*u, g(H_1) A^*v) &= -\frac{1}{2\pi i} \int_{\Lambda} f(\lambda) ([G_+(\lambda) - G_-(\lambda)] u, h(\lambda, g, v))_{\mathcal{K}} d\lambda. \end{aligned} \quad (3.20)$$

*Proof.* Multiply (3.19) by  $f(\lambda)$  and integrate over  $\Lambda$ . The left side becomes  $(f(H) A^*u, g(H_1) A^*v)$ . The first term on the right becomes

$$\begin{aligned} & \int_{\Lambda} f(\lambda) \overline{g(\lambda)} (T_{\lambda}u, T_{+\lambda}v)_{\mathcal{H}} d\lambda \\ &= \int_{\Lambda} (Ff(H) E(\Lambda) A^*u, g(\lambda) T_{+\lambda}v)_{\mathcal{H}} d\lambda \\ &= \int_{\Lambda} (Ff(H) A^*u, F_{+}g(H_1) A^*v)_{\mathcal{H}} d\lambda \\ &= (W_{+}f(H) A^*u, g(H_1) A^*v), \end{aligned}$$

by Lemmas 3.14, 3.15 and 3.18.  $\square$

LEMMA 3.24. *Let  $\varphi(\lambda)$  be a real valued Borel measurable function on  $\Lambda$  such that*

$$\int_0^{\infty} \left| \int_I e^{-it\varphi(\lambda) - i\xi\lambda} d\lambda \right|^2 d\xi \rightarrow 0 \quad \text{as } t \rightarrow \infty \quad (3.21)$$

*for each  $I \subset \subset \Lambda - e$ . Then*

$$e^{it\varphi(H_1)} e^{-it\varphi(H)} E(\Lambda) u \rightarrow W_{+} E(\Lambda) u \quad \text{as } t \rightarrow \infty$$

*for each  $u \in \mathcal{H}$ .*

*Proof.* Suppose  $I \subset \subset \Lambda$  and  $J \subset \subset \Lambda - e$ . Put  $f(\lambda) = \chi_I e^{-it\varphi(\lambda)}$  and  $g(\lambda) = \chi_J e^{-it\varphi(\lambda)}$ .

By Lemma 3.23,

$$\begin{aligned} & ([W_{+} - I] e^{-it\varphi(H)} E(I) A^*u, e^{-it\varphi(H_1)} E_1(J) A^*v) \\ &= -\frac{1}{2\pi i} \int_I e^{-it\varphi(\lambda)} ([G_{+}(\lambda) - G_{-}(\lambda)] u, h(\lambda, g, v))_{\mathcal{H}} d\lambda \end{aligned}$$

holds for all  $u, v \in D(A^*)$ . In absolute value the square of this is

$$\leq \frac{1}{4\pi^2} \int_I \| [G_{+}(\lambda) - G_{-}(\lambda)] \|^2_{\mathcal{H}} d\lambda \int_I \| h(\lambda, g, v) \|^2_{\mathcal{H}} d\lambda.$$

By (3.18)

$$\int_{-\infty}^{\infty} \|h(\lambda, g, v)\|_{\mathcal{H}}^2 d\lambda = 2\pi \int_0^{\infty} d\xi \left\| \int_I e^{-i\xi\mu - it\varphi(\mu)} \tilde{M}(\mu) v d\mu \right\|_{\mathcal{H}}^2 \rightarrow 0$$

by (3.21). Since linear combinations of elements of the form  $E(I) A^* u$  are dense in  $E(\wedge) \mathcal{H}$  and linear combinations of elements of the form  $E_1(J) A^* v$  are dense in  $E_1(\wedge - e) \mathcal{H}$ , we have

$$([W_+ - I] e^{-it\varphi(H)} u, e^{-it\varphi(H_1)} v) \rightarrow 0$$

as  $t \rightarrow \infty$  for  $u \in E(\wedge) \mathcal{H}$  and  $v \in E_1(\wedge - e) \mathcal{H}$ . By Lemma 2.20, this implies

$$(e^{it\varphi(H_1)} e^{-it\varphi(H)} u, v) \rightarrow (W_+ u, v) \quad (3.22)$$

for all such  $u$  and  $v$ . Thus for such  $u$  we have

$$\begin{aligned} \|[e^{it\varphi(H_1)} e^{-it\varphi(H)} - W_+] u\|^2 &= \|e^{it\varphi(H_1)} e^{-it\varphi(H)} u\|^2 \\ &\quad + \|W_+ u\|^2 - 2\operatorname{Re}(e^{it\varphi(H_1)} e^{-it\varphi(H)} u, W_+ u) \\ &= 2\|u\|^2 - 2\operatorname{Re}(e^{it\varphi(H_1)} e^{-it\varphi(H)} u, W_+ u). \end{aligned}$$

Since  $R(W_+) = E_2(\wedge - e) \mathcal{H}$ , this tends to 0 as  $t \rightarrow \infty$ .  $\square$

The conclusions of Theorem 3.1 are contained in those of Lemmas 3.2–3.24.

We now make an important observation due to Kato and Kuroda [2].

**LEMMA 3.25.** *Assumption 6 is implied by 6'. There is a function  $N(\lambda)$  from  $\wedge$  to  $B(\mathcal{H})$  which is locally Holder continuous and such that*

$$\frac{d}{d\lambda} (E(\lambda) A^* u, B^* v) = (N(\lambda) u, v)_{\mathcal{H}}, \quad u \in D(A^*), \quad v \in D(B^*). \quad (3.23)$$

*Proof.* As in the case of Lemma 3.8 we get

$$[Bf(H) E(I) A^*] = \int_I f(\lambda) N(\lambda) d\lambda \quad (3.24)$$

for each Borel  $I \subset \wedge$  and each bounded Borel measurable function  $f(\lambda)$ . Now suppose  $I \subset \subset \wedge$ . Then

$$\begin{aligned} Q(\zeta) &= B[R(i) A^*]^a + (i - \zeta) BR(i) E(R - I) [R(i) A^*]^a \\ &\quad + (i - \zeta)^2 BR(i) R(\zeta) E(R - I) [R(i) A^*]^a \\ &\quad + (i - \zeta) BR(\zeta) E(I) [R(i) A^*]^a. \end{aligned}$$



All but the last term on the right can be easily extended to be continuous in  $\Pi_I^\pm$ . By (3.24)

$$BR(\zeta) E(I) [R(i) A^*]^a = \int_I \frac{1}{(\zeta - \mu)(i - \mu)} N(\lambda) d\lambda.$$

Since  $N(\lambda)$  is Hölder continuous in  $I$ , Privalov's theorem implies that this too can be extended to be continuous in  $\Pi_I^\pm$ . Thus  $Q(\zeta)$  has a continuous extension to  $\Pi_\wedge^\pm$ .  $\square$

#### 4. The Application

We now turn to the proofs of the theorems of Section 2. We take  $H$  in Theorem 3.1 to be  $P_0$ , the closure in  $L^2$  of  $P(D)$  on  $C_0^\infty$ . We know that

$$\sigma(P_0) = \{P(\xi), \xi \in E^n\}$$

([14, p. 65]). One checks easily that this set is merely the interval  $[\lambda_0, \infty)$ , where

$$\lambda_0 = \min_{\xi \in E^n} P(\xi).$$

We take  $\wedge = \sigma(P_0) - \Gamma$ . Thus  $E(\wedge)$  is the identity on  $L^2$ . Assumption 2 of Theorem 3.1 is easily verified (cf. [8] or [7, II]). Next, let  $\mathcal{K}$  be the direct sum of  $N^2$  copies of  $L^2$ , with each summand having indices  $j, k$  each running from 1 to  $N$ . Elements of  $\mathcal{K}$  will be denoted by  $\{u_{jk}\}$ . We define

$$Av = \{\bar{g}_{jk} P_j(D) v\}, Bu = \{h_{jk} P_k(D) u\}.$$

If we define  $A$  and  $B$  first on  $C_0^\infty$ , we see that they are closable. By considering the closures, we obtain closed operators from  $H = L^2$  to  $\mathcal{K}$ . Note that

$$A^* \{u_{jk}\} = \sum \bar{P}_j(D) g_{jk} u_{jk} \quad (4.1)$$

and

$$B^* \{v_{jk}\} = \sum \bar{P}_k(D) h_{jk} v_{jk}. \quad (4.2)$$

In particular we have

$$(Q(x, D) u, v) = (Bu, Au)_{\mathcal{K}} \quad (4.3)$$

For  $s$  real, let  $H^{s,2}$  denote the completion of  $C_0^\infty$  with respect to the norm given by

$$\|\varphi\|_{s,2}^2 = \int (1 + |\xi|^2)^s |F\varphi|^2 d\xi,$$

where  $F$  denotes the Fourier transform. When  $s$  is a positive integer, the norm of  $H^{s,2}$  is equivalent to the sum of the  $L^2$  norms of all derivatives up to order  $s$ . By (2.3)

$$(P_0 u, u) \geq a \|u\|_{m,2}^2 - K_1 \|u\|^2,$$

where  $K_1$  is some constant independent of  $u$ . Moreover, hypothesis (II) implies that there is a constant  $K_2$  such that

$$\|Au\|^2 + \|Bu\|^2 \leq \frac{1}{2} a \|u\|_{m,2}^2 + K_2 \|u\|^2$$

(cf. [14, p. 140]). Thus the symmetric bilinear form

$$h(u, v) = (P_0 u, v) + (Bu, Av)_{\mathcal{H}}, \quad u, v \in H^{m,2} \quad (4.4)$$

is bounded from below. The operator associated with it is self-adjoint (cf. Theorem 7.6, chapter 1 of [14]). Since  $C_0^\infty$  is dense in  $H^{m,2}$ , this operator is unique, and Lemma 2.1 is proved. This also shows that Assumption 3 of Theorem 3.1 is satisfied. It also follows from hypothesis (II) that  $h_{jk} P_k(D)$  is  $P_0$ -compact for each  $j$  and  $k$  ([14, p. 112]). Thus  $B$  is  $P_0$ -compact and Assumption 5 of Section 3 is verified. We have also verified Assumption 8.

Next we note that we can always arrange things so that  $A$  is injective. In fact, if we set  $P_0(\xi) = 1$ ,  $h_{0k}(x) \equiv 0$  and let  $g_{0k}(x)$  be functions which do not vanish anywhere and satisfy hypotheses (I)–(III), then

$$Q(x, D) = \sum_{j,k=0}^N \bar{P}_j(D) g_{jk}(x) h_{jk}(x) P_k(D),$$

and nothing is changed. Now  $\mathcal{H}$  becomes  $(N+1)^2$  copies of  $L^2$  and  $A$  is injective. It therefore remains only to verify that Assumptions 6 and 7 are satisfied.

A simple computation shows that

$$\pi F [\delta_\varepsilon (P_0 - \lambda) w] = \frac{\varepsilon}{[P(\xi) - \lambda]^2 + \varepsilon^2} F w.$$

Put

$$\begin{aligned} \psi_{\lambda, \varepsilon}(\xi) &= \frac{\varepsilon}{[P(\xi) - \lambda]^2 + \varepsilon^2}, \quad |P(\xi) - \lambda| < 1 \\ &= 0, \quad |P(\xi) - \lambda| \geq 1, \end{aligned}$$

and

$$\varphi_{\lambda, \varepsilon}(\xi) = \frac{\varepsilon}{[P(\xi) - \lambda]^2 + \varepsilon^2} - \psi_{\lambda, \varepsilon}(\xi).$$

Then by definition,

$$K_\lambda(x) = \lim F(\psi_{\lambda, \varepsilon}),$$

when  $F$  denotes the inverse Fourier transform. Since  $|\varphi_{\lambda, \varepsilon}| \leq \varepsilon$ , we have

$$(\varphi_{\lambda, \varepsilon} u, v) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0, u, v \in L^2.$$

Thus if  $u, v$  are functions in  $L^2$  with compact supports, then

$$\begin{aligned} \pi(\delta_\varepsilon(P_0 - \lambda) u, v) &= ([\psi_{\lambda, \varepsilon} + \varphi_{\lambda, \varepsilon}] Fu, Fv) \\ &= (F(\psi_{\lambda, \varepsilon}) * u, v) + (\varphi_{\lambda, \varepsilon} Fu, Fv) \rightarrow (K_\lambda * u, v) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Thus if  $\{E(\lambda)\}$  denotes the spectral family of  $P_0$ , we have

$$\pi \frac{d}{d\lambda} (E(\lambda) u, v) = (K_\lambda * u, v). \quad (4.6)$$

Thus Assumption 7 will be verified if we can show that the operator  $AK_\lambda * A^*$  can be extended to a bounded operator on  $\mathcal{H}$  which is locally Hölder continuous with respect to the norm topology. Likewise by Lemma 3.25, Assumption 6 will be satisfied if we can do the same for the operator  $BK_\lambda * A^*$ . We note that hypothesis (III) implies just that for both of these operators. In fact we have

$$\begin{aligned} & |(K_\lambda * [P_j(D) g_{jk} u], \bar{P}_t(D) h_{st} v)|^2 \\ & \leq \int \int |g_{jk}(g)|^2 |P_j(D) P_t(D) K_\lambda(x-y)| |v(x)|^2 dx dy \\ & \times \int \int |h_{s,t}(x)|^2 |P_j(D) P_t(D) K_\lambda(x-y)| |u(y)|^2 dx dy \\ & \leq C \|u\|^2 \|v\|^2. \end{aligned} \quad (4.7)$$

with the other inequalities similarly verified. This completes the proof of Theorem 3. A different variation of the above inequality proves Theorem 2.4. In fact we have

$$\begin{aligned} & |(K_\lambda * [P_j(D) g_{jk} u], \bar{P}_t(D) h_{st} v)|^2 \\ & \leq \int \int |g_{jk}(y) h_{st}(x)|^2 |P_j(D) P_t(D) K_\lambda(x-y)|^2 dx dy \\ & \times \int \int |u(y) v(x)|^2 dx dy \leq C \|u\|^2 \|v\|^2, \end{aligned} \quad (4.8)$$

with similar calculations for the other inequalities. This proves Theorem 2.4.

We postpone the proof of Theorem 2.5 until the next section, and turn our atten-

tion to Theorems 2.6 and 2.7. First we need some lemmas. Let  $S^{n-1}$  denote the unit sphere  $|x|=1$  in  $E^n$ , and let  $(\cdot, \cdot)_S$  denote the scalar product in  $L^2(S^{n-1})$ .

LEMMA 4.1. *If  $s > \frac{1}{2}$ , then*

$$\gamma_\lambda h(\omega) = h(\lambda \omega), \quad \omega \in S^{n-1}, \quad (4.9)$$

*is a mapping from  $(0, \infty)$  to  $B(H^{s,2}, L^2(S^{n-1}))$  which is locally Hölder continuous. This was proved by Kuroda [6, Lemma 4.1].*

LEMMA 4.2. *Let  $I$  be any open interval in  $\mathbb{R}$ . Then there are a finite number of functions  $\gamma_k$  each mapping  $I$  into  $B(H^{s,2}, L^2(S^{n-1}))$  and locally Hölder continuous and such that*

$$\frac{d}{d\lambda} (E(\lambda) u, v) = \sum (\gamma_k F u, \gamma_k F v)_S, \quad u, v \in C_0^\infty. \quad (4.10)$$

*Proof.* Let  $r_1, \dots, r_l$  be the distinct positive roots of  $p'(r^2) = 0$  in ascending order. Let  $r_0 = 0, r_{l+1} = \infty$  and  $J_k = [r_{k-1}, r_k]$ . Let  $I_k$  be the range of  $p(r^2)$  on  $J_k$ . By the implicit function theorem there is a function  $f_k(t) \in C^\infty(I_k)$  such that

$$p(f_k(t)^2) = t, \quad t \in I_k \quad (4.11)$$

$$f_k(p(r^2)) = r, \quad r \in J_k. \quad (4.12)$$

Now

$$\begin{aligned} \pi(\delta_\varepsilon(P_0 - \lambda) u, v) &= \int_{-\infty}^{\infty} \frac{\varepsilon}{[p(|\xi|^2) - \lambda]^2 + \varepsilon^2} F u \overline{F v} d\xi \\ &= \int_0^\infty \frac{\varepsilon}{[p(r^2) - \lambda]^2 + \varepsilon^2} (F u(r\omega), F v(r\omega))_S r^{n-1} dr \\ &= \sum_k \int_{I_k} \frac{\varepsilon}{(t - \lambda)^2 + \varepsilon^2} (F u(f_k(t)\omega), F v(f_k(t)\omega))_S \\ &\quad \times f_k(t)^{n-1} |f'_k(t)| dt. \end{aligned}$$

Taking the limit as  $\varepsilon \rightarrow 0$ , we obtain

$$\frac{d}{d\lambda} (E(\lambda) u, v) = \sum_{\lambda \in I_k} (\gamma_k F u, \gamma_k F v)_S, \quad (4.12)$$

where

$$\gamma_k h(\omega) = f_k(\lambda)^{(n-1)/2} |f'_k(\lambda)|^{1/2} h(f_k(\lambda)\omega). \quad (4.13)$$

By Lemma 4.1,  $\gamma_k$  is a mapping from  $I_k$  to  $B(H^{s,2}, L^2(S^{n-1}))$  which is locally Hölder continuous. Since the endpoints of the  $I_k$  are contained in  $\Gamma$ , an arbitrary open interval  $I \subset \wedge$  is either contained in  $I_k$  or does not intersect it. This gives the lemma.

For a locally square integrable function  $h$  we put

$$\tilde{h}(x) = \int |h(y)|^2 e^{-|x-y|^2} dy.$$

LEMMA 4.3. *If*

$$\sup_x (1+|x|^2)^s \tilde{h}(x) < \infty, \quad (4.14)$$

*then the operator  $S$  given by*

$$Su = (1+|x|^2)^{1/2s} \int h(y) u(y) e^{-|x-y|^2} dy \quad (4.15)$$

*is bounded on  $L^2$ .*

*Proof.* We have

$$(Su, v) = \int \int (1+|x|^2)^{s/2} e^{-|x-y|^2} h(y) u(y) \overline{v(x)} dx dy.$$

The square of this is bounded by

$$\begin{aligned} & \int \int (1+|x|^2)^s e^{-|x-y|^2} |h(y) v(x)|^2 dx dy \\ & \times \int \int e^{-|x-y|^2} |u(y)|^2 dx dy \leq C \|v\|^2 \|u\|^2. \end{aligned}$$

COROLLARY 4.4. *If (4.14) holds, then the operator  $L$  given by*

$$Lu = F[e^{-|x|^2} * (hu)] \quad (4.16)$$

*is bounded from  $L^2$  to  $H^{s,2}$*

*Proof.* We have

$$\|Lu\|_{s,2} = \|(1+|x|^2)^{s/2} [e^{-|x|^2} * (hu)]\|.$$

Apply Lemma 4.3.  $\square$

LEMMA 4.5. *If (4.14) holds, then there exists a mapping  $T_\lambda$  of  $\wedge$  to  $B(L^2)$  which is locally Hölder continuous and such that*

$$\frac{d}{d\lambda} (E(\lambda)(hu), hv) = (T_\lambda u, v), \quad u, v \in L^2. \quad (4.17)$$

*Proof.* It clearly suffices to prove (4.12) for  $u, v \in C_0^\infty$ . Let  $I$  be a component of  $\Lambda$ . For those  $I_k$  which contain  $I$ , set

$$\psi_k(\xi) = \exp \left\{ \frac{1}{4} (f_k(\lambda)^2 - |\xi|^2) \right\} \quad (4.18)$$

(see the proof of Lemma 4.2). Then

$$F\psi_k = \exp [f_k(\lambda)^2 - |x|^2] \equiv \tilde{\psi}_k, \quad (4.19)$$

and

$$\psi_k(f_k(\lambda) \omega) = 1. \quad (4.20)$$

Thus

$$\begin{aligned} (\gamma_k F(hu), \gamma_k F(hv))_S &= (\gamma_k \psi_k F(hu), \gamma_k \psi_k F(hv))_S \\ &= (\gamma_k F[\tilde{\psi}_k * (hu)], \gamma_k F[\tilde{\psi}_k * (hv)])_S \\ &= (T_k u, v), \end{aligned} \quad (4.21)$$

where

$$T_k = e^{2f_k(\lambda)^2} L^* \gamma_k^* \gamma_k L. \quad (4.22)$$

By Lemma 4.1 and Corollary 4.4, this is a locally Hölder continuous map of  $I_k$  into  $B(L^2)$ . If we put

$$T_\lambda = \sum_{I \in I_k} T_k \quad (4.23)$$

and apply Lemma 4.2, we obtain (4.17).

LEMMA 4.6. *If there is a  $\beta < \frac{1}{2}(n-1)$  such that*

$$\sup_x \int \tilde{h}(y) (1 + |x - y|)^{-\beta} dy < \infty, \quad (4.24)$$

*then (4.17) holds for some locally Hölder continuous map  $T_\lambda$  from  $\Lambda$  to  $B(L^2)$ .*

*Proof.* By (4.21), (4.10) and (4.6)

$$\begin{aligned} \sum (\gamma_k F(hu), \gamma_k F(hv))_S &= \sum (\gamma_k F[\tilde{\psi}_k * (hu)], \gamma_k F[\tilde{\psi}_k * (hv)])_S \\ &= \frac{d}{d\lambda} (E(\lambda) [\tilde{\psi}_k * (hu)], \tilde{\psi}_k * (hv))_S \\ &= (K_\lambda * [\tilde{\psi}_k * (hu)], \tilde{\psi}_k * (hv))_S \\ &= \sum e^{2f_k(\lambda)^2} \int \int \int \int K_\lambda(x - y) e^{-|y - t|^2} h(z) u(z) \\ &\quad \times e^{-|x - t|^2} \overline{h(t) v(t)} dx dy dz dt. \end{aligned}$$

By (2.18) and (2.19), it suffices to show that the integral

$$\int \int \int \int (1+|x-y|)^{-\beta} e^{-|y-z|^2-|x-t|^2} |h(z) u(z) h(t) v(t)| dx dy dz dt$$

is bounded by a constant times  $\|u\| \|v\|$ . But its square is bounded by

$$\begin{aligned} & \int \int \int \int (1+|x-y|)^{-\beta} e^{-|y-z|^2-|x-t|^2} |h(z) v(t)|^2 dx dy dz dt \\ & \times \int \int \int \int (1+|x-y|)^{-\beta} e^{-|y-z|^2-|x-t|^2} |h(t) u(z)|^2 dx dy dz dt. \end{aligned}$$

By (4.24), this is bounded by a constant times

$$\int \int e^{-|x-t|^2} |v(t)|^2 dx dt \int \int e^{-|y-z|^2} |u(z)|^2 dy dz$$

which equals a constant times  $\|u\|^2 \|v\|^2$ .  $\square$

For  $h$  locally square integrable and  $a > 0$  set

$$\hat{h}_a(x) = \int_{|x-y| < a} |h(y)|^2 dy \quad (4.25)$$

LEMMA 4.7. *For each  $a > 0$  there is a constant  $C_a$  depending only on  $a$  and  $n$  such that*

$$\|\tilde{h}\|_p \leq C_a \|\hat{h}_a\|_p, \quad 1 \leq p \leq \infty. \quad (4.26)$$

*Proof.* For each  $k$ , let  $z_1^{(k)}, \dots, z_{N(k)}^{(k)}$  be points such that the set  $ka < |x| < (k+1)a$  is covered by  $N(k)$  balls of radius  $a$  and centers at the  $z_j^{(k)}$ . We know that there is a constant  $c_n$  depending only on  $n$  such that

$$N(k) \leq c_n (ka)^{n-1}.$$

Now

$$A_k(x) \equiv \int_{ka < |x-y| < (k+1)a} |h(y)|^2 e^{-|x-y|^2} dy \leq e^{-k^2 a^2} \sum_{j=1}^{N(k)} \hat{h}_a(x + z_j^{(k)}).$$

Thus

$$\|\tilde{h}\|_p \leq \sum_{k=0}^{\infty} \|A_k\|_p \leq \sum_{k=0}^{\infty} e^{-k^2 a^2} \sum_{j=1}^{N(k)} \|\hat{h}_a\|_p \leq c_n \|\hat{h}_a\|_p \sum_{k=0}^{\infty} (ka)^{n-1} e^{-k^2 a^2}. \quad \square$$

LEMMA 4.8. *For each  $a > 0$  and  $\alpha \geq 0$  there is a constant  $C$  depending only on*

$a, \alpha$  and  $n$  such that

$$\|(1+|x|)^\alpha \tilde{h}\|_\infty \leq C \|(1+|x|)^\alpha \hat{h}_a\|_\infty. \quad (4.27)$$

*Proof.* We may assume  $a \leq 1$ . Set

$$A_k(x) = (1+|x|)^\alpha \int_{ka < |x-y| < (k+1)a} |h(y)|^2 e^{-|x-y|^2} dy.$$

Let  $c_n$  be as in the preceding proof. If  $|x| \leq (2k+2)a$ , then

$$A_k \leq (2k+3)^\alpha c_n (ka)^{n-1} e^{-k^2 a^2} \hat{C},$$

where  $\hat{C}$  denotes the right side of (4.27). On the other hand, if  $|x| > (2k+2)a$ , then  $|x-y| < (k+1)a$  implies  $|y| > \frac{1}{2}|x|$ . Thus in this case

$$A_k \leq (1+|x|)^\alpha e^{-k^2 a^2} c_n (ka)^{n-1} (1+\frac{1}{2}|x|)^{-\alpha} \hat{C},$$

since the centers of the covering balls are a distance  $> \frac{1}{2}|x|$  from the origin. Combining the two estimates, we get

$$A_k \leq \hat{C} c_n 2^\alpha (2k+3)^\alpha (ka)^{n-1} e^{-k^2 a^2}$$

for any  $x$ . Since the left hand side of (4.27) is  $\|\sum A_k\|_\infty$ , the result follows.  $\square$

**THEOREM 4.9.** *Suppose there are constants  $a > 0$ ,  $a \geq 0$  and  $p \leq \infty$  satisfying (1.5) such that*

$$(1+|x|)^\alpha \hat{h}_a \in L^p \quad (4.28)$$

*Then there is a mapping  $T_\lambda$  from  $\wedge$  to  $N(L^2)$  which is locally Hölder continuous and such that (4.17) holds.*

*Proof.* Let  $c$  be a number such that

$$\alpha > c > 1 - \frac{2n}{(n+1)p},$$

and set

$$\begin{aligned} t &= p, & p < 2n/(n+1) \\ &= \max[(1-c)p, 0], & p \geq 2n/(n+1). \end{aligned}$$

Thus  $t < 2n/(n+1)$  and consequently

$$\|(1+|x|)^{-\beta}\|_{t'} < \infty, \quad t' = t/(t-1). \quad (4.29)$$



Put  $\theta = t/p$  and  $s = a/2(1 - \theta)$ . Then  $s > \frac{1}{2}$ . Let  $I$  be a component of  $\Lambda$ , and define  $T_\lambda$  by (4.23). Then  $T_\lambda$  maps  $C_0^\infty$  into the distributions and

$$\sum_{I \in I_k} (\gamma_k F(hu), \gamma_k F(hu))_S = (T_\lambda u, v), \quad u, v \in C_0^\infty. \quad (4.30)$$

If (4.24) holds, then  $T_\lambda$  can be extended to a bounded operator on  $L^2$ , and there is a  $b > 0$  such that

$$[T_\lambda] \equiv \|T_\lambda\| + \sup_{\lambda'} |\lambda - \lambda'|^{-b} \|T_\lambda - T_{\lambda'}\| \leq \text{constant} \|(1 + |x|)^{-\beta} * \tilde{h}\|_\infty$$

(Lemma 4.6). By Hölder's inequality and (4.29), this is bounded by a constant times  $\|\tilde{h}\|_r$ . By Lemma 4.7 this implies

$$[T_\lambda] \leq \text{constant} \|\hat{h}_a\|_r. \quad (4.31)$$

For the case  $t = p$ , we are finished. For then  $a = 0$  satisfies (1.5), and the result follows from (4.31). To take care of the other case, note that by Lemmas 4.5 and 4.8 there is a  $b > 0$  such that

$$[T_\lambda] \leq \text{constant} \|(1 + |x|^2)^s \hat{h}_a\|_\infty. \quad (4.32)$$

If we apply a general interpolation theorem due to Stein and Weiss [20] to (4.31) and (4.32), we obtain

$$\begin{aligned} [T_\lambda] &\leq \text{constant} \|(1 + |x|^2)^{(1-\theta)s} \hat{h}_a\|_{t/\theta} \\ &\leq \text{constant} \|(1 + |x|)^a \hat{h}_a\|_p. \end{aligned}$$

## 5. The Remaining Proofs

In this section we shall prove Lemma 2.2 and Theorem 2.5.

*Proof of Lemma 2.2.* Let  $I$  be an open interval containing  $\lambda$  and such that  $I \subset \Lambda$ . Set

$$\Omega = \{\xi \in E^n \mid P(\xi) \in I\}, \quad \Sigma = \{\xi \in E^n \mid P(\xi) = \lambda\}.$$

Then there is a  $C^\infty$  diffeomorphism  $\varphi$  of  $I \times \Sigma$  onto  $\Omega$  (cf. [7, II, Proposition 2.2]). Let  $J(t, \omega)$  denote the Jacobian of this transformation. Then

$$\begin{aligned} K_\lambda(x) &= \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \frac{\varepsilon e^{ix\xi} d\xi}{[P(\xi) - \lambda]^2 + \varepsilon^2} \\ &= \lim_{\varepsilon \rightarrow 0} \int_I \int_{\Sigma} \frac{\varepsilon J(t, \omega) \exp\{i\varphi(t, \omega)x\} dt d\omega}{(t - y)^2 + \varepsilon^2} \\ &= \pi \int_{\Sigma} J(\lambda, \omega) \exp\{i\varphi(\lambda, \omega)x\} d\omega, \end{aligned} \quad (5.1)$$

and the convergence is uniform for bounded  $x$ . The last formula shows that  $K_\lambda$  is infinitely differentiable.  $\square$

*Proof of Theorem 2.5.* Let the intervals  $I_k$ ,  $J_k$  and the functions  $f_k$  be given as in Lemma 4.2. Suppose  $\lambda$  is any point in  $\Lambda$ . If  $\lambda \in I_k$ , set  $t_k = f_k(\lambda)$ . By Bochner's formula [21, p. 235],  $K_\lambda(x)$  is the limit of

$$\begin{aligned}
 & 2\pi (2\pi/|x|)^\gamma \int_0^\infty \frac{\varepsilon r^{\gamma+1} J_\gamma(r|x|) dr}{[p(r^2) - \lambda]^2 + \varepsilon^2} \\
 &= 2\pi (2\pi/|x|)^\gamma \sum_k \int_{J_k} \\
 &= \pi (2\pi/|x|)^\gamma \sum_k \int_{I_k} \frac{\varepsilon f_k(s)^\gamma J_\gamma(f_k(s)|x|) ds}{[(s-\lambda)^2 + \varepsilon^2] p'(f_k(s)^2)} \\
 &\rightarrow \pi^2 (2\pi/|x|)^\gamma \sum \frac{t_k^\gamma J_\gamma(t_k|x|)}{p'(t_k^2)},
 \end{aligned} \tag{5.2}$$

where the summation is taken over those  $k$  for which  $\lambda \in I_k$ . This proves (2.17). To prove the rest of the lemma, we note that a simple induction shows that  $D^\mu K_\lambda$  is a finite sum of terms of the form

$$R(x) h(t) J_\nu(|x|t)/(|x|t)^\nu,$$

where  $\nu - \frac{1}{2}n$  is an integer  $\geq -1$ ,  $R(x)$  is a polynomial of degree  $\leq \nu + 1 - \frac{1}{2}n$ ,  $t$  is one of the  $t_k$  and  $h$  is infinitely differentiable. This immediately implies (2.18). To prove (2.19), set  $f(s) = J_\nu(s)/s^\nu$ . Then there is a constant  $c$  depending only on  $\nu$  such that

$$|f(s)| + |f'(s)| \leq c/(1+s)^{\nu+1/2}, \quad s \geq 0. \tag{5.3}$$

Moreover, for each positive integer  $k$  there is a constant  $C_k$  such that

$$|t-1|^k \leq C_k |t^k - 1|, \quad t \text{ real}. \tag{5.4}$$

Thus

$$\begin{aligned}
 |f(s_1) - f(s_2)|^k &\leq C_k |f(s_1)^k - f(s_2)^k| \\
 &= k C_k |f(\tilde{s})^{k-1} f'(\tilde{s}) (s_1 - s_2)| \\
 &\leq C |s_1 - s_2| (1 + \tilde{s})^{k(\nu+1/2)},
 \end{aligned}$$

where  $\tilde{s}$  is some value between  $s_1$  and  $s_2$ . Thus

$$|f(Rt_1) - f(Rt_2)| \leq C |t_1 - t_2|^{1/k} (1+R)^{(1/k) - \nu - 1/2}$$

This implies (2.19).  $\square$

## REFERENCES

- [1] KATO, TOSIO and KURODA, S. T., *Theory of simple scattering and eigenfunction expansions*, Functional Analysis and Related Field, Springer, 1970, 99–131.
- [2] —, *The abstract theory of scattering*, Rocky Mountain J. Math. 1 (1971), 127–171.
- [3] KATO, TOSIO, *Wave operators and similarity of non-self-adjoint operators*, Math. Ann. 162 (1966), 258–279.
- [4] —, *Perturbation Theory for Linear Operators*, Springer, 1966.
- [5] —, *Proc. Conf. Functional Analysis and Related Topics*, Tokyo Univ. Press, 1970, 206–215.
- [6] KURODA, S. T., *Some remarks on scattering for Schrödinger operators*, J. Fac. Sci. Univ. Tokyo, 17 (1970), 315–329.
- [7] —, *Scattering theory for differential operators, I and II*, to appear.
- [8] BEREZANSKII, Y. M., *Expansions in Eigenfunctions of self-adjoint Operators*, Amer. Math. Soc., Providence, 1968.
- [9] IKEBE, TERUO, *Eigenfunction expansions associated with the Schrödinger operator and their applications to scattering theory*, Arch. Rat. Mech. Anal. 5 (1960), 1–34.
- [10] ALSHOLM, P. and SCHMIDT, G., *Spectral and scattering theory for Schrödinger operators*, *ibid.*, 40 (1971), 281–311.
- [11] REJTO, P. A., *On partly gentle perturbations III*, J. Math. Anal. Appl. 27 (1969), 21–67.
- [12] GREIFENEGGER, V., JÖRGENS, K., WEIDMANN, J., and WINKLER, M., *Streutheorie für Schrödinger-Operatoren*, to appear.
- [13] SIMON, BARRY, *Quantum Mechanics for Hamiltonians Defined as Quadratic Forms*, Princeton Univ. Press, 1971.
- [14] SCHECHTER, MARTIN, *Spectra of Partial Differential Operators*, North-Holland, Amsterdam, 1971.
- [15] —, *Hamiltonians for singular potentials*, Indiana Univ. Math. J., 22 (1972), 483–503.
- [16] SCHECHTER, MARTIN and BULKA, ISAAC, to appear.
- [17] BEALS, RICHARD, *On spectral theory and scattering for elliptic operators with singular potentials*, Report, Math. Dept., Yale Univ., 1969.
- [18] BIRMAN, M. S., *Scattering problems for differential operators with constant coefficients*, Funk. Anal. Publ. 3 (1969), 1–16.
- [19] AGMON, SHMUEL, *Lecture at the Mathematics Research Institute, Oberwolfach*, 1971.
- [20] STEIN, E. M., and WEISS, GUIDO, *Interpolation of operators with change of measures*, Trans. Amer. Math. Soc., 87 (1958), 159–172.
- [21] BOCHNER, SALOMON, *Lectures on Fourier Integrals*, Annals of Mathematics Studies, Princeton Univ. Press, 1959.
- [22] SCHECHTER, MARTIN, *Principles of Functional Analysis*, Academic Press, New York, 1971.

Yeshiva University, New York  
The Hebrew University, Jerusalem

Received June 26, 1973.