

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 49 (1974)

Artikel: Algebraic L-Theory
Autor: Ranicki, A.A.
DOI: <https://doi.org/10.5169/seals-37984>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 21.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Algebraic L-Theory

IV. Polynomial Extension Rings

by A. A. RANICKI, Trinity College, Cambridge

Introduction

In Chapter XII of [1] Bass defines the notion of a *contracted functor*, as a functor

$$F: (\text{rings}) \rightarrow (\text{abelian groups})$$

such that the sequence

$$0 \rightarrow F(A) \xrightarrow{(-\bar{\varepsilon}_+)} F(A[x]) \oplus F(A[x^{-1}]) \xrightarrow{(E+E-)} F(A[x, x^{-1}]) \xrightarrow{B} LF(A) \rightarrow 0$$

is naturally split exact for any ring A (associative with 1), where

$$\bar{\varepsilon}_\pm: A \rightarrow A[x^{\pm 1}] \quad \bar{E}_\pm: A[x^{\pm 1}] \rightarrow A[x, x^{-1}]$$

are inclusions in polynomial extensions of A , and

$$\begin{aligned} B: F(A[x, x^{-1}]) &\rightarrow LF(A) \\ &= \text{coker}((\bar{E}_+ \bar{E}_-): F(A[x]) \oplus F(A[x^{-1}]) \rightarrow F(A[x, x^{-1}])) \end{aligned}$$

is the natural projection. Theorem 7.4 of Chapter XII of [1], the “Fundamental Theorem” of algebraic K -theory, states that

$$K_1: (\text{rings}) \rightarrow (\text{abelian groups})$$

is a contracted functor such that

$$LK_1(A) = K_0(A)$$

up to natural isomorphism. Here, we obtain analogous results for the groups of algebraic L -theory considered in the previous instalments of this series ([5], [6], [7] – we shall refer to these as Parts I, II, III respectively). In Part I we defined L -theoretic functors

$$U_n, V_n: (\text{rings with involution}) \rightarrow (\text{abelian groups})$$

for $n \pmod{4}$, using quadratic forms on $\begin{cases} \text{f.g. projective} \\ \text{f.g. free} \end{cases} A$ -modules for the $\begin{cases} U_- \\ V_- \end{cases}$ groups.

(The definitions are reviewed in §3 below, allowing this part to be read independently of the previous parts). It was shown in Part II that

$$V_n(A[x, x^{-1}]) = V_n(A) \oplus U_{n-1}(A)$$

if the involution $\bar{}: A \rightarrow A; a \mapsto \bar{a}$ is extended to $A[x, x^{-1}]$ by $\bar{x} = x^{-1}$. The main result of this part of the paper (Theorem 4.1) is a split exact sequence

$$0 \rightarrow V_n(A) \xrightarrow{(-\bar{\epsilon}_+)} V_n(A[x]) \oplus V_n(A[x^{-1}]) \xrightarrow{(E+E_-)} V_n(A[x, x^{-1}]) \xrightarrow{B} U_n(A) \rightarrow 0$$

for each $n \pmod{4}$, with the involution on A extended to $A[x^{\pm 1}]$, $A[x, x^{-1}]$ by $\bar{x} = x$. The proof depends on L -theoretic analogues (Lemmas 4.2, 4.3) of the Higman linearization trick (quoted in Lemma 2.2) and of a result from [2] (quoted in Lemma 2.3) on the automorphisms of $A[x, x^{-1}]$ -modules which are linear in x . A similar result has been obtained independently by Karoubi ([4]), using an L -theoretic analogue of the localization sequence of Chapter IX of [1].

Adopting the terminology of [1], we can say that each

$$V_n: (\text{rings with involution}) \rightarrow (\text{abelian groups})$$

is a contracted functor, with

$$LV_n(A) = U_n(A)$$

up to natural isomorphism. Corollary 4.4 generalizes this “Fundamental Theorem” of algebraic L -theory to describe the intermediate L -groups $V_n^Q(A[x, x^{-1}])$, as defined in Part III, for suitable subgroups $Q \subseteq \tilde{K}_1(A[x, x^{-1}])$. Corollary 4.5 identifies the “lower L -theories” of Part II with the functors

$$L^m U_n: (\text{rings with involution}) \rightarrow (\text{abelian groups}) \quad (m > 0)$$

derived from U_n . (There is an obvious analogy here with the “lower K -theories” of Chapter XII of [1],

$$K_{-m} = L^m K_0: (\text{rings}) \rightarrow (\text{abelian groups}).)$$

Corollary 4.6 describes the L -groups of polynomial extensions in several variables.

The work presented here was stimulated by a course of lectures on algebraic K -theory given by Hyman Bass at Cambridge University in the Lent Term of 1973.

§1. Contracted Functors

Let (rings) be the category of associative rings with 1, and 1-preserving ring morphisms. Let x be an invertible indeterminate over such a ring A commuting with every element of A , and define $A[x, x^{-1}]$, the ring of finite polynomials $\sum_{j=-\infty}^{\infty} x^j a_j$ in x, x^{-1} with coefficients $a_j \in A$. Let $A[x^{\pm 1}]$ be the subring of $A[x, x^{-1}]$ of poly-

nomials involving only non-negative powers of $x^{\pm 1}$. Let

$$\bar{\varepsilon}_\pm : A \rightarrow A[x^{\pm 1}], \quad \bar{E}_\pm : A[x^{\pm 1}] \rightarrow A[x, x^{-1}], \quad \bar{\varepsilon} = \bar{E}_\pm \bar{\varepsilon}_\pm : A \rightarrow A[x, x^{-1}]$$

be the inclusions, and define left inverses

$$\varepsilon_\pm : A[x^{\pm 1}] \rightarrow A, \quad \varepsilon : A[x, x^{-1}] \rightarrow A$$

for $\bar{\varepsilon}_\pm, \bar{\varepsilon}$ by $x^{\pm 1} \mapsto 1$.

A functor

$$F : (\text{rings}) \rightarrow (\text{abelian groups})$$

is *contracted* if the sequence

$$0 \rightarrow F(A) \xrightarrow{\begin{pmatrix} -\bar{\varepsilon}_+ \\ -\bar{\varepsilon}_- \end{pmatrix}} F(A[x]) \oplus F(A[x^{-1}]) \xrightarrow{(E_+ E_-)} F(A[x, x^{-1}]) \xrightarrow{B} LF(A) \rightarrow 0$$

is exact for each A , and there is given a natural right inverse

$$\bar{B} : LF(A) \rightarrow F(A[x, x^{-1}])$$

for the natural projection

$$\begin{aligned} B : F(A[x, x^{-1}]) &\rightarrow LF(A) \\ &= \text{coker}((\bar{E}_+ \bar{E}_-) : F(A[x]) \oplus F(A[x^{-1}]) \rightarrow F(A[x, x^{-1}])), \end{aligned}$$

that is $B\bar{B} = 1 : LF(A) \rightarrow LF(A)$. (This is just Definition 7.1 of Chapter XII of [1]).

LEMMA 1.1. *Let*

$$F, G : (\text{rings}) \rightarrow (\text{abelian groups})$$

be functors, and suppose given

i) a natural left inverse

$$E_+ : F(A[x, x^{-1}]) \rightarrow F(A[x])$$

for

$$\bar{E}_+ : F(A[x]) \rightarrow F(A[x, x^{-1}])$$

such that the square

$$\begin{array}{ccc} F(A[x^{-1}]) & \xrightarrow{E_-} & F(A[x, x^{-1}]) \\ \varepsilon_- \downarrow & & \downarrow E_+ \\ F(A) & \xrightarrow{\bar{\varepsilon}_+} & F(A[x]) \end{array}$$

commutes,

ii) natural morphisms

$$\bar{\eta}_+ : G(A) \rightarrow L_+ F(A) = \text{coker}(\bar{E}_+ : F(A[x]) \rightarrow F(A[x, x^{-1}]))$$

$$\eta_+ : L_+ F(A) \rightarrow G(A)$$

such that $\eta_+ \bar{\eta}_+ = 1$, and such that the square

$$\begin{array}{ccc} L_+ F(A) & \xrightarrow{\eta_+} & G(A) \\ \downarrow \alpha_+ & & \downarrow \bar{\eta}_- \\ F(A[x, x^{-1}]) & \xrightarrow{\delta_-} & L_- F(A) \end{array}$$

commutes, where

$$\Delta_+: L_+ F(A) \rightarrow F(A[x, x^{-1}])$$

is the right inverse for the natural projection

$$\delta_+: F(A[x, x^{-1}]) \rightarrow L_+ F(A)$$

induced by

$$1 - \bar{E}_+ E_+ : F(A[x, x^{-1}]) \rightarrow F(A[x, x^{-1}]),$$

and δ_- , $\bar{\eta}_-$ are defined as δ_+ , $\bar{\eta}_+$ but with x^{-1} replacing x .


Then F is a contracted functor, and

$$B = \eta_+ \delta_+ : F(A[x, x^{-1}]) \rightarrow G(A)$$

induces a natural isomorphism

$$LF(A) = \text{coker}((\bar{E}_+ \bar{E}_-): F(A[x]) \oplus F(A[x^{-1}]) \rightarrow F(A[x, x^{-1}])) \rightarrow G(A).$$

Proof. The diagrams



are commutative exact braids, where E_- , Δ_- , η_- are defined as E_+ , Δ_+ , η_+ but with x^{-1} replacing x . It follows that

$$0 \rightarrow F(A) \xrightarrow{\begin{pmatrix} \bar{\varepsilon}_+ \\ -\bar{\varepsilon}_- \end{pmatrix}} F(A[x]) \oplus F(A[x^{-1}]) \xrightarrow{(E_+ E_-)} F(A[x, x^{-1}]) \xrightarrow{B} G(A) \rightarrow 0$$

is an exact sequence, with

$$\bar{B} = \Delta_{\pm} \bar{\eta}_{\pm} : G(A) \rightarrow F(A[x, x^{-1}])$$

a natural right inverse for

$$B = \eta_{\pm} \delta_{\pm} : F(A[x, x^{-1}]) \rightarrow G(A).$$

Thus F is a contracted functor, with

$$LF(A) = G(A)$$

up to natural isomorphism. \square

(The conditions of Lemma 1.1 are necessary, as well as sufficient, for a functor to be contracted. If

$$F : (\text{rings}) \rightarrow (\text{abelian groups})$$

is a contracted functor, then

$$F(A[x, x^{-1}]) = \bar{\varepsilon}F(A) \oplus \bar{E}_+ N_+ F(A) \oplus \bar{E}_- N_- F(A) \oplus \bar{B}LF(A)$$

where

$$N_{\pm} F(A) = \ker(\varepsilon_{\pm} : F(A[x^{\pm 1}]) \rightarrow F(A)),$$

and the morphisms

$$\begin{aligned} E_+ : F(A[x, x^{-1}]) &\rightarrow F(A[x]) = \bar{\varepsilon}_+ F(A) \oplus N_+ F(A); \\ \bar{\varepsilon}(r) \oplus \bar{E}_+(s_+) \oplus \bar{E}_-(s_-) \oplus \bar{B}(t) &\mapsto \bar{\varepsilon}_+(r) \oplus s_+ \\ \bar{\eta}_+ : LF(A) &\rightarrow L_+ F(A) = \bar{E}_- N_- F(A) \oplus \bar{B}LF(A); t \mapsto 0 \oplus \bar{B}(t) \\ \eta_+ : L_+ F(A) &\rightarrow LF(A); \bar{E}_-(s_-) \oplus \bar{B}(t) \mapsto t \end{aligned}$$

satisfy the conditions of Lemma 1.1, with $G = LF$.)

§2. K-Theory of Polynomial Extensions

Let $\mathbf{P}(A)$ be the category of finitely generated (f.g.) projective left A -modules. Write $|\mathbf{P}(A)|$ for the class of objects, and $\text{Hom}_A(P, Q)$ for the additive group of

morphisms $g: P \rightarrow Q \in \mathbf{P}(A)$. A ring morphism

$$f: A \rightarrow A'$$

induces a functor

$$f: \mathbf{P}(A) \rightarrow \mathbf{P}(A'); \begin{cases} P \in |\mathbf{P}(A)| \mapsto fP = A' \otimes_A P \in |\mathbf{P}(A')| \\ g \in \text{Hom}_A(P, Q) \mapsto fg = 1 \otimes g \in \text{Hom}_{A'}(fP, fQ). \end{cases}$$

Given $P \in |\mathbf{P}(A)|$, let

$$P[x^{\pm 1}] = \bar{\varepsilon}_{\pm} P \in |\mathbf{P}(A[x^{\pm 1}])|, P_x = \bar{\varepsilon} P \in |\mathbf{P}(A[x, x^{-1}])|.$$

Defining complementary A -submodules

$$P^+ = \sum_{j=0}^{\infty} x^j P, \quad P^- = \sum_{j=-\infty}^{-1} x^j P$$

of P_x (where $x^j P = x^j \otimes P$) we shall identify

$$P^+ = P[x], \quad xP^- = P[x^{-1}]$$

in the obvious way.

Let $\mathbf{N}(A)$ be the category with objects pairs

$$(P \in |\mathbf{P}(A)|, v \in \text{Hom}_A(P, P) \text{ nilpotent})$$

and morphisms

$$f: (P, v) \rightarrow (P', v') \in \mathbf{N}(A)$$

isomorphisms $f \in \text{Hom}_A(P, P')$ such that

$$v'f = fv \in \text{Hom}_A(P, P').$$

As usual, there are defined functors

$$K_i: (\text{rings}) \rightarrow (\text{abelian groups}); \quad A \mapsto K_i(\mathbf{P}(A))$$

for $i=0,1$. Theorem 7.4 of Chapter XII of [1], the “Fundamental Theorem” of algebraic K -theory, may be stated and proved as follows:

THEOREM 2.1 *The functor K_1 is contracted, with*

$$L_+ K_1(A) = K_0 \mathbf{N}(A), \quad LK_1(A) = K_0(A)$$

up to natural isomorphism.

Proof. Given an automorphism

$$f: G_x \rightarrow G_x \in \mathbf{P}(A[x, x^{-1}]) \quad (G \in |\mathbf{P}(A)|)$$

let $F = f(G) \subseteq G_x$, and define

$$(P, v) = (G^-/x^{-N}F^-, x^{-1}) \in |\mathbf{N}(A)|$$

for $N \geq 0$ so large that $x^{-N}F^- \subseteq G^-$. Then

$$E_+: K_1(A[x, x^{-1}]) \rightarrow K_1(A[x]);$$

$$\tau(f: G_x \rightarrow G_x) \mapsto \bar{\varepsilon}_+ \tau(\varepsilon f: G \rightarrow G) \oplus \tau((1-v)^{-1}(1-xv): P^+ \rightarrow P^+)$$

is a well-defined morphism.

LEMMA 2.2 *Every element of $K_1(A[x])$ can be represented by an automorphism*

$$f = f_0 + xf_1: G^+ \rightarrow G^+ \in \mathbf{P}(A[x])$$

with $f_0, f_1 \in \text{Hom}_A(G, G)$.

Proof. Given an automorphism

$$f = f_0 + xf_1 + x^2f_2 + \cdots + x^rf_r \in \text{Hom}_{A[x]}(G^+, G^+) \quad (f_j \in \text{Hom}_A(G, G), 0 \leq j \leq r)$$

we can apply the usual Higman linearization trick (first used in the proof of Theorem 15 of [3]), the identity

$$\begin{aligned} & \begin{pmatrix} 1 & -x^{r-1} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} f & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ xf_r & 1 \end{pmatrix} \\ & = \begin{pmatrix} f_0 + xf_1 + \cdots + x^{r-1}f_{r-1} & -x^{r-1} \\ xf_r & 1 \end{pmatrix}: G^+ \oplus G^+ \rightarrow G^+ \oplus G^+ \end{aligned}$$

($r-1$) times, to obtain a representative automorphism for $\tau(f) \in K_1(A[x])$ which is linear in x (with $r=1$). \square

Given an automorphism

$$f = f_0 + xf_1 \in \text{Hom}_{A[x]}(G^+, G^+)$$

let $\gamma = (f_0 + f_1)^{-1} f_1 \in \text{Hom}_A(G, G)$. Then

$$f = (f_0 + f_1)(1 + (x-1)\gamma): G^+ \rightarrow G^+$$

and (up to isomorphism)

$$(G^-/x^{-1}f(G^-), x^{-1}) = (G^-/x^{-1}(1 + (x-1)\gamma)G^-, x^{-1}) = (G, -\gamma(1-\gamma)^{-1}) \in |\mathbf{N}(A)|.$$

It follows that

$$\begin{aligned} E_+ \bar{E}_+ \tau(f) &= \tau(f_0 + f_1 : G^+ \rightarrow G^+) \oplus \tau((1 + \gamma(1 - \gamma)^{-1})^{-1} \\ &\quad \times (1 + x\gamma(1 - \gamma)^{-1}) : G^+ \rightarrow G^+) \\ &= \tau(f_0 + f_1 : G^+ \rightarrow G^+) \oplus \tau(1 + (x - 1)\gamma : G^+ \rightarrow G^+) \\ &= \tau(f) \in K_1(A[x]). \end{aligned}$$

Thus the composite

$$K_1(A[x]) \xrightarrow{E_+} K_1(A[x, x^{-1}]) \xrightarrow{E_+} K_1(A[x])$$

is the identity. Similarly, it can be shown that the square

$$\begin{array}{ccc} K_1(A[x^{-1}]) & \xrightarrow{E_-} & K_1(A[x, x^{-1}]) \\ \varepsilon_- \downarrow & & \downarrow E_+ \\ K_1(A) & \xrightarrow{\bar{\varepsilon}_+} & K_1(A[x]) \end{array}$$

commutes.

Higman's trick also shows that every element of $K_1(A[x, x^{-1}])$ may be expressed as

$$\tau = \tau(f_0 + xf_1 : P_x \rightarrow P_x) \oplus \tau(x^N : Q_x \rightarrow Q_x) \in K_1(A[x, x^{-1}])$$

for some $P, Q \in \mathbf{P}(A)$, $f_0, f_1 \in \text{Hom}_A(P, P)$, $N \in \mathbf{Z}$.

LEMMA 2.3. *If $\gamma \in \text{Hom}_A(P, P)$ is such that*

$$1 + (x - 1)\gamma \in \text{Hom}_{A[x, x^{-1}]}(P_x, P_x)$$

is an isomorphism then there exist integers $r, s \geq 0$ such that

$$\gamma^r(1 - \gamma)^s = 0 \in \text{Hom}_A(P, P),$$

and $R = \ker \gamma^r$, $S = \ker(1 - \gamma)^s$ are complementary submodules of P , such that

$$\gamma = \begin{pmatrix} \gamma_R & 0 \\ 0 & \gamma_S \end{pmatrix} : P = R \oplus S \rightarrow P = R \oplus S$$

with $\gamma_R \in \text{Hom}_A(R, R)$, $1 - \gamma_S \in \text{Hom}_A(S, S)$ nilpotent.

Proof. See Corollary 2.4 of [2] and pp. 232–34 of [8]. \square

If $f_0, f_1 \in \text{Hom}_A(P, P)$ are such that

$$f = f_0 + xf_1 \in \text{Hom}_{A[x, x^{-1}]}(P_x, P_x)$$

is an isomorphism, then

$$ef = f_0 + f_1 \in \text{Hom}_A(P, P)$$

is an isomorphism, and $\gamma = (f_0 + f_1)^{-1} f_1 \in \text{Hom}_A(P, P)$ satisfies the hypothesis of Lemma 2.3. Hence

$$\begin{aligned} \tau(f) &= \bar{\varepsilon} \tau(f_0 + f_1 : P \rightarrow P) \oplus \tau(1 + (x-1) \gamma : P_x \rightarrow P_x) \\ &= \bar{\varepsilon} \tau(f_0 + f_1 : P \rightarrow P) \\ &\oplus \bar{E}_+ \tau(1 + (x-1) \gamma_R : R[x] \rightarrow R[x]) \\ &\oplus \bar{E}_- \tau(1 + (x^{-1}-1) (1-\gamma_S) : S[x^{-1}] \rightarrow S[x^{-1}]) \\ &\oplus \tau(x : S_x \rightarrow S_x) \in K_1(A[x, x^{-1}]) \end{aligned}$$

It is now easy to verify that

$$K_1(A[x]) \xrightarrow[E_+]{\quad} K_1(A[x, x^{-1}]) \xrightarrow[\delta_+]{\quad} K_0 \mathbf{N}(A)$$

is a direct sum system, with

$$\begin{aligned} \Delta_+ : K_0 \mathbf{N}(A) &\rightarrow K_1(A[x, x^{-1}]); [P, v] \mapsto \tau((1-v)^{-1} (x-v) : P_x \rightarrow P_x) \\ \delta_+ : K_1(A[x, x^{-1}]) &\rightarrow K_0 \mathbf{N}(A); \tau(f : G_x \rightarrow G_x) \mapsto [G^+ / x^N F^+, x] - [F^+ / x^N F^+, x] \end{aligned}$$

where $F = f(G) \subseteq G_x$ (as before) and $N \geq 0$ is so large that $x^N F^+ \subseteq G^+$, (so that, in particular,

$$\delta_+ \tau(f_0 + x f_1 : P_x \rightarrow P_x) = [S, -\gamma_S^{-1} (1-\gamma_S)] \in K_0 \mathbf{N}(A).$$

Identifying

$$L_+ K_1(A) = K_0 \mathbf{N}(A)$$

in this way, note that the morphisms

$$\begin{aligned} \eta_+ : K_0 \mathbf{N}(A) &\rightarrow K_0(A); [P, v] \mapsto [P] \\ \bar{\eta}_+ : K_0(A) &\rightarrow K_0 \mathbf{N}(A); [P] \mapsto [P, 0] \end{aligned}$$

are such that the conditions of Lemma 1.1 are satisfied. Hence

$$K_1 : (\text{rings}) \rightarrow (\text{abelian groups})$$

is a contracted functor, with

$$LK_1(A) = K_0(A)$$

up to natural isomorphism. This completes the proof of Theorem 2.1. \square

§3. Review of the Definitions of the L-Groups

Let (rings with involution) be the category of rings A (as in §1) with involution ${}^-: A \rightarrow A; a \mapsto \bar{a}$ such that

$$\bar{1} = 1, \bar{a+b} = \bar{a} + \bar{b}, \bar{ab} = \bar{b} \cdot \bar{a}, a = a \quad \text{for all } a, b \in A.$$

As in Part I it will be assumed that f.g. free A -modules have a well-defined dimension.

Given a ring with involution A define a *duality* involution

$${}^*: \mathbf{P}(A) \rightarrow \mathbf{P}(A) \begin{cases} P \in |\mathbf{P}(A)| \mapsto P^* = \text{Hom}_A(P, A), \text{ left } A\text{-action by} \\ \quad A \times P^* \rightarrow P^*; (a, p^*) \mapsto (p \mapsto p^*(p) \cdot \bar{a}) \\ f \in \text{Hom}_A(P, Q) \mapsto (f^*: Q^* \rightarrow P^*; q^* \mapsto (p \mapsto q^*(f(p)))), \end{cases}$$

using the natural isomorphisms

$$P \rightarrow P^{**}; p \mapsto (\bar{p}^* \mapsto \overline{p^*(p)}) \quad (P \in |\mathbf{P}(A)|)$$

to identify

$${}^{**} = 1: \mathbf{P}(A) \rightarrow \mathbf{P}(A).$$

An ε -hermitian product (over A) is a morphism

$$\theta: Q \rightarrow Q^* \in \mathbf{P}(A)$$

such that

$$\theta^* = \varepsilon \theta \in \text{Hom}_A(Q, Q^*),$$

where $\varepsilon = \pm 1$. A \pm form (over A) is a pair

$$(Q \in |\mathbf{P}(A)|, \varphi \in \text{Hom}_A(Q, Q^*)),$$

and

$$\theta = \varphi \pm \varphi^* \in \text{Hom}_A(Q, Q^*)$$

is the *associated* \pm hermitian product. An *isomorphism* of \pm forms

$$(f, \chi): (Q, \varphi) \rightarrow (Q', \varphi')$$

is an isomorphism $f \in \text{Hom}_A(Q, Q')$ together with a morphism $\chi \in \text{Hom}_A(Q, Q^*)$ such that

$$f^* \varphi' f - \varphi = \chi \mp \chi^* \in \text{Hom}_A(Q, Q^*).$$

Such an isomorphism preserves the associated \pm hermitian products, in that

$$f^*(\varphi' \pm \varphi'^*)f = (\varphi \pm \varphi^*) \in \text{Hom}_A(Q, Q^*).$$

A \pm form (Q, φ) is *non-singular* if the associated \pm hermitian product $(\varphi \pm \varphi^*) \in \text{Hom}_A(Q, Q^*)$ is an isomorphism. The *hamiltonian* \pm form on $P \in |\mathbf{P}(A)|$,

$$H\pm(P) = (P \oplus P^*, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix})$$

is non-singular. A *sublagrangian* of a non-singular \pm form (Q, φ) is a direct summand L of Q such that

$$j^*\varphi j = \lambda \mp \lambda^* \in \text{Hom}_A(L, L^*)$$

for some $\lambda \in \text{Hom}_A(L, L^*)$, denoting by $j \in \text{Hom}_A(L, Q)$ the inclusion. It was shown in Theorem 1.1 of Part I that if L is a sublagrangian of (Q, φ) there is defined a non-singular \pm form $(L^\perp/L, \hat{\varphi})$ on a direct complement L^\perp/L to L in the *annihilator* of L in (Q, φ) ,

$$L^\perp = \ker(j^*(\varphi \pm \varphi^*): Q \rightarrow L^*),$$

and that there is defined an isomorphism of \pm forms

$$(f, \chi): (Q, \varphi) \rightarrow H\pm(L) \oplus (L^\perp/L, \hat{\varphi})$$

with f the identity on $L^\perp = L \oplus L^\perp/L$. A *lagrangian* is a sublagrangian L such that

$$L^\perp = L,$$

in which case there is defined an isomorphism of \pm forms

$$(f, \chi): (Q, \varphi) \rightarrow H\pm(L).$$

A \pm *formation* (over A), $(Q, \varphi; F, G)$, is a triple consisting of

- i) a non-singular \pm form over A , (Q, φ) ,
- ii) a lagrangian F of (Q, φ) ,
- iii) a sublagrangian G of (Q, φ) .

An *isomorphism* of \pm formations

$$(f, \chi): (Q, \varphi; F, G) \rightarrow (Q', \varphi'; F', G')$$

is an isomorphism of \pm forms

$$(f, \chi): (Q, \varphi) \rightarrow (Q', \varphi')$$

such that $f(F)=F', f(G)=G'$. A *stable isomorphism* of \pm formations

$$[f, \chi]: (Q, \varphi; F, G) \rightarrow (Q', \varphi'; F', G')$$

is an isomorphism of \pm formations

$$(f, \chi): (Q, \varphi; F, G) \oplus (H \pm (P); P, P^*) \rightarrow (Q', \varphi'; F', G') \oplus (H \pm (P'); P', P'^*)$$

defined for some $P, P' \in |\mathbf{P}(A)|$.

Let $T \subseteq \tilde{K}_0(A) = \text{coker}(K_0(\mathbf{Z}) \rightarrow K_0(A))$ be a subgroup invariant under the duality involution

$$*: \tilde{K}_0(A) \rightarrow \tilde{K}_0(A); [P] \mapsto [P^*] \quad (\text{that is, } * (T) = T).$$

For $n \pmod{4}$ define the abelian monoid $X_n^T(A)$ of $\begin{cases} \text{isomorphism} \\ \text{stable isomorphism} \end{cases}$

classes of $\begin{cases} \pm \text{ forms } (Q, \varphi) \\ \pm \text{ formations } (Q, \varphi; F, G) \end{cases}$ over A such that the projective class

$\begin{cases} [Q] \\ [G] - [F^*] \end{cases}$ lies in $T \subseteq \tilde{K}_0(A)$, under the direct sum \oplus , with $\pm = (-)^i$ if $n = \begin{cases} 2i \\ 2i+1 \end{cases}$.

The monoid morphisms

$$\partial^T: X_n^T(A) \rightarrow X_{n-1}^T(A); \begin{cases} (Q, \varphi) \mapsto (H_{\mp}(Q); Q, \Gamma_{(Q, \varphi)}) \\ (Q, \varphi; F, G) \mapsto (G^{\perp}/G, \hat{\varphi}) \end{cases} \quad n = \begin{cases} 2i \\ 2i+1 \end{cases}$$

are such that $(\partial^T)^2 = 0$, where

$$\Gamma_{(Q, \varphi)} = \{(x, (\varphi \pm \varphi^*) x) \mid x \in Q\} \subseteq Q \oplus Q^*.$$

Define an equivalence relation \sim on $\ker(\partial^T: X_n^T(A) \rightarrow X_{n-1}^T(A))$ by $z_1 \sim z_2$ if there exist $b_1, b_2 \in X_{n+1}^T(A)$ such that $z_1 \oplus \partial^T b_1 = z_2 \oplus \partial^T b_2 \in X_n^T(A)$. It was shown in Theorem 2.1 of Part III that the quotient monoids

$$U_n^T(A) = \ker(\partial^T: X_n^T(A) \rightarrow X_{n-1}^T(A)) / \overline{\text{im}(\partial^T: X_{n+1}^T(A) \rightarrow X_n^T(A))}$$

of equivalence classes are abelian groups, generalizing the definitions in Part I of

$$U_n(A) = U_n^{K_0(A)}(A), \quad V_n(A) = U_n^{\{0\}}(A).$$

Theorem 2.3 of Part III established an exact sequence

$$\cdots \rightarrow H^{n+1}(T'/T) \rightarrow U_n^T(A) \rightarrow U_n^{T'}(A) \rightarrow H^n(T'/T) \rightarrow U_{n-1}^T(A) \rightarrow \cdots$$

for $*$ -invariant subgroups $T \subseteq T' \subseteq \tilde{K}_0(A)$, where

$$H^n(G) = \{g \in G \mid g^* = (-)^n g\} / \{h + (-)^n h^* \mid h \in G\}$$

are the Tate cohomology groups (abelian, of exponent 2).

There are analogous definitions and results for L -groups associated with subgroups $R \subseteq \tilde{K}_1(A) = \text{coker}(K_1(\mathbf{Z}) \rightarrow K_1(A))$ invariant under the duality involution

$$*: \tilde{K}_1(A) \rightarrow \tilde{K}_1(A); \tau(f: \tilde{P} \rightarrow \tilde{Q}) \mapsto \tau(f^*: \tilde{Q}^* \rightarrow \tilde{P}^*)$$

denoting by \tilde{P} a f.g. free A -module P with a prescribed base, and by \tilde{P}^* the dual based A -module.

A *based \pm form* (\tilde{Q}, φ) is a \pm form (Q, φ) on a based A -module \tilde{Q} . The *torsion* of a based \pm form (\tilde{Q}, φ) is

$$\tau(\tilde{Q}, \varphi) = \begin{cases} \tau(\varphi \pm \varphi^*: \tilde{Q} \rightarrow \tilde{Q}^*) \in \tilde{K}_1(A) & \text{if } (Q, \varphi) \text{ is non-singular} \\ 0 \in \tilde{K}_1(A) & \text{otherwise.} \end{cases}$$

An *R-isomorphism* of based \pm forms

$$(f, \chi): (\tilde{Q}, \varphi) \rightarrow (\tilde{Q}', \varphi')$$

is an isomorphism of the underlying forms

$$(f, \chi): (Q, \varphi) \rightarrow (Q', \varphi')$$

such that

$$\tau(f: \tilde{Q} \rightarrow \tilde{Q}') \in R \subseteq \tilde{K}_1(A).$$

A *based \pm formation* $(Q, \varphi; \tilde{F}, \tilde{G})$ is a \pm formation $(Q, \varphi; F, G)$ with bases for F, G and G^\perp/G . The *torsion* $\tau(Q, \varphi; \tilde{F}, \tilde{G}) \in \tilde{K}_1(A)$ of a based \pm formation is the torsion of the isomorphism

$$f: \tilde{F} \oplus \tilde{F}^* \rightarrow \tilde{G} \oplus \tilde{G}^* \oplus \underline{G^\perp/G}$$

in the isomorphism of \pm forms

$$(f, \chi): H \pm (F) \rightarrow H \pm (G) \oplus (G^\perp/G, \hat{\varphi})$$

given by Theorem 1.1 of Part I. An *R-isomorphism* of based \pm formations

$$(f, \chi): (Q, \varphi; \tilde{F}, \tilde{G}) \rightarrow (Q', \varphi'; \tilde{F}', \tilde{G}')$$

is an isomorphism of the underlying \pm formations such that the restrictions

$$\tilde{F} \rightarrow \tilde{F}', \tilde{G} \rightarrow \tilde{G}', \underline{G^\perp/G} \rightarrow \underline{G'^\perp/G'}$$

of f have torsions in $R \subseteq \tilde{K}_1(A)$. A *stable R-isomorphism* of based \pm formations

$$[f, \chi]: (Q, \varphi; F, G) \rightarrow (Q', \varphi'; \tilde{F}', \tilde{G}')$$

is an R -isomorphism

$$(f, \chi): (Q, \varphi; \underline{F}, \underline{G}) \oplus (H \pm (P); \underline{P}, \underline{P}^*) \rightarrow (Q', \varphi'; \underline{F}', \underline{G}') \oplus (H \pm (P'); \underline{P}', \underline{P}'^*)$$

defined for some based A -modules $\underline{P}, \underline{P}'$.

For $n \pmod{4}$ define the abelian monoid $Y_n^R(A)$ of $\begin{cases} R\text{-isomorphism} \\ \text{stable } R\text{-isomorphism} \end{cases}$ classes of based $\begin{cases} \pm \text{ forms} \\ \pm \text{ formations} \end{cases}$ over A with torsion in $R \subseteq \tilde{K}_1(A)$, under the direct sum \oplus , with $\pm = (-)^i$ if $n = \begin{cases} 2i \\ 2i+1 \end{cases}$. The monoid morphisms

$$\partial^R: Y_n^R(A) \rightarrow Y_{n-1}^R(A); \begin{cases} (Q, \varphi) \mapsto (H \mp (Q); \underline{Q}, \Gamma_{(Q, \varphi)}) \\ (Q, \varphi; \underline{F}, \underline{G}) \mapsto (\underline{G}^\perp / \underline{G}, \hat{\varphi}) \end{cases} \quad n = \begin{cases} 2i \\ 2i+1 \end{cases}$$

are such that $(\partial^R)^2 = 0$, and the quotient monoids

$$V_n^R(A) = \ker(\partial^R: Y_n^R(A) \rightarrow Y_{n-1}^R(A)) / \overline{\text{im}(\partial^R: Y_{n+1}^R(A) \rightarrow Y_n^R(A))}$$

are abelian groups (by Theorem 3.1 of Part III) generalizing the definitions in Part I of

$$V_n(A) = V_n^{\tilde{K}_1(A)}(A) (= U_n^{\{0\}}(A)), \quad W_n(A) = V_n^{\{0\}}(A).$$

Theorem 3.3 in Part III established an exact sequence

$$\cdots \rightarrow H^{n+1}(R'/R) \rightarrow V_n^R(A) \rightarrow V_n^{R'}(A) \rightarrow H^n(R'/R) \rightarrow V_{n-1}^R(A) \rightarrow \cdots$$

for $*$ -invariant subgroups $R \subseteq R' \subseteq \tilde{K}_1(A)$.

A morphism of rings with involution

$$f: A \rightarrow A'$$

such that $f(T) \subseteq T'$ (for some $*$ -invariant subgroups $T \subseteq \tilde{K}_0(A)$, $T' \subseteq \tilde{K}_0(A')$) induces abelian group morphisms

$$f: U_n^T(A) \rightarrow U_n^{T'}(A'); \begin{cases} (Q, \varphi) \mapsto (fQ, f\varphi) \\ (Q, \varphi; F, G) \mapsto (fQ, f\varphi; fF, fG) \end{cases} \quad n = \begin{cases} 2i \\ 2i+1 \end{cases}.$$

Similarly, if $f(R) \subseteq R'$ (for $*$ -invariant subgroups $R \subseteq \tilde{K}_1(A)$, $R' \subseteq \tilde{K}_1(A')$) there are induced morphisms

$$f: V_n^R(A) \rightarrow V_n^{R'}(A') \quad (n \pmod{4}).$$

§4. L-Theory of Polynomial Extensions

Given a ring with involution A and an indeterminate x over A commuting with

every element of A extend the involution on A to the involution

$$- : A[x, x^{-1}] \rightarrow A[x, x^{-1}]; \quad \sum_{j=-\infty}^{\infty} x^j a_j \mapsto \sum_{j=-\infty}^{\infty} x^j \bar{a}_j$$

on $A[x, x^{-1}]$. This restricts to involutions on the subrings $A[x]$, $A[x^{-1}]$ of $A[x, x^{-1}]$. F. g, free $A[x]$ -modules have well-defined dimension, as do those over $A[x^{-1}]$, $A[x, x^{-1}]$. Thus the rings with involution $A[x^{\pm 1}]$, $A[x, x^{-1}]$ satisfy the conditions imposed on A in §3.

Call a functor

$$F: (\text{rings with involution}) \rightarrow (\text{abelian groups})$$

contracted if the sequence

$$0 \rightarrow F(A) \xrightarrow{(-\bar{e}_+)} F(A[x]) \oplus F(A[x^{-1}]) \xrightarrow{(E_+ E_-)} F(A[x, x^{-1}]) \xrightarrow{B} LF(A) \rightarrow 0$$

is exact for every ring with involution A and there is given a natural right inverse

$$\bar{B}: LF(A) \rightarrow F(A[x, x^{-1}])$$

for the natural projection

$$\begin{aligned} B: F(A[x, x^{-1}]) &\rightarrow LF(A) \\ &= \text{coker}((\bar{E}_+ \bar{E}_-): F(A[x]) \oplus F(A[x^{-1}]) \rightarrow F(A[x, x^{-1}])). \end{aligned}$$

The obvious analogue to Lemma 1.1 holds for functors

$$(\text{rings with involution}) \rightarrow (\text{abelian groups})$$

as does the following analogue of Theorem 2.1 for the L -theoretic functors of §3:

THEOREM 4.1. *Each of the functors*

$$V_n: (\text{rings with involution}) \rightarrow (\text{abelian groups}) \quad (n \pmod{4})$$

is contracted, with

$$LV_n(A) = U_n(A), \quad L_{\pm} V_n(A) = U_n^{K_0(A)}(A[x^{\mp 1}])$$

up to natural isomorphism, where $\tilde{K}_0(A) \equiv \bar{e}_{\mp} \tilde{K}_0(A) \subseteq \tilde{K}_0(A[x^{\mp 1}])$. \square

The proof of Theorem 4.1 in the case $n=2i$ will be similar to the proof of Theorem 2.1. The case $n=2i+1$ will follow by an application of the results of Part II on the L -theory of Laurent extensions (that is, of the ring $A[x, x^{-1}]$ with involution by $\bar{x}=x^{-1}$).

Recall from Part II that a *modular A-base* of an $A[x, x^{-1}]$ -module Q is an A -submodule Q_0 of Q such that every element q of Q has a unique expression as

$$q = \sum_{j=-\infty}^{\infty} x^j q_j \quad (q_j \in Q_0, \{j \mid q_j \neq 0\} \text{ finite}),$$

so that $Q = A[x, x^{-1}] \otimes_A Q_0$ up to $A[x, x^{-1}]$ -module isomorphism. For example the A -modules generated by the bases of free $A[x, x^{-1}]$ -modules are modular A -bases.

Define a morphism

$$\begin{aligned} \delta_+ : V_{2i}(A[x, x^{-1}]) &\rightarrow U_{2i}^{K_0(A)}(A[x^{-1}]); \\ (Q, \varphi) &\mapsto (P[x^{-1}], [\varphi]_{-1} - x^{-1}[\varphi]_{-2}) \end{aligned}$$

by choosing a modular A -base Q_0 for Q (which is a f.g. free $A[x, x^{-1}]$ -module) and an integer $N \geq 0$ so large that

$$(\varphi \pm \varphi^*) (x^N Q_0^+) \subseteq x^{-N} Q_0^{*+} \quad (\pm = (-)^i),$$

defining

$$P = x^N Q_0^- \cap (\varphi \pm \varphi^*)^{-1} (x^{-N} Q_0^{*+}) \in |\mathbf{P}(A)|,$$

with $[\varphi]_j \in \text{Hom}_A(P, P^*)$ given by

$$[\varphi]_j(y)(y') = a_j \in A \quad (y, y' \in P, j \in \mathbf{Z})$$

if

$$\varphi(y)(y') = \sum_{j=-\infty}^{\infty} x^j a_j \in A[x, x^{-1}] \quad (a_j \in A),$$

and writing $P[x^{-1}]$ for $\bar{\epsilon}_- P = A[x^{-1}] \otimes_A P \in |\mathbf{P}(A[x^{-1}])|$.

The A -module isomorphism

$$[\varphi \pm \varphi^*]_{-1} : Q \rightarrow Q^*$$

may be expressed as

$$[\varphi \pm \varphi^*]_{-1} = \begin{pmatrix} [\varphi]_{-1} \pm ([\varphi]_{-1})^* & 0 & 0 \\ 0 & 0 & 1 \\ 0 & \pm 1 & 0 \end{pmatrix} : P \oplus L \oplus L^* \rightarrow P^* \oplus L^* \oplus L$$

where $L = (\varphi \pm \varphi^*)^{-1} (x^{-N} Q_0^{*-})$, $L^* = x^N Q_0^+ \subseteq Q$, so that $(P, [\varphi]_{-1})$ is a non-singular \pm form over A .

For any $y, y' \in P$

$$\begin{aligned} [\varphi \pm \varphi^*]_{-2}(y)(y') &= [\varphi \pm \varphi^*]_{-1}(xy)(y') \\ &= [\varphi \pm \varphi^*]_{-1}(xy - x^N y_{N-1})(y') \in A, \end{aligned}$$

where $y_{N-1} \in Q_0$ is such that

$$y - x^{N-1} y_{N-1} \in x^{N-1} Q_0^- \cap (\varphi \pm \varphi^*)^{-1} (x^{-N-1} Q_0^*) = x^{-1} P.$$

Thus

$$(P, ([\varphi \pm \varphi^*]_{-1})^{-1} ([\varphi \pm \varphi^*]_{-2})) = ((\varphi \pm \varphi^*)^{-1} (x^{-N} Q_0^{*+}) / x^N Q_0^+, x) \in |\mathbf{N}(A)|,$$

and $(P[x^{-1}], [\varphi]_{-1} - x^{-1} [\varphi]_{-2})$ is a non-singular \pm form over $A[x^{-1}]$.

Suppose that Q'_0 is a different modular A -base of Q . Let $M \geq 0$ be so large that

$$Q'_0 \subseteq \sum_{j=-M}^M x^j Q_0, \quad Q_0 \subseteq \sum_{j=-M}^M x^j Q'_0.$$

Then $N' = N + M$ is so large that

$$(\varphi \pm \varphi^*) (x^{N'} Q_0'^{*+}) \subseteq x^{-N'} Q_0'^{*+},$$

and

$$\begin{aligned} P' &= x^{N'} Q_0'^{-} \cap (\varphi \pm \varphi^*)^{-1} (x^{-N'} Q_0'^{*+}) \quad (\text{definition}) \\ &= x^N (x^M Q_0'^{-} \cap Q_0^+) \oplus P \oplus x^{-N} (\varphi \pm \varphi^*)^{-1} (Q_0'^{-} \cap x^{-M} Q_0'^{*+}). \end{aligned}$$

Now

$$L = (x^N (x^M Q_0'^{-} \cap Q_0^+)) [x^{-1}] \subseteq P'[x^{-1}]$$

is a sublagrangian of $(P'[x^{-1}], [\varphi]_{-1} - x^{-1} [\varphi]_{-2})$ with $L^\perp / L = P[x^{-1}]$, so that

$$\begin{aligned} (P'[x^{-1}], [\varphi]_{-1} - x^{-1} [\varphi]_{-2}) &= (P[x^{-1}], [\varphi]_{-1} - x^{-1} [\varphi]_{-2}) \oplus H_\pm(L) \\ &= (P[x^{-1}], [\varphi]_{-1} - x^{-1} [\varphi]_{-2}) \in U_{2i}^{K_0(A)}(A[x^{-1}]). \end{aligned}$$

Thus the choice of N and Q_0 is immaterial to the definition of δ_+ .

Finally, suppose that

$$(Q, \varphi) = \bar{E}_+ (Q_0^+, \varphi_0) \in V_{2i}(A[x, x^{-1}])$$

for some $(Q_0^+, \varphi_0) \in V_{2i}(A[x])$. Then we can choose $N = 0$, and

$$\delta_+ (Q, \varphi) = 0 \in U_{2i}^{K_0(A)}(A[x^{-1}]).$$

Hence the morphism

$$\delta_+ : V_{2i}(A[x, x^{-1}]) \rightarrow U_{2i}^{K_0(A)}(A[x^{-1}])$$

is well-defined, and such that the composite

$$V_{2i}(A[x]) \xrightarrow{E_+} V_{2i}(A[x, x^{-1}]) \xrightarrow{\delta_+} U_{2i}^{K_0(A)}(A[x^{-1}])$$

is zero. Before going on to show that this sequence is in fact split exact, we need an L -theoretic analogue of Lemma 2.2 (the Higman linearization trick):

LEMMA 4.2. *Every element of $U_{2i}^{K_0(A)}(A[x])$ (resp. $V_{2i}(A[x, x^{-1}])$) can be represented by a linear \pm form, $(Q^+, \varphi_0 + x\varphi_1)$ over $A[x]$ (resp. $(Q_x, \varphi_0 + x\varphi_1)$ over $A[x, x^{-1}]$) where $\varphi_0, \varphi_1 \in \text{Hom}_A(Q, Q^*)$.*

Proof. Given $(Q^+, \varphi) \in U_{2i}^{K_0(A)}(A[x])$, let

$$\varphi = \sum_{j=0}^N x^j \varphi_j \text{Hom}_{A[x]}(Q^+, Q^{*+}) \quad (\varphi_j \in \text{Hom}_A(Q, Q^*)),$$

and suppose $N > 1$. Now

$$\begin{aligned} & \left(\left(\begin{pmatrix} 1 & 0 & 0 \\ -x & 1 & 0 \\ \pm x^{N-1} \varphi_N^* & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & -x^{N-1} \varphi_N & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right) \right. \\ & \left. : (Q^+, \varphi) \oplus H_{\pm}(Q^+) \rightarrow \left(Q^+ \oplus Q^+ \oplus Q^{*+}, \begin{pmatrix} \varphi - x^N \varphi_N & -x^{N-1} \varphi_N & x \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \right) \right) \end{aligned}$$

is an isomorphism of \pm forms over $A[x]$, so that

$$(Q'^+, \varphi') = (Q^+, \varphi) \in U_{2i}^{K_0(A)}(A[x])$$

with $Q' = Q \oplus Q \oplus Q^*$ such that

$$\varphi' = \sum_{j=0}^{N-1} x^j \varphi'_j \in \text{Hom}_{A[x]}(Q'^+, Q'^{*+}) \quad (\varphi'_j \in \text{Hom}_A(Q', Q'^*)).$$

Iterating this procedure $(N-1)$ times we obtain a representative for

$$(Q^+, \varphi) \in U_{2i}^{K_0(A)}(A[x]) \text{ with } N=1.$$

The same method works for elements $(Q_x, \varphi) \in V_{2i}(A[x, x^{-1}])$ provided we can assume that

$$(\varphi \pm \varphi^*) (Q^+) \subseteq Q^{*+}.$$

Choosing $N \geq 0$ so large that

$$(\varphi \pm \varphi^*) (x^N Q^+) \subseteq x^{-N} Q^{*+},$$

note that

$$(x^N, 0) : (Q_x, \varphi' = x^{2N} \varphi) \rightarrow (Q_x, \varphi)$$

as an isomorphism of \pm forms over $A[x, x^{-1}]$, so that

$$(Q_x, \varphi') = (Q_x, \varphi) \in V_{2i}(A[x, x^{-1}]),$$

and that

$$(\varphi' \pm \varphi'^*) (Q^+) \subseteq Q^{*+}. \quad \square$$

The morphism

$$\begin{aligned} \Delta_+ : U_{2i}^{K_0(A)}(A[x^{-1}]) &\rightarrow V_{2i}(A[x, x^{-1}]); \\ (Q[x^{-1}], \varphi) &\mapsto (Q_x, x\varphi) \oplus \tilde{\varepsilon}\varepsilon_-(Q[x^{-1}], -\varphi) \oplus H_{\pm}(-Q_x) \end{aligned}$$

is clearly well-defined, with $-Q \in |\mathbf{P}(A)|$ such that $Q \oplus -Q$ is f.g. free.

The composite

$$U_{2i}^{K_0(A)}(A[x^{-1}]) \xrightarrow{\Delta_+} V_{2i}(A[x, x^{-1}]) \xrightarrow{\delta_+} U_{2i}^{K_0(A)}(A[x^{-1}])$$

is the identity: by Lemma 4.2 it is sufficient to consider $\delta_+ \Delta_+(Q[x^{-1}], \varphi)$ with

$$\varphi = \varphi_0 + x^{-1}\varphi_{-1} \in \text{Hom}_{A[x^{-1}]}(Q[x^{-1}], Q^*[x^{-1}]) \quad (\varphi_0, \varphi_{-1} \in \text{Hom}_A(Q, Q^*)),$$

and

$$\begin{aligned} \delta_+ \Delta_+(Q[x^{-1}], \varphi_0 + x^{-1}\varphi_{-1}) &= \delta_+((Q_x, x\varphi_0 + \varphi_{-1}) \oplus (Q_x, -(\varphi_0 + \varphi_{-1})) \oplus H_{\pm}(-Q_x)) \\ &= ((Q \cap (x(\varphi_0 \pm \varphi_0^*) + (\varphi_{-1} \pm \varphi_{-1}^*))^{-1}(Q^{*+})) [x^{-1}], \\ &[x\varphi_0 + \varphi_{-1}]_{-1} - x^{-1}[x\varphi_0 + \varphi_{-1}]_{-2}) \\ &= ((1 + x^{-1}\gamma)^{-1}(x^{-1}Q), [x\varphi_0 + \varphi_{-1}]_{-1} - x^{-1}[x\varphi_0 + \varphi_{-1}]_{-2}) \end{aligned}$$

where $\gamma = (\varphi_0 \pm \varphi_0^*)^{-1}(\varphi_{-1} \pm \varphi_{-1}^*) \in \text{Hom}_A(Q, Q)$ is nilpotent. Now

$$(1 + x^{-1}\gamma)^{-1} = \sum_{j=0}^{\infty} (-)^j x^{-j} \gamma^j \in \text{Hom}_{A[x^{-1}]}(Q[x^{-1}], Q[x^{-1}]),$$

so that

$$\begin{aligned} [x\varphi_0 + \varphi_{-1}]_j (1 + y^{-1}\gamma)^{-1} (x^{-1}y) (1 + x^{-1}\gamma)^{-1} (x^{-1}y') \\ = \begin{cases} \varphi_0(y)(y') & \text{if } j = 1 \\ (\varphi_{-1} - \varphi_0\gamma - \gamma^*\varphi_0)(y)(y') & \text{if } j = -2 \end{cases} \quad (y, y' \in Q), \end{aligned}$$

and

$$\varphi_{-1} - \varphi_0\gamma - \gamma^*\varphi_0 = -\varphi_{-1} + \chi \mp \chi^* \in \text{Hom}_A(Q, Q^*),$$

where $\chi = \varphi_{-1} - \gamma^*\varphi_0 \in \text{Hom}_A(Q, Q^*)$. Thus

$$\begin{aligned} \delta_+ \Delta_+(Q[x^{-1}], \varphi_0 + x^{-1}\varphi_{-1}) &= (Q[x^{-1}], \varphi_0 + x^{-1}(\varphi_{-1} - (\chi \mp \chi^*))) \\ &= (Q[x^{-1}], \varphi_0 + x^{-1}\varphi_{-1}) \in U_{2i}^{K_0(A)}(A[x^{-1}]) \end{aligned}$$

and

$$\delta_+ \Delta_+ = 1 : U_{2i}^{K_0(A)}(A[x^{-1}]) \rightarrow U_{2i}^{K_0(A)}(A[x^{-1}]).$$

It is therefore sufficient to prove that $V_{2i}(A[x, x^{-1}])$ is generated by the images of $\bar{E}_+ : V_{2i}(A[x]) \rightarrow V_{2i}(A[x, x^{-1}])$, $\Delta_+ : U_{2i}^{K_0(A)}(A[x^{-1}]) \rightarrow V_{2i}(A[x, x^{-1}])$ for the exactness of

$$V_{2i}(A[x]) \xrightarrow{E_+} V_{2i}(A[x, x^{-1}]) \xrightarrow{\delta_+} U_{2i}^{K_0(A)}(A[x^{-1}]).$$

We shall do this using the following L-theoretic analogue of Lemma 2.3:

LEMMA 4.3. *Let (Q_x, φ) be a non-singular \pm form over $A[x, x^{-1}]$ such that*

$$\varphi = \mu + (x-1)v \in \text{Hom}_{A[x, x^{-1}]}(Q_x, Q_x^*) \quad (\mu, v \in \text{Hom}_A(Q, Q^*)).$$

Then (Q_x, φ) is isomorphic to the sum

$$(R_x, \mu_R + (x-1)v_R) \oplus (S_x, \mu_S + (x-1)v_S)$$

of non-singular \pm forms over $A[x, x^{-1}]$ such that

$$(R[x], \mu_R + (x-1)v_R)$$

is a non-singular \pm form over $A[x]$, and

$$(S[x^{-1}], x^{-1}(\mu_S + (x-1)v_S))$$

is a non-singular \pm form over $A[x^{-1}]$.

Proof. The invertibility of

$$\varphi \pm \varphi^* = (\mu \pm \mu^*) + (x-1)(v \pm v^*) \in \text{Hom}_{A[x, x^{-1}]}(Q_x, Q_x^*)$$

implies that

$$\varepsilon(\varphi \pm \varphi^*) = \mu \pm \mu^* \in \text{Hom}_A(Q, Q^*)$$

$$(\mu \pm \mu^*)^{-1}(\varphi \pm \varphi^*) = 1 + (x-1)\gamma \in \text{Hom}_{A[x, x^{-1}]}(Q_x, Q_x)$$

are isomorphisms, where

$$\gamma = (\mu \pm \mu^*)^{-1}(v \pm v^*) \in \text{Hom}_A(Q, Q).$$

Hence, by Lemma 2.3,

$$\gamma = \begin{pmatrix} \gamma_R & 0 \\ 0 & \gamma_S \end{pmatrix} : Q = R \oplus S \rightarrow Q = R \oplus S$$

with $\gamma_R \in \text{Hom}_A(R, R)$, $1 - \gamma_S \in \text{Hom}_A(S, S)$ nilpotent.

Adding on some \mp hermitian products of type $\chi \mp \chi^* \in \text{Hom}_A(Q, Q^*)$ to μ and ν if necessary, it may be assumed that $\mu(R)(S) = 0$, $\nu(R)(S) = 0$. Let

$$\mu = \begin{pmatrix} \mu_R & \mu_{RS} \\ 0 & \mu_S \end{pmatrix} : R \oplus S \rightarrow R^* \oplus S^*, \quad \nu = \begin{pmatrix} \nu_R & \nu_{RS} \\ 0 & \nu_S \end{pmatrix} : R \oplus S \rightarrow R^* \oplus S^*$$

so that

$$\begin{pmatrix} \mu_R \pm \mu_R^* & \mu_{RS} \\ \pm \mu_{RS}^* & \mu_S \pm \mu_S^* \end{pmatrix} \begin{pmatrix} \gamma_R & 0 \\ 0 & \gamma_S \end{pmatrix} = \begin{pmatrix} \nu_R \pm \nu_R^* & \nu_{RS} \\ \pm \nu_{RS}^* & \nu_S \pm \nu_S^* \end{pmatrix} : R \oplus S \rightarrow R^* \oplus S^*.$$

Working as in the calculation of $\delta_+ \Delta_+$ above,

$$\begin{aligned} \delta_+(Q_x, \varphi) &= ((Q^- \cap (\varphi \pm \varphi^*)^{-1}(Q^{*+})) [x^{-1}], [\varphi]_{-1} - x^{-1} [\varphi]_{-2}) \\ &= ((1 + (x-1) \gamma_S)^{-1}(S) [x^{-1}], [\mu_S + (x-1) \nu_S]_{-1} - x^{-1} [\mu_S + (x-1) \nu_S]_{-2}) \\ &= (S [x^{-1}], x^{-1} (\mu_S + (x-1) \nu_S)) \in U_{2i}^{K_0(A)}(A [x^{-1}]). \end{aligned}$$

Thus $\varepsilon_- \delta_+(Q_x, \varphi) = (S, \mu_S)$ is a non-singular \pm form over A , and hence so is (S, ν_S) , because

$$(\nu_S \pm \nu_S^*) = (\mu_S \pm \mu_S^*) \gamma_S \in \text{Hom}_A(S, S^*)$$

and $\gamma_S \in \text{Hom}_A(S, S)$ is an isomorphism (being unipotent). Let

$$\begin{aligned} g &= \pm(\nu_S \pm \nu_S^*)^{-1} \nu_{RS}^* \in \text{Hom}_A(R, S) \\ \mu' &= \begin{pmatrix} \mu'_R = \mu_R - g^* \mu_S g & 0 \\ 0 & \mu_S \end{pmatrix} : R \oplus S \rightarrow R^* \oplus S^* \\ \nu' &= \begin{pmatrix} \nu'_R = \nu_R - g^* \nu_S g & 0 \\ 0 & \nu_S \end{pmatrix} : R \oplus S \rightarrow R^* \oplus S^*. \end{aligned}$$

Now

$$\begin{aligned} (f, \chi) &= \left(\begin{pmatrix} 1 & 0 \\ g & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ (\mu_S + (x-1) \nu_S) g & 0 \end{pmatrix} \right) \\ : (Q_x, \varphi) &= (R_x \oplus S_x, \mu + (x-1) \nu) \rightarrow (Q_x, \varphi') = (R_x \oplus S_x, \mu' + (x-1) \nu') \end{aligned}$$

is an isomorphism of \pm forms over $A[x, x^{-1}]$. It follows that

$$f^*(\varphi' \pm \varphi'^*) f = (\varphi \pm \varphi^*) \in \text{Hom}_{A[x, x^{-1}]}(Q_x, Q_x^*)$$

and as f is defined over A

$$\begin{aligned} f^*(\mu' \pm \mu'^*) f &= (\mu \pm \mu^*) \in \text{Hom}_A(Q, Q^*) \\ f^*(\nu' \pm \nu'^*) f &= (\nu \pm \nu^*) \in \text{Hom}_A(Q, Q^*). \end{aligned}$$

Defining

$$\gamma' = (\mu' \pm \mu'^*)^{-1} (v' \pm v'^*) = \begin{pmatrix} \gamma'_R = (\mu'_R \pm \mu'^*_R)^{-1} (v_R \pm v^*_R) & 0 \\ 0 & \gamma_S \end{pmatrix} : R \oplus S \rightarrow R \oplus S,$$

we have that

$$\gamma' = f\gamma f^{-1} = \begin{pmatrix} 1 & 0 \\ g & 1 \end{pmatrix} \begin{pmatrix} \gamma_R & 0 \\ 0 & \gamma_S \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -g & 1 \end{pmatrix} = \begin{pmatrix} \gamma_R & 0 \\ g\gamma_R - \gamma_S g & \gamma_S \end{pmatrix} : R \oplus S \rightarrow R \oplus S.$$

Hence

$$\gamma'_R = \gamma_R \in \text{Hom}_A(R, R)$$

is nilpotent, and $(R[x], \mu'_R + (x-1)v'_R)$ is a non-singular \pm form over $A[x]$. This completes the proof of Lemma 4.3. \square

Given $(Q_x, \varphi) \in V_{2i}(A[x, x^{-1}])$ it may be assumed, by Lemma 4.2, that $\varphi = \mu + (x-1)v \in \text{Hom}_{A[x, x^{-1}]}(Q_x, Q_x^*)$ ($\mu, v \in \text{Hom}_A(Q, Q^*)$). Applying the decomposition of Lemma 4.3,

$$\begin{aligned} (Q_x, \varphi) &= (R_x, \mu_R + (x-1)v_R) \oplus (S_x, \mu_S + (x-1)v_S) \\ &= \{(R_x, \mu_R + (x-1)v_R) \oplus (S_x, \mu_S)\} \oplus \{(S_x, \mu_S + (x-1)v_S) \\ &\quad \oplus (S_x, -\mu_S) \oplus H_{\pm}(-S_x)\} \\ &= \bar{E}_+((R[x], \mu_R + (x-1)v_R) \oplus (S[x], \mu_S)) \\ &\quad \oplus \Delta_+(S[x^{-1}], x^{-1}(\mu_S + (x-1)v_S)) \in V_{2i}(A[x, x^{-1}]). \end{aligned}$$

As pointed out above, this suffices to prove the exactness of

$$V_{2i}(A[x]) \xrightarrow{E_+} V_{2i}(A[x, x^{-1}]) \xrightarrow{\delta_+} U_{2i}^{K_0(A)}(A[x^{-1}]).$$

Define next a morphism

$$\begin{aligned} E_+ : V_{2i}(A[x, x^{-1}]) &\rightarrow V_{2i}(A[x]); \\ (Q_x, \varphi) &\mapsto ((\varphi \pm \varphi^*)^{-1} (x^{N_1+1} Q^{*-}) \cap x^{-N_1} Q^{*+})[x], [\varphi]_0 - x([\varphi]_1) \\ &\quad \oplus ((x^N Q^- \cap (\varphi \pm \varphi^*)^{-1} (x^{-N} Q^{*+})) [x], [\varphi]_{-1} - [\varphi]_{-2}) \end{aligned}$$

for $N, N_1 \geq 0$ so large that

$$(\varphi \pm \varphi^*)(Q) \subseteq \sum_{j=-2N}^{2N_1+1} x^j Q^*$$

with $Q \in |\mathbf{P}(A)|$ f.g. free. The verification that E_+ is well-defined is by analogy with that for δ_+ . Moreover, if

$$(Q_x, \varphi) = (R_x, \mu_R + (x-1)v_R) \oplus (S_x, \mu_S + (x-1)v_S)$$

(as in Lemma 4.3), then

$$E_+(Q_x, \varphi) = (R[x], \mu_R + (x-1)v_R) \oplus (S[x], \mu_S) \in V_{2i}(A[x]),$$

so that the composites

$$\begin{aligned} U_{2i}^{K_0(A)}(A[x^{-1}]) &\xrightarrow{A_+} V_{2i}(A[x, x^{-1}]) \xrightarrow{E_+} V_{2i}(A[x]) \\ V_{2i}(A[x]) &\xrightarrow{E_+} V_{2i}(A[x, x^{-1}]) \xrightarrow{E_+} V_{2i}(A[x]) \end{aligned}$$

are 0, 1 respectively. Thus

$$V_{2i}(A[x]) \xrightarrow[E_+]{E_+} V_{2i}(A[x, x^{-1}]) \xrightarrow{\delta_+} U_{2i}^{K_0(A)}(A[x^{-1}])$$

defines a direct sum system, and we can identify

$$L_+ V_{2i}(A) = U_{2i}^{K_0(A)}(A[x^{-1}]).$$

Similarly, replacing x with x^{-1} , there is defined a direct sum system

$$V_{2i}(A[x^{-1}]) \xrightarrow[E_-]{E_-} V_{2i}(A[x, x^{-1}]) \xrightarrow{\delta_-} U_{2i}^{K_0(A)}(A[x]),$$

allowing the identification

$$L_- V_{2i}(A) = U_{2i}^{K_0(A)}(A[x]).$$

The proof of Lemma 4.2 shows that every element $(Q[x^{-1}], \varphi) \in V_{2i}(A[x^{-1}])$ has a representative with

$$\varphi = \varphi_0 + x^{-1}\varphi_{-1} \in \text{Hom}_{A[x^{-1}]}(Q[x^{-1}], Q^*[x^{-1}]) \quad (\varphi_0, \varphi_{-1} \in \text{Hom}_A(Q, Q^*)).$$

The composite

$$V_{2i}(A[x^{-1}]) \xrightarrow{E_-} V_{2i}(A[x, x^{-1}]) \xrightarrow{E_+} V_{2i}(A[x])$$

sends such a representative to

$$\begin{aligned} E_+ E_-(Q[x^{-1}], \varphi) &= (((\varphi \pm \varphi^*)^{-1}(xQ^*) \cap Q^+) [x], [\varphi]_0 - [\varphi]_1) \\ &\oplus ((xQ^* \cap (\varphi \pm \varphi^*)^{-1}(x^{-1}Q^*)) [x], [\varphi]_{-1} - [\varphi]_{-2}) \\ &= (Q[x], \varphi_0) \oplus ((\varphi \pm \varphi^*)^{-1}(Q^* \oplus x^{-1}Q^*) [x], [\varphi]_{-1} \\ &\quad - [\varphi]_{-2}) \in V_{2i}(A[x, x^{-1}]). \end{aligned}$$

The A -module isomorphism

$$\begin{aligned} Q \oplus Q &\rightarrow (\varphi \pm \varphi^*)^{-1}(Q^* \oplus x^{-1}Q^*); \\ (y, y') &\mapsto (\varphi \pm \varphi^*)^{-1}((\varphi_0 \pm \varphi_0^*)y, x^{-1}(((\varphi_0 \pm \varphi_0^*) + \varphi_{-1} \pm \varphi_{-1}^*)y + (\varphi_0 \pm \varphi_0^*)y')) \end{aligned}$$

defines an isomorphism of \pm forms over A

$$(Q \oplus Q, \begin{pmatrix} \varphi_0 + \varphi_{-1} & 0 \\ 0 & -\varphi_0 \end{pmatrix}) \rightarrow ((\varphi \pm \varphi^*)^{-1} (Q^* \oplus x^{-1}Q^*), [\varphi]_{-1} - [\varphi]_{-2}).$$

Therefore

$$\begin{aligned} E_+ \bar{E}_- (Q[x^{-1}], \varphi_0 + x^{-1}\varphi_{-1}) &= (Q[x], \varphi_0 + \varphi_{-1}) \oplus (Q[x] \oplus Q[x], \varphi_0 \oplus -\varphi_0) \\ &= (Q[x], \varphi_0 + \varphi_{-1}) \\ &= \bar{\varepsilon}_+ \varepsilon_- (Q[x^{-1}], \varphi_0 + x^{-1}\varphi_{-1}) \in V_{2i}(A[x]), \end{aligned}$$

and the square

$$\begin{array}{ccc} V_{2i}(A[x^{-1}]) & \xrightarrow{E_-} & V_{2i}(A[x, x^{-1}]) \\ \varepsilon_- \downarrow & & \downarrow E_+ \\ V_{2i}(A) & \xrightarrow{\bar{\varepsilon}_+} & V_{2i}(A[x]) \end{array}$$

commutes. Similarly, we can verify that the square

$$\begin{array}{ccc} U_{2i}^{K_0(A)}(A[x^{-1}]) & \xrightarrow{\eta_+} & U_{2i}(A) \\ \Delta_+ \downarrow & & \downarrow \bar{\eta}_- \\ V_{2i}(A[x, x^{-1}]) & \xrightarrow{\delta_-} & U_{2i}^{K_0(A)}(A[x]) \end{array}$$

commutes, where

$$\eta_{\pm} : U_{2i}^{K_0(A)}(A[x^{\mp 1}]) \rightarrow U_{2i}(A), \quad \bar{\eta}_{\pm} : U_{2i}(A) \rightarrow U_{2i}^{K_0(A)}(A[x^{\mp 1}])$$

are the morphisms induced by

$$\eta_{\pm} : A[x^{\mp 1}] \rightarrow A; \sum_{j=0}^{\infty} x^{\mp j} a_j \mapsto a_0, \quad \bar{\varepsilon}_{\mp} : A \rightarrow A[x^{\mp 1}]$$

respectively (so that $\eta_{\pm} \bar{\eta}_{\pm} = 1$). For

$$\begin{aligned} \delta_- \Delta_+ (Q[x^{-1}], \varphi = \varphi_0 + x^{-1}\varphi_{-1}) &= \delta_- ((Q_x, x\varphi) \oplus (Q_x, -(\varphi_0 + \varphi_{-1})) \oplus H_{\pm}(-Q_x)) \\ &= ((x^{-1}Q^+ \cap (\varphi \pm \varphi^*)^{-1}(Q^*))[x], [x\varphi]_{-1} - x[x\varphi]_0) \\ &= ((x^{-1}Q)[x], [x\varphi]_{-1}) = (Q[x], \varphi_0) \\ &= \bar{\eta}_- \eta_+ (Q[x^{-1}], \varphi) \in U_{2i}^{K_0(A)}(A[x]). \end{aligned}$$


The conditions of Lemma 1.1 are now satisfied, and so

$$V_{2i} : (\text{rings with involution}) \rightarrow (\text{abelian groups})$$

is a contracted functor, with

$$L_{\pm} V_{2i}(A) = U_{2i}^{K_0(A)}(A[x^{\mp 1}]), \quad LV_{2i}(A) = U_{2i}(A)$$

(up to natural isomorphisms), and the diagram




incorporates two commutative exact braids.

Let $S_0 \subseteq \tilde{K}_1(A[x, x^{-1}])$ be the infinite cyclic subgroup generated by $\bar{B}([A]) = \tau(x: A_x \rightarrow A_x)$, and define

$$\tilde{W}_n(A[x, x^{-1}]) = V_n^{S_0}(A[x, x^{-1}]) \quad (n \pmod 4).$$

Working as for $V_{2i}(A[x, x^{-1}])$, it is possible to define morphisms to fit into a diagram



(with $E_+ \bar{E}_+ = 1$ etc.) incorporating two commutative exact braids. For example,

$$\delta_+ : \tilde{W}_{2i}(A[x, x^{-1}]) \rightarrow V_{2i}^{K_1(A)}(A[x^{-1}]); (\tilde{Q}_x, \varphi) \mapsto (\tilde{P}[x^{-1}], [\varphi]_{-1} - x^{-1}[\varphi]_{-2})$$

$$E_+ : \tilde{W}_{2i}(A[x, x^{-1}]) \rightarrow W_{2i}(A[x]);$$

$$(\tilde{Q}_x, \varphi) \mapsto (\tilde{P}_1[x], [\varphi]_0 - x[\varphi]_1) \oplus (\tilde{P}[x], [\varphi]_{-1} - [\varphi]_{-2})$$

for any A -base P of $P = x^N Q^- \cap (\varphi \pm \varphi^*)^{-1} (x^{-N} Q^{*+})$ (which is free for sufficiently large $N \geq 0$, as $\tau(\tilde{Q}_x, \varphi) \in S_0$ and $[P] = B\tau(\tilde{Q}_x, \varphi) = 0 \in \tilde{K}_0(A)$) with

$$P_1 = (\varphi \pm \varphi^*)^{-1} (x^N Q^*) \oplus (\varphi \pm \varphi^*)^{-1} (\tilde{Q}^*)$$

the corresponding A -base of $P_1 = (\varphi \pm \varphi^*)^{-1} (x^{N+1} Q^{*-}) \cap x^{-N} Q^+$, for N so large that

$$(\varphi \pm \varphi^*) (Q) \subseteq \sum_{j=-2N}^{2N+1} x^j Q^*.$$

Also, let

$$A_+: V_{2i}^{K_1(A)}(A[x^{-1}]) \rightarrow \tilde{W}_{2i}(A[x, x^{-1}]); (\tilde{Q}[x^{-1}], \varphi) \mapsto (\tilde{Q}_x, x\varphi) \oplus (\tilde{Q}_x, -\bar{e}\varepsilon_- \varphi)$$

where $\tilde{Q} = (\varepsilon_- (\varphi \pm \varphi^*))^{-1} (\tilde{Q}^*)$.

Given an invertible indeterminate z over A commuting with every element of A define A_z as $A[z, z^{-1}]$ but with involution by $\bar{z} = z^{-1}$. Similarly, define $A[x^{\pm 1}]_z$, $A[x, x^{-1}]_z$, and identify

$$A[x^{\pm 1}]_z = A_z[x^{\pm 1}], \quad A[x, x^{-1}]_z = A_z[x, x^{-1}].$$


Let $S'_0 \subseteq \tilde{K}_1(A_z)$ be the infinite cyclic subgroup generated by $\tau(z: A_z \rightarrow A_z)$ and define

$$\tilde{W}_n(A_z) = V_n^{S'0}(A_z)$$

$$\tilde{W}_n(A[x^{\pm 1}]_z) = V_n^{\tilde{e}_\pm(x)S'0}(A[x^{\pm 1}]_z)$$

$$\tilde{\tilde{W}}_n(A[x, x^{-1}]_z) = V_n^{\tilde{\epsilon}(z)S_0 \oplus \tilde{\epsilon}(x)S'0}(A[x, x^{-1}]_z)$$


for $n \pmod{4}$. By analogy with $\tilde{W}_{2i}(A[x, x^{-1}])$, $\tilde{\tilde{W}}_{2i}(A[x, x^{-1}]_z)$ fits into a diagram incorporating two commutative exact braids (where $A_z = A[z, z^{-1}]$, with $\bar{z} = z^{-1}$).



We can now apply the decompositions

$$\begin{aligned}
 \tilde{W}_{2i}(A_z) &= \bar{\varepsilon}(z) W_{2i}(A) \oplus \bar{B}(z) V_{2i-1}(A) \\
 \tilde{W}_{2i}(A[x]_z) &= \bar{\varepsilon}(z) W_{2i}(A[x]) \oplus \bar{B}(z) V_{2i-1}(A[x]) \\
 \tilde{W}_{2i}(A[x, x^{-1}]_z) &= \bar{\varepsilon}(z) \tilde{W}_{2i}(A[x, x^{-1}]) \oplus \bar{B}(z) V_{2i-1}(A[x, x^{-1}]) \\
 V_{2i}^{K_1(A_z)}(A[x]_z) &= \bar{\varepsilon}(z) V_{2i}^{K_1(A)}(A) \oplus \bar{B}(z) U_{2i}^{K_0(A)}(A) \\
 V_{2i}(A_z) &= \bar{\varepsilon}(z) V_{2i}(A) \oplus \bar{B}(z) U_{2i-1}(A)
 \end{aligned}$$

given by Theorem 1.1 of Part II (and extended to the intermediate L -groups in Part III). The above diagram splits naturally (via $\bar{\varepsilon}(z), \bar{B}(z)$) into two similar ones: the diagram for $\tilde{W}_{2i}(A[x, x^{-1}])$ and the diagram



where

$$\begin{aligned}
 E_+ : V_{2i-1}(A[x, x^{-1}]) &\xrightarrow{\bar{B}(z)} \tilde{W}_{2i}(A[x, x^{-1}]_z) \xrightarrow{E_+} \tilde{W}_{2i}(A[x]_z) \xrightarrow{B(z)} V_{2i-1}(A[x]) \\
 \delta_+ : V_{2i-1}(A[x, x^{-1}]) &\xrightarrow{\bar{B}(z)} \tilde{W}_{2i}(A[x, x^{-1}]_z) \\
 &\xrightarrow{\delta_+} V_{2i}^{K_1(A_z)}(A[x^{-1}]_z) \xrightarrow{B(z)} U_{2i-1}^{K_0(A)}(A[x^{-1}]) \\
 \Delta_+ : U_{2i-1}^{K_0(A)}(A[x^{-1}]) &\xrightarrow{\bar{B}(z)} V_{2i}^{K_1(A_z)}(A[x^{-1}]_z) \\
 &\xrightarrow{\Delta_+} \tilde{W}_{2i}(A[x, x^{-1}]_z) \xrightarrow{B(z)} V_{2i-1}(A[x, x^{-1}])
 \end{aligned}$$

(and similarly for E_-, δ_-, Δ_-). Thus the conditions of Lemma 1.1 are also satisfied in the odd-dimensional case, and

$$V_{2i-1} : (\text{rings with involution}) \rightarrow (\text{abelian groups})$$

is a contracted functor, with identifications

$$L_{\pm} V_{2i-1}(A) = U_{2i-1}^{K_0(A)}(A[x^{\pm 1}]), \quad LV_{2i-1}(A) = U_{2i-1}(A).$$

This completes the proof of Theorem 4.1 \square

The groups

$$\text{Nil}_{\pm}(A) = \ker(\varepsilon_{\pm}: K_1(A[x^{\pm 1}]) \rightarrow K_1(A))$$

are such that

$$\begin{aligned} K_1(A[x^{\pm 1}]) &= \bar{\varepsilon}_{\pm} K_1(A) \oplus \text{Nil}_{\pm}(A) \\ K_1(A[x, x^{-1}]) &= \bar{\varepsilon} K_1(A) \oplus \bar{E}_+ \text{Nil}_+(A) \oplus \bar{E}_- \text{Nil}_-(A) \oplus \bar{B} K_0(A), \end{aligned}$$

fitting into direct sum systems

$$\text{Nil}_{\pm}(A) \xrightleftharpoons[E_{\pm} A_{\pm}]{\delta_{\pm} E_{\pm}} K_0 \text{N}(A) \xrightleftharpoons[\bar{\eta}_{\pm}]{\eta_{\pm}} K_0(A)$$

(by Theorem 2.1).

Given $*$ -invariant subgroups $S_{\pm} \subseteq \text{Nil}_{\pm}(A)$, define

$$\begin{aligned} N_{\pm} V_n^{S_{\pm}}(A) &= \ker(\varepsilon_{\pm}: V_n^{\bar{\varepsilon}_{\pm} K_1(A) \oplus S_{\pm}}(A[x^{\pm 1}]) \rightarrow V_n(A)) \quad (n \pmod{4}) \\ \text{writing} \quad \begin{cases} N_{\pm} V_n(A) \\ N_{\pm} W_n(A) \end{cases} \quad \text{for} \quad \begin{cases} N_{\pm} V_n^{\text{Nil}_{\pm}(A)}(A) \\ N_{\pm} V_n^{\{0\}}(A) \end{cases}. \end{aligned}$$

COROLLARY 4.4. *Given $*$ -invariant subgroups*

$$R \subseteq \tilde{K}_1(A), \quad S_{\pm} \subseteq \text{Nil}_{\pm}(A), \quad \tilde{T} \subseteq \tilde{K}_0(A)$$

there are direct sum decompositions

$$\begin{aligned} V_n^{\bar{\varepsilon}_{\pm} R \oplus S_{\pm}}(A[x^{\pm 1}]) &= \bar{\varepsilon}_{\pm} V_n^R(A) \oplus N_{\pm} V_n^{S_{\pm}}(A) \\ U_n^{\bar{\varepsilon}_{\pm} T}(A[x^{\pm 1}]) &= \bar{\varepsilon}_{\pm} U_n^T(A) \oplus N_{\pm} V_n(A) \\ V_n^Q(A[x, x^{-1}]) &= \bar{\varepsilon} V_n^R(A) \oplus \bar{E}_+ N_+ V_n^{S_+}(A) \oplus \bar{E}_- N_- V_n^{S_-}(A) \oplus \bar{B} U_n^T(A) \end{aligned}$$

for $n \pmod{4}$, where

$$\begin{aligned} Q &= \bar{\varepsilon} R \oplus \bar{E}_+ S_+ \oplus \bar{E}_- S_- \oplus \bar{B} T \subseteq \tilde{K}_1(A[x, x^{-1}]) \\ &= \bar{\varepsilon} \tilde{K}_1(A) \oplus \bar{E}_+ \text{Nil}_+(A) \oplus \bar{E}_- \text{Nil}_-(A) \oplus \bar{B} K_0(A) \end{aligned}$$

with $T \subseteq K_0(A)$ the preimage of \tilde{T} under the natural projection $K_0(A) \rightarrow \tilde{K}_0(A)$.

Proof. The forgetful map

$$V_n(A[x^{\pm 1}]) \rightarrow U_n^{\bar{\varepsilon}_{\pm} T}(A[x^{\pm 1}])$$

fits into the exact sequence of Theorem 2.3 of Part III, which splits, via $\bar{\varepsilon}_\pm, \varepsilon_\pm$ into two exact sequences

$$\begin{array}{ccccccc}
 \rightarrow & 0 & \rightarrow & N_\pm V_n(A) & \rightarrow & N_\pm V_n(A) & \rightarrow 0 \rightarrow \\
 & \uparrow & & \uparrow & & \uparrow & \uparrow \\
 \rightarrow & H^{n+1}(\bar{\varepsilon}_\pm \tilde{T}) & \rightarrow & V_n(A[x^{\pm 1}]) & \rightarrow & U_n^{\bar{\varepsilon}_\pm T}(A[x^{\pm 1}]) & \rightarrow H^n(\bar{\varepsilon}_\pm \tilde{T}) \rightarrow \\
 & \bar{\varepsilon}_\pm \uparrow \varepsilon_\pm & & \bar{\varepsilon}_\pm \uparrow \varepsilon_\pm & & \bar{\varepsilon}_\pm \uparrow \varepsilon_\pm & \bar{\varepsilon}_\pm \uparrow \varepsilon_\pm \\
 \rightarrow & H^{n+1}(\tilde{T}) & \rightarrow & V_n(A) & \rightarrow & U_n^T(A) & \rightarrow H^n(\tilde{T}) \rightarrow .
 \end{array}$$

Hence $N_\pm V_n(A) \subseteq V_n(A[x^{\pm 1}])$ is mapped isomorphically to $\ker(\varepsilon_\pm : U_n^{\bar{\varepsilon}_\pm T}(A[x^{\pm 1}]) \rightarrow U_n^T(A))$ and so (up to isomorphism)

$$U_n^{\bar{\varepsilon}_\pm T}(A[x^{\pm 1}]) = \bar{\varepsilon}_\pm U_n^T(A) \oplus N_\pm V_n(A).$$

In particular,

$$\begin{aligned}
 U_n^{R_0(A)}(A[x^{\pm 1}]) &= \bar{\varepsilon}_\pm U_n(A) \oplus N_\pm V_n(A), \\
 V_n(A[x^{\pm 1}]) &= \bar{\varepsilon}_\pm V_n(A) \oplus N_\pm V_n(A).
 \end{aligned}$$

It now follows from Theorem 4.1 that

$$V_n(A[x, x^{-1}]) = \bar{\varepsilon} V_n(A) \oplus \bar{E}_+ N_+ V_n(A) \oplus \bar{E}_- N_- V_n(A) \oplus \bar{B} U_n(A).$$

The expressions for $V_n^{\bar{\varepsilon}_\pm R \oplus S_\pm}(A[x^{\pm 1}])$, $V_n^Q(A[x, x^{-1}])$ may be deduced from those for $V_n(A[x^{\pm 1}])$, $V_n(A[x, x^{-1}])$, working as for $U_n^{\bar{\varepsilon}_\pm T}(A[x^{\pm 1}])$ above. (In particular, for $R=0$, $S_+=0$, $S_-=0$, $\tilde{T}=0$ we have

$$Q = S_0 \subseteq \tilde{K}_1(A[x, x^{-1}])$$

and

$$\begin{aligned}
 W_n(A[x^{\pm 1}]) &= \bar{\varepsilon}_\pm W_n(A) \oplus N_\pm W_n(A), \\
 \tilde{W}_n(A[x, x^{-1}]) &= \bar{\varepsilon} W_n(A) \oplus \bar{E}_+ N_+ W_n(A) \oplus \bar{E}_- N_- W_n(A) \oplus \bar{B} V_n(A). \quad \square
 \end{aligned}$$

In §4 of Part II there were defined lower L -theories, functors

$$L_n^{(m)} : (\text{rings with involution}) \rightarrow (\text{abelian groups})$$

for $m < 0$, $n \pmod{4}$ by

$$L_n^{(m)}(A) = \ker(\varepsilon : L_{n+1}^{(m+1)}(A_z) \rightarrow L_{n+1}^{(m+1)}(A))$$

with $L_n^{(0)}(A) = U_n(A)$. By convention, $L_n^{(1)}(A) = V_n(A)$.

COROLLARY 4.5. *The lower L -theories $L_n^{(m)}$ coincide (up to natural isomorphism)*

with the functors LV_n , L^2V_n , ... derived from V_n , with

$$L_n^{(m)}(A) = L^{1-m}V_n(A) \quad (m \leq 0, n \pmod{4}).$$

Proof. By Theorem 4.1,

$$LV_n(A) = U_n(A) = L_n^{(0)}(A).$$

Assume inductively that

$$L_n^{(p)}(A) = L^{1-p}V_n(A) \quad (n \pmod{4})$$

for $0 \geq p > m$, for some $m \leq -1$. Then

$$\begin{aligned} L_n^{(m)}(A) &= \ker(\varepsilon: L_{n+1}^{(m+1)}(A_z) \rightarrow L_{n+1}^{(m+1)}(A)) \\ &= \ker(\varepsilon: L^{-m}V_{n+1}(A_z) \rightarrow L^{-m}V_{n+1}(A)) \\ &= L(\ker(\varepsilon: L^{-m-1}V_{n+1}(A_z) \rightarrow L^{-m-1}V_{n+1}(A))) \\ &= L(\ker(\varepsilon: L_{n+1}^{(m+2)}(A_z) \rightarrow L_{n+1}^{(m+2)}(A))) \\ &= LL_n^{(m+1)}(A) \\ &= LL^{-m}V_n(A) = L^{1-m}V_n(A) \end{aligned}$$

giving the induction step. \square

Given a functor

$$F: (\text{rings with involution}) \rightarrow (\text{abelian groups})$$

define

$$N_{\pm}F(A) = \ker(\varepsilon_{\pm}: F(A[x^{\pm 1}]) \rightarrow F(A)).$$

(By Corollary 4.4, the previous definitions of $N_{\pm}V_n(A)$, $N_{\pm}W_n(A)$ agree with this, up to natural isomorphism).

By analogy with the first part of Corollary 7.6 of Chapter XII of [1] we have

COROLLARY 4.6. *Let x_1, x_2, \dots, x_p be independent commuting indeterminates over A , with $\bar{x}_j = x_j$ ($1 \leq j \leq p$). Then*

$$\begin{aligned} L_n^{(m)}(A[x_1, x_2, \dots, x_p]) &= (1 \oplus N_+)^p L_n^{(m)}(A) \\ L_n^{(m)}(A[x_1, x_1^{-1}, x_2, x_2^{-1}, \dots, x_p, x_p^{-1}]) &= (1 \oplus N_+ \oplus N_- \oplus L)^p L_n^{(m)}(A) \end{aligned}$$

up to natural isomorphism, for $m \leq 1$, $n \pmod{4}$, $p \geq 1$. \square

REFERENCES

- [1] BASS H., *Algebraic K-theory*, Benjamin (1968).
- [2] BASS H., HELLER A. and SWAN R. G., *The Whitehead group of a polynomial extension*, Publ. Math. IHES no. 22 (1964).
- [3] HIGMAN, G., *The units of group-rings*, Proc. London Math. Soc. (2) 46 (1940), 231–48.
- [4] KAROUBI, M., *Localisation de formes quadratiques*, (preprint).
- [5] RANICKI, A. A., *Algebraic L-theory, I: Foundations*, Proc. London Math. Soc. (3) 27 (1973), 101–25.
- [6] ——, *Algebraic L-theory, II: Laurent extensions*, Proc. London Math. Soc. (3) 27 (1973), 126–58.
- [7] ——, *Algebraic L-theory, III: Twisted Laurent extensions*, in *Algebraic K-theory III*, Springer Lecture Notes No. 343 (1973), 412–463.
- [8] SWAN, R. G., *Algebraic K-theory*, Springer Lecture Notes No. 76 (1968).

Trinity College, Cambridge, England

Received August 17, 1973