

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 49 (1974)

Artikel: Functions of Bounded mean Oscillation and Quasiconformal Mappings
Autor: Reimann, H.M.
DOI: <https://doi.org/10.5169/seals-37994>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Functions of Bounded mean Oscillation and Quasiconformal Mappings

H. M. REIMANN

1. Introduction

A locally integrable real valued function u is said to be of bounded mean oscillation (BMO) in \mathbf{R}^n , if

$$\frac{1}{|Q|} \int_Q \left| u(x) - \frac{1}{|Q|} \int_Q u(x) dx \right| dx \leq K$$

for every cube $Q \subset \mathbf{R}^n$ and some constant K . The notations

$$u_Q = \frac{1}{|Q|} \int_Q u(x) dx = \frac{1}{|Q|} \int_Q u(x) dx \quad \text{with} \quad |Q| = \int_Q dx$$

will be used. On the space of BMO-functions modulo constants a norm can be defined by

$$\|u\|_* = \sup_{Q \subset \mathbf{R}^n} \int_Q |u(x) - u_Q| dx. \quad (1.1)$$

With this norm BMO/\mathbf{R} is a Banach space. The space of BMO-functions was introduced by John and Nirenberg [8]. We will make use of their fundamental lemma:

LEMMA 1. *Assume that $u \in \text{BMO}$. Then, if $\mu(\sigma) = |\{x \in Q : |u(x) - u_Q| > \sigma\}|$ is the measure of the set of points in the cube Q where $|u(x) - u_Q| > \sigma$, we have*

$$\mu(\sigma) \leq a e^{-b\sigma/\|u\|_*} |Q|, \quad (1.2)$$

where a and b are constants depending on n only.

BMO-functions have been used in many different contexts, first in a paper of John on rotation and strain [7] and at the same time by Moser [9] in his work on the continuity of solutions of elliptic differential equations. Later on applications arose in connection with singular integral operators (Stein [12]) and as spaces of interpolation (Stampacchia [11], Stein and Zygmund [13]). Most recently, Fefferman and Stein [3] characterized the space of BMO-functions as the dual of the Hardy space H^1 .

It seems that BMO-functions also have their place in the theory of quasiconformal mappings. We propose to show that the logarithm of the Jacobian determinant of a

quasiconformal mapping $f: \mathbf{R}^n \rightarrow \mathbf{R}^n$ is in BMO. We then proceed to study transformations of BMO-functions by quasiconformal mappings. It turns out that a quasiconformal mapping of \mathbf{R}^n onto itself induces a continuous bijective isomorphism $\varphi: u \rightarrow u \circ f^{-1}$ of BMO/\mathbf{R} . Moreover this situation is in a certain way typical for quasiconformal mappings: If $\varphi: u \rightarrow u \circ f^{-1}$ is a continuous bijective isomorphism of BMO/\mathbf{R} which is induced by a homeomorphism f of \mathbf{R}^n satisfying certain regularity conditions, then f is quasiconformal.

2. The Jacobian of a Quasiconformal Mapping

For our considerations we adopt the so called analytic definition of quasiconformality. A function $f: G \rightarrow \mathbf{R}^n$ defined in a domain $G \subset \mathbf{R}^n$ is said to be absolutely continuous on lines (ACL), if it is continuous and if for each interval $I = \{x \in \mathbf{R}^n: a_i \leq x_i \leq b_i\} \subset G$ f is absolutely continuous on almost all line segments in I , parallel to the coordinate axes. The partial derivatives of an ACL-function f exist a.e. and the Jacobian matrix of f at x will be denoted by $F(x)$, its determinant by $J_f(x)$. By definition a K -quasiconformal mapping is a homeomorphism $f: G \rightarrow \mathbf{R}^n$ such that $f \in \text{ACL}$, f is totally differentiable a.e. and

$$\sup_{\xi \in \mathbf{R}^n, |\xi|=1} |F(x) \xi|^n \leq K J_f(x) \quad \text{a.e.} \quad (2.1)$$

According to a theorem of Väisälä [14], in this definition the regularity conditions $f \in \text{ACL}$ and f differentiable a.e. can be replaced by the single hypothesis, that f has generalized derivatives which are locally L^n -integrable.

THEOREM 1. *If f is a quasiconformal mapping of \mathbf{R}^n onto itself with Jacobian determinant J_f then $\log J_f \in \text{BMO}$.*

The proof of Theorem 1 is based on a converse to the lemma of John and Nirenberg (Lemma 3) and on the following result due to Gehring [5]:

LEMMA 2. *Assume that f is a K -quasiconformal mapping of G onto $G' \subset \mathbf{R}^n$ and that Q is a cube in the domain G with*

$$\text{dia } Q' < \text{dist}(Q', \partial G') \quad (2.2)$$

($Q' = fQ$). Then there exist constants c and p , $p > n$, which depend on K and n only, such that

$$\left(\int_Q J_f^{p/n} dx \right)^{n/p} \leq c \int_Q J_f dx. \quad (2.3)$$

We set

$$L_f(x) = \limsup_{y \rightarrow x} \frac{|f(y) - f(x)|}{|y - x|}.$$

Since f is K -quasiconformal, inequality (2.1) and the total differentiability imply $L_f^n \leq KJ_f$ a.e. and $J_f \leq L_f^n$ a.e. Hence according to Lemma 4 in [3] there exists a constant c_0 (depending on K and n) such that for every cube $Q \subset G$ with $\text{dia } Q' < \text{dist}(Q', \partial G')$

$$\int_Q J_f dx \leq c_0 \left(\int_Q J_f^{1/n} dx \right)^n, \quad (2.4)$$

If Q is such a cube (satisfying (2.3)), then (2.4) holds for any cube contained in Q and Lemma 3 in [5] shows that for some constants c and p , $p > n$,

$$\left(\int_Q J_f^{p/n} dx \right)^{n/p} \leq c \int_Q J_f dx.$$

For later reference let us note a simple consequence of this lemma (cf. [5] Theorem 2)

COROLLARY. Assume that f is a K -quasiconformal mapping of G onto $G' \subset \mathbb{R}^n$ and that Q is a cube in the domain G with $\text{dia } Q' < \text{dist}(Q', \partial G')$. Then

$$\frac{|A'|}{|Q'|} \leq c \left(\frac{|A|}{|Q|} \right)^{(p-n)/p} \quad (2.5)$$

for every measurable set $A \subset Q$ with image $A' = fA$. (As above $|A|$ stands for the n -dimensional measure of the set A .)

If A is a measurable set, $A \subset Q$, then

$$\begin{aligned} \frac{|A'|}{|A|} &= \int_A J_f dx \leq \left(\int_A J_f^{p/n} dx \right)^{n/p} \\ &\leq \left(\frac{|A|}{|Q|} \right)^{-n/p} \left(\int_Q J_f^{p/n} dx \right)^{n/p} \leq c \left(\frac{|A|}{|Q|} \right)^{-n/p} \frac{|Q'|}{|Q|} \end{aligned}$$

by Lemma 2 and Hölder's inequality.

Let us remark that for the case of plane quasiconformal mappings results similar to Lemma 2 have previously been established by Bojarski [1] and by Gehring and Reich [6].

LEMMA 3. $f = \log u \in \text{BMO}$ if and only if for all cubes $Q \subset \mathbb{R}^n$

$$\left(\int_Q u^a dx \right)^{1/a} \leq k \left(\int_Q u^{-b} dx \right)^{-1/b} \quad (2.6)$$

for some positive constants a, b and k .

The fact, that inequality (2.6) is a consequence of Lemma 1 has already been pointed out by John and Nirenberg. (The result has been stated in this form by Moser [9].) Let us therefore assume that (2.6) holds for $u = e^f$. It is well known that

$$M_t = M_t(u) = \begin{cases} \left(\int_Q u^t dx \right)^{1/t} & t \neq 0 \\ \exp \int_Q \log u dx & t = 0 \end{cases}$$

is a monotone increasing function of t . Our assumption therefore implies $M_s \leq KM_0$ and $M_0 \leq KM_{-s}$ for $s = \min(a, b)$. If we set

$$Q_1 = \left\{ x \in Q : \log u(x) \geq \int_Q \log u dx \right\}$$

and $Q_2 = Q \setminus Q_1$ we obtain the inequalities

$$|Q|^{-1} \int_{Q_1} u^s dx \leq \int_Q u^s dx = M_s^s \leq k^s M_0^s$$

and

$$|Q|^{-1} \int_{Q_2} u^{-s} dx \leq k^{-s} M_0^{-s}.$$

After adding these two inequalities and inserting the expressions

$$f = \log u \quad \text{and} \quad f_Q = \int_Q f dx = \log M_0$$

we have

$$\int_{Q_1} e^{s(f-f_Q)} dx + \int_{Q_2} e^{-s(f-f_Q)} dx = \int_Q e^{s|f-f_Q|} dx \leq (k^s + k^{-s}) |Q|.$$

Finally,

$$\exp \int_Q s |f - f_Q| dx \leq \int_Q e^{s|f-f_Q|} dx$$

upon applying Jensen's inequality. This shows that $f \in \text{BMO}$ with

$$\|f\|_* \leq s^{-1} \log(k^s + k^{-s}). \quad (2.7)$$

We shall also need a distortion lemma for quasiconformal mappings, to the effect that the image of a cube can still be compared with a cube. This kind of result is typical for the geometric theory of quasiconformal mappings. We choose a formulation, which is particularly suited for our purposes.

LEMMA 4. *Let $f: G \rightarrow \mathbb{R}^n$ be a K -quasiconformal mapping. There exists a constant k (which depends on K and n) such that to every cube $P' \subset G' = fG$ with*

$$\text{dist}(P', \partial G') > 2k \text{ dia } P'$$

there exists a cube $Q \subset G$ with $fQ = Q' \supset P'$, $\text{dia } Q' < \text{dist}(Q', \partial G')$ and

$$|Q'| \leq k^n n^{n/2} |P'|. \quad (2.8)$$

The proof is based on the geometric definition of quasiconformality, according to which a homeomorphism $f: G \rightarrow \mathbb{R}^n$ is K -quasiconformal if and only if

$$\text{mod } R' \leq K^{1/(n-1)} \text{mod } R$$

for every ring $R \subset G$. We refer the reader to the literature (see e.g. [10], [2]) for the precise definitions and for a proof of the equivalence of the analytic and geometric definitions of quasiconformality.

Using preliminary translations, we can assume that the given cube P' is centered at 0 and that $f(0) = 0$. Consider the spherical ring $R' \subset G'$ with complementary components

$$C'_0 = \{z: |z| \leq r' = 2^{-1} \text{dia } P'\} \quad \text{and} \quad C'_1 = \{z: |z| \geq s' = k2^{-1} \text{dia } P'\}.$$

The constant $k > 1$ will be determined later on. The modulus of the ring R' is given by

$$\text{mod } R' = \log \frac{s'}{r'} = \log k. \quad (2.9)$$

We set $s = \inf_{z \in C'_1} |f^{-1}(z)|$ and $r = \sup_{z \in C'_0} |f^{-1}(z)|$ and observe that $|f^{-1}(z)| \leq r$ for all $z \in P'$.

The inner complementary component C_0 of $R = f^{-1}R'$ contains 0 and a point x_0 with $|x_0| = r$, the outer component C_1 a continuum connecting ∞ with a point x_1 , $|x_1| = s$. According to a theorem of Teichmüller and its space analogue (see [4], [2], [10])

$$\text{mod } R \leq \log \left(\lambda^2 \left(\frac{s}{r} + 1 \right) \right) \quad (2.10)$$

for some constant λ which depends on n only. Since K -quasiconformal mappings satisfy

$$\text{mod } R' \leq K_0 \text{ mod } R$$

with $K_0 = K^{1/(n-1)}$, we then have by (2.9) and (2.10)

$$\log k \leq K_0 \log \left(\lambda^2 \left(\frac{s}{r} + 1 \right) \right)$$

which is equivalent to

$$k^{-K_0} \leq \lambda^2 \left(\frac{s}{r} + 1 \right).$$

This shows that $s/r \geq n^{1/2}$ if we choose $k = (\lambda^2(1 + n^{1/2}))^{K_0}$. In this situation any cube $Q \subset G$ centered at the origin with side length $2r$ (and diameter $2r n^{1/2}$) satisfies the requirements of the lemma: The construction shows that $\text{dia } Q' < k \text{ dia } P'$ and

$$|Q'| \leq k^n n^{n/2} |P'|$$

since $|z| \leq k 2^{-1} \text{ dia } P'$ for all $z \in Q'$. If we further assume that $\text{dist}(P', \partial G') > 2k \text{ dia } P' = 4s'$, then it is clear that $\text{dia } Q' \leq 2s' \leq \text{dist}(Q', \partial G')$.

LEMMA 5. *The Jacobian determinant $J = J_f$ of a K -quasiconformal mapping $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies*

$$\int_Q J \, dx \leq c_1 \left(\int_Q J^{-b} \, dx \right)^{-1/b} \quad (2.11)$$

for all cubes $Q \subset \mathbb{R}^n$. The constants b and c_1 depend on n and K only.

The inverse f^{-1} of a K -quasiconformal mapping is K^{n-1} -quasiconformal and its determinant J^{-1} satisfies Gehring's inequality (see (2.3)):

$$\left(\int_{P'} J^{-p/n} \, dy \right)^{n/p} \leq c \int_{P'} J^{-1} \, dy \quad (2.12)$$

for every cube $P' \subset \mathbb{R}^n$. To any cube $Q \subset \mathbb{R}^n$ let us choose in accordance with Lemma 4 a cube P' with $P = f^{-1} P' \supset Q$ and

$$|P| \leq k^n n^{n/2} |Q| = k' |Q|. \quad (2.13)$$

A transformation of variables for the integrals in inequality (2.12) then shows that

$$\left(|P'|^{-1} \int_P J^{-p/n} J \, dx \right)^{n/p} \leq c |P| |P'|^{-1}.$$

Because $|P'| = \int_P J \, dx$ this can be rewritten in the form

$$\left(\int_P J^{(n-p)/n} \, dx \right)^{n/p} \leq c |P| \left(\int_P J \, dx \right)^{(n-p)/p}$$

and together with inequality (2.13) this leads to

$$\left(\int_Q J^{(n-p)/n} \, dx \right)^{n/(n-p)} \geq (ck')^{p/(n-p)} \int_Q J \, dx.$$

If the two Lemmata 3 and 5 are combined, a bound for $\|\log J\|_*$ can be given by

$$\|\log J\|_* \leq \frac{n}{p-n} \log((ck')^{p/n} + (ck')^{-p/n})$$

provided that $p \leq 2n$ and by

$$\|\log J\|_* \leq \log((ck')^{p/(p-n)} + (ck')^{p/(n-p)})$$

otherwise.

Remark 1. By definition, a K -quasiconformal mapping $f = (f_1, \dots, f_n)$ of \mathbf{R}^n onto itself satisfies for $i = 1, \dots, n$

$$K^{-n+1} J_f \leq |\operatorname{grad} f_i|^n \leq K J_f \quad \text{a.e. in } \mathbf{R}^n.$$

Hence there exist $g_i \in L^\infty(\mathbf{R}^n)$, $\|g_i\|_\infty \leq (n-1) \log K$, such that $n \log |\operatorname{grad} f_i| = \log J_f + g_i$ ($i = 1, \dots, n$). But functions in $L^\infty(\mathbf{R}^n)$ are also in BMO and therefore $\log |\operatorname{grad} f_i| \in \text{BMO}$ ($i = 1, \dots, n$).

Remark 2. Local variants of Theorem 1 can be obtained. If $f: G \rightarrow G'$ is a quasiconformal mapping and if $Q \subset G$ is a cube such that both $\operatorname{dist}(Q, \partial G)$ and $\operatorname{dist}(Q', \partial G')$ are big enough, then $\log J_f$ considered as a function in Q is in BMO.

3. The Invariance of the Space BMO

THEOREM 2. *If f is a K -quasiconformal mapping of \mathbf{R}^n onto itself, then $\varphi: u \rightarrow u' = u \circ f^{-1}$ is a bijective isomorphism of BMO and*

$$\|u'\|_* \leq C \|u\|_* \tag{3.1}$$

for all $u \in \text{BMO}$, where C is a constant depending on K and n only.

We note that since the inverse of a quasiconformal mapping is also a quasiconformal mapping, all that has to be shown is inequality (3.1). It then follows directly from the definition that φ is an isomorphism of BMO.

For the proof of Theorem 2 we assume that $u \in \text{BMO}$ and set $u' = u \circ f^{-1}$. To a given cube P' we determine Q as in Lemma 4 such that $P' \subset Q'$ and $|Q'| \leq k^n n^{n/2} |P'|$. The

set $A_\sigma = \{x \in Q : |u(x) - u_Q| > \sigma\}$ is mapped onto the set $A'_\sigma = \{z \in Q' : |u'(z) - u_Q| > \sigma\}$ and by the corollary to Lemma 2 one knows that

$$\frac{|A'_\sigma|}{|Q'|} \leq c \left(\frac{|A_\sigma|}{|Q|} \right)^{(p-n)/p}.$$

Because of Lemma 1

$$|A_\sigma| \leq a e^{-b\sigma/\|u\|_*} |Q|$$

hence

$$\frac{|A'_\sigma|}{|Q'|} \leq ca^{(p-n)/p} \exp\left(\frac{-b\sigma(p-n)}{\|u\|_* p}\right).$$

An integration of this inequality with respect to σ shows that

$$\int_{Q'} |u'(z) - u_Q| dz = |Q'|^{-1} \int_0^\infty |A'_\sigma| d\sigma \leq ca^{(p-n)/p} b^{-1} p(p-n)^{-1} \|u\|_*$$

and in combination with inequality (2.8) of Lemma 4

$$\int_{P'} |u'(z) - u_Q| dz \leq k^n n^{n/2} \int_{Q'} |u'(z) - u_Q| dz \leq \text{const } \|u\|_*$$

One is left with the task of replacing u_Q by

$$u'_{P'} = \int_{P'} u'(z) dz,$$

but

$$|u'_{P'} - u_Q| = \left| \int_{P'} (u'(z) - u_Q) dz \right|,$$

so

$$\int_{P'} |u'(z) - u'_{P'}| dz \leq |u'_{P'} - u_Q| + \int_{P'} |u'(z) - u_Q| dz \leq 2 \int_{P'} |u'(z) - u_Q| dz.$$

This shows that $\|u'\|_* \leq C\|u\|_*$ with $C = 2k^n n^{n/2} c a^{(p-n)/p} b^{-1} p(p-n)^{-1}$.

THEOREM 3. Assume that f is a (orientation preserving) homeomorphism of \mathbf{R}^n onto itself, that $f \in \text{ACL}$ and that f is totally differentiable a.e. If the induced mapping $\varphi: u \rightarrow u' = u \circ f^{-1}$ is a bijective isomorphism of BMO and if

$$\|u'\|_* \leq C\|u\|_* \quad \text{for all } u \in \text{BMO} \quad (3.1)$$

then f is a quasiconformal mapping.

Let us make precise that the hypothesis of φ being a bijective isomorphism of BMO is meant to include the assumption that f and its inverse are absolutely continuous with respect to n -dimensional measure. If this were not the case, the isomorphism φ could not be defined properly: if the zero set N were mapped onto a set N' of positive measure, then both $u' = 0$ and $u'' = \chi_{N'}$, the characteristic function of N' , were in BMO and both would satisfy

$$u' = u \circ f^{-1} \quad u'' = u \circ f^{-1}$$

with $u = 0$ a.e. in \mathbf{R}^n , $u \in \text{BMO}$.

Our first aim is to construct suitable functions $u \in \text{BMO}$.

LEMMA 6. (John-Nirenberg).

$$u(x) = \log^+ \frac{1}{|x|} = \begin{cases} \log \frac{1}{|x|} & |x| \leq 1 \\ 0 & |x| \geq 1 \end{cases}$$

is in BMO.

For a proof see [9].

LEMMA 7. Assume that g is a continuous function defined on \mathbf{R} with

$$k_g = \sup_{x \in \mathbf{R}} |g(x)| + \sup_{x, y \in \mathbf{R}} |g(x) - g(y)| \left(1 + \log^+ \frac{1}{|x - y|} \right) < \infty. \quad (3.2)$$

If $u \in \text{BMO}(\mathbf{R}^n)$ and if

$$k_u = \sup_{|Q| \geq 1} \left| \int_Q u(x) dx \right| < \infty,$$

then $v(x, y) = u(x)g(y) \in \text{BMO}(\mathbf{R}^{n+1})$.

Let $Q_r \subset \mathbf{R}^n$ denote the cube with side length r , centred at the origin. If $u \in \text{BMO}$, then for $r \leq 1$

$$|u_{Q_1} - u_{Q_r}| \leq (2^n + 1) \left(1 - \frac{\log r}{\log 2} \right) \|u\|_* \quad (3.3)$$

This can be seen as follows: Set $u_s = u_{Q_{2^{-s}}}$ $s = 0, 1, \dots$ Then

$$\begin{aligned} |u_s - u_{s-1}| &= 2^{ns} \int_{Q_{2^{-s}}} |u_s - u_{s-1}| dx \\ &\leq 2^{ns} \int_{Q_{2^{-s+1}}} |u(x) - u_{s-1}| dx + \|u\|_* \leq (2^n + 1) \|u\|_*, \end{aligned}$$

$$|u_s - u_{Q_1}| \leq \sum_{k=1}^s |u_k - u_{k-1}| \leq s(2^n + 1) \|u\|_*.$$

For $r = 2^{-s}$ this is equivalent with

$$|u_{Q_r} - u_{Q_1}| \leq (2^n + 1) \frac{-\log r}{\log 2} \|u\|_*.$$

For arbitrary r , $0 < r \leq 1$, inequality (3.3) can now easily be derived.

A cube $Q \subset \mathbf{R}^{n+1}$ with sides parallel to the coordinate axes can be represented as a direct product $Q = P \times S$ of cubes $P \subset \mathbf{R}^n$, $S \subset \mathbf{R}$. Set $a_Q = u_P g_0$, where g_0 is the value of g at the center of S . Then

$$\begin{aligned} \int_Q |v(x, y) - a_Q| \, dx \, dy \\ \leq \int_S |g(y)| \, dy \int_P |u(x) - u_P| \, dx + |u_P| \int_S |g(y) - g_0| \, dy. \end{aligned}$$

If $|Q| \geq 1$ this gives immediately

$$\int_Q |v(x, y) - a_Q| \, dx \, dy \leq k_g (\|u\|_* + k_u).$$

If $|Q| < 1$, we make use of (3.3) and (3.2) to conclude that

$$|u_P| \leq (2^n + 1) \left(1 - \frac{\log |P|}{n \log 2} \right) + k_u \|u\|_*$$

and

$$\int_S |g(y) - g_0| \, dy \leq k_g (1 - n^{-1} \log |P|)^{-1}.$$

Therefore

$$\int_Q |v(x, y) - a_Q| \, dx \, dy \leq 2k_g (k_u + 2^{n+1} \|u\|_*)$$

for any cube with sides parallel to the coordinate axes. If $Q \subset \mathbf{R}^{n+1}$ is an arbitrary cube, there exists a cube $Q_0 \supset Q$ with sides parallel to the coordinate axes and with

$$|Q_0| \leq (n+1)^{(n+1)/2} |Q|.$$

So the estimate

$$\int_Q |v - v_Q| \, dx \, dy \leq 2 \int_Q |v - a_{Q_0}| \, dx \, dy \leq 2(n+1)^{(n+1)/2} \int_{Q_0} |v - a_{Q_0}| \, dx \, dy$$

for the mean oscillation over Q shows that

$$\|v\|_* \leq 4(n+1)^{(n+1)/2} k_g (k_u + 2^{n+1} \|u\|_*).$$

As an application set $u(x) = \log^+ 1/|x|$, $x \in \mathbf{R}$ and let $g(y)$ be the piecewise linear, continuous odd function defined for $y \in \mathbf{R}$ by

$$g(y) = \begin{cases} 1 - |y - 1| & 0 \leq y \leq 2, \\ 0 & 2 \leq y, \\ -g(-y) & y \leq 0. \end{cases}$$

Since $|g(x) - g(y)| \leq \min\{2, |x - y|\}$, g satisfies the assumptions of Lemma 7 (with $k_g \leq 3$). From the Lemmata 6 and 7 we conclude that

$$v(x_1, x_2) = g(x_2) \log^+ \frac{1}{|x_1|}$$

is in BMO. For dimensions $n > 2$ let us define $v \in \text{BMO}$ by

$$v(x) = \log^+ 1/|x_1| \, h(x_2) \dots h(x_{n-1}) \, g(x_n)$$

with

$$h(x) = \begin{cases} 1 & |x| \leq 1, \\ 2 - |x| & 1 \leq |x| \leq 2, \\ 0 & |x| \geq 2, \end{cases}$$

v then has compact support and $v(x) = g(x_n) \log^+ 1/|x_1|$ for $|x_i| \leq 1$, $i = 1, \dots, n$. Finally, for $r > 0$ let v_r be defined by

$$v_r(x) = v\left(\frac{x}{r}\right).$$

Certainly $\|v_r\|_* = \|v\|_*$, since the space of BMO-functions is invariant under dilations.

With $a = (\alpha_1, \dots, \alpha_n) \in R_+^n$ (i.e. $\alpha_i > 0$, $i = 1, \dots, n$) we associate the sets $U_{a,r} = \{x : |x_i| \leq \alpha_i r, i = 1, \dots, n\}$ and the functions

$$\varphi_{a,r} = \begin{cases} |U_{a,r}|^{-1} v_r(x) & x \in U_{a,r} \\ 0 & \text{otherwise} \end{cases}$$

and

$$\psi_{a,r} = |\varphi_{a,r}|.$$

LEMMA 8. If $h \in L^1(\mathbb{R}^n)$, then there exists a sequence r_j converging to 0 such that a.e. in \mathbb{R}^n

$$\lim_{j \rightarrow \infty} \varphi_{a, r_j} * h(t) = \lim_{j \rightarrow \infty} \int_{\mathbb{R}^n} \varphi_{a, r_j}(x) h(t-x) dx = 0$$

and

$$\lim_{j \rightarrow \infty} \psi_{a, r_j} * h(t) = c_a h(t)$$

with

$$c_a = \int_{\mathbb{R}^n} \psi_{a, r} dx = \int_{U_{a, 1}} |v(x)| dx.$$

c_a is a continuous function of $a \in \mathbb{R}_+^n$. For $a = (\alpha_1, \dots, \alpha_{n-1}, 1)$, $\alpha_i \leq 1$, c_a can easily be calculated:

$$c_a = \frac{1}{2}(1 - \log \alpha_1). \quad (3.4)$$

For the proof of Lemma 8 consider the differences

$$d_1(t) = \varphi_{a, r} * h(t) - 0 = \int_{\mathbb{R}^n} \varphi_{a, r}(x) (h(t-x) - h(t)) dx$$

and

$$d_2(t) = \psi_{a, r} * h(t) - c_a h(t) = \int_{\mathbb{R}^n} \psi_{a, r}(x) (h(t-x) - h(t)) dx.$$

They satisfy

$$\lim_{r \rightarrow 0} \int_{\mathbb{R}^n} |d_k(t)| dt \leq \lim_{r \rightarrow 0} \int_{\mathbb{R}^n} \psi_{a, r}(x) \int_{\mathbb{R}^n} |h(t-x) - h(t)| dt dx = 0.$$

$k = 1, 2$ because

$$\lim_{x \rightarrow 0} \int_{\mathbb{R}^n} |h(t-x) - h(t)| dt = 0$$

and $\psi_{a, r}$ has its support in $U_{a, r}$. For some sequence r_j with $\lim_{j \rightarrow \infty} r_j = 0$ the differences d_1 and d_2 will therefore converge pointwise a.e.

For any rotation ϱ of \mathbb{R}^n set $\varphi_{\varrho, a, r}(x) = \varphi_{a, r}(\varrho^{-1}x)$, $U_{\varrho, a, r} = \varrho^{-1}(U_{a, r})$ and $v_{\varrho, r}(x) = v_r(\varrho^{-1}x) = v(\varrho^{-1}x/r)$. Observe that $\|v_{\varrho, r}\|_* = \|v\|_*$, since BMO is invariant under rotations. We think of ϱ as being given by an element in $O(n)$, the group of orthogonal $n \times n$ -matrices. As an immediate consequence of Lemma 8 let us note:

COROLLARY. Let $\{\varrho_i\}$ and $\{a_m\}$ be countable dense subsets of $O(n)$ and \mathbf{R}_+^n respectively. If $h \in L^1(\mathbf{R}^n)$, then for any pair ϱ_i, a_m there exists a sequence $\{r_j\}$ with $\lim_{j \rightarrow \infty} r_j = 0$ such that

$$\lim_{j \rightarrow \infty} \varphi_{\varrho_i, a_m, r_j} * h(t) = 0 \quad (3.5)$$

and

$$\lim_{j \rightarrow \infty} \psi_{\varrho_i, a_m, r_j} * h(t) = c_{a_m} h(t) \quad (3.6)$$

for all $t \in \mathbf{R}^n \setminus N$, where N is a set of measure zero which is independent of the pair ϱ_i, a_m .

The proof of Theorem 3 consists in showing that $\sup_{\xi \in \mathbf{R}^n, |\xi|=1} |F(x) \xi|^n \leq K \det F(x)$ a.e. in \mathbf{R}^n . ($F(x)$ is the Jacobian matrix of f at x .) This clearly is a local property. Based on our hypothesis and on the corollary to Lemma 8 we can assume that at $x=0$ the following conditions are satisfied:

- i) f is (totally) differentiable
- ii) $J_f = \det F \neq 0$, in view of the remark following Theorem 3
- iii) (3.5) and (3.6) hold for $h(x) = \begin{cases} J_f(x) & |x| < 1 \\ 0 & |x| \geq 1 \end{cases}$

F can be written in the form $F = \varrho D \sigma$, where $\varrho, \sigma \in O(n)$ and $D = \begin{pmatrix} \lambda_1 & 0 \\ & \ddots \\ 0 & \lambda_n \end{pmatrix}$ is a di-

agonal matrix with $\lambda_1 \geq \lambda_2 \dots \geq \lambda_n > 0$ (this is true for any $n \times n$ -matrix M with $\det M \neq 0$). Let us exclude the case $\det \sigma = -1$ by possibly interchanging the order of the coordinates. If we compose f with the rotation σ^{-1} , then the resulting mapping $g = \sigma^{-1} \circ f$ still satisfies the assumptions of Theorem 3 and the three additional conditions above. Note that $J_g = J_f$ and that the Jacobian matrix of g at 0 is $G = \varrho D$. The same is true if we consider the mapping cf , $c > 0$, instead of f (with $J_{cf} = c^n J_f$). There-

fore we can assume without loss of generality, that $F = \varrho D$ with $D = \begin{pmatrix} \lambda_1 & 0 \\ & \ddots \\ 0 & \lambda_n \end{pmatrix}$ and

$\lambda_1 \geq \lambda_2 \dots \geq \lambda_n = 1$. We then have to show that $\lambda_1^n \leq K \lambda_1 \dots \lambda_n$.

With this in mind let us choose ϱ_i and $a_m = (\alpha_{m1}, \dots, \alpha_{mn})$ in such a way that

$$|\lambda_k \alpha_{mk} - 1| < \varepsilon \quad k = 1, \dots, n \quad (3.7)$$

$$|c_{a_m} - \frac{1}{2}(1 - \log \alpha_{m1})| < \varepsilon \quad (3.8)$$

(cf. (3.4)) and such that for r small enough, say $r < \delta_1$, $U'_r = f U_{\varrho_i, a_m, r}$ contains the cube $S = \{z: |z_i| \leq r(1 - \varepsilon)\}$ and is contained in the cube $Q = \{z: |z_i| \leq r(1 + \varepsilon)\}$. We remind that $U_{\varrho, a, r}$ was defined by $U_{\varrho, a, r} = \varrho^{-1} \{x: |x_i| \leq r \alpha_i, i = 1, \dots, n\}$.

By the main hypothesis (3.1) $w_r = v_{\varrho, r} \circ f^{-1}$ is in BMO and $\|w_r\|_* \leq C \|v\|_*$. So with

$$\begin{aligned} w_Q &= \int_Q w_r(z) dz \\ \int_{U'_r} |w_r(z) - w_Q| dz &\leq |S|^{-1} \int_Q |w_r(z) - w_Q| dz \\ &\leq \left(\frac{1+\varepsilon}{1-\varepsilon}\right)^n \int_Q |w_r(z) - w_Q| dz \leq \left(\frac{1+\varepsilon}{1-\varepsilon}\right)^n C \|v\|_*. \end{aligned}$$

In this inequality we can replace w_Q by the mean value

$$\tilde{v}_r = \int_{U'_r} w_r(z) dz$$

if we instead write

$$\int_{U'_r} |w_r(z) - \tilde{v}_r| dz \leq 2 \left(\frac{1+\varepsilon}{1-\varepsilon}\right)^n C \|v\|_*.$$

Due to the absolute continuity of f with respect to n -dimensional Lebesgue measure

$$\begin{aligned} \tilde{v}_r &= \frac{|U_{\varrho_i, a_m, r}|}{|U'_r|} \int_{U_{\varrho_i, a_m, r}} v_{\varrho_i, r}(x) J_f(x) dx \\ &= \frac{|U_{\varrho_i, a_m, r}|}{|U'_r|} \varphi_{\varrho_i, a_m, r} * h(0) \end{aligned}$$

so by the corollary to Lemma 6

$$\lim_{j \rightarrow \infty} \tilde{v}_{r_j} = 0.$$

Hence there exists $\delta_2 > 0$ ($\delta_2 \leq \delta_1$) such that for $r = r_j < \delta_2$

$$\int_{U'_r} |w_r(z)| dz \leq \varepsilon + 2 \left(\frac{1+\varepsilon}{1-\varepsilon}\right)^n C \|v\|_*. \quad (3.9)$$

On the other side

$$\int_{U'_r} |w_r(z)| dz = \frac{|U_{\varrho_i, a_m, r}|}{|U'_r|} \psi_{\varrho_i, a_m, r} * h(0)$$

so by (3.6)

$$\lim_{j \rightarrow \infty} \int_{\tilde{U}'_{r_j}} |w_{r_j}(z)| dz = (J_f(0))^{-1} c_{a_m} J_f(0) = c_{a_m}. \quad (3.10)$$

Combining the two results (3.9) and (3.10) with (3.8) we conclude that for j big enough

$$\frac{1}{2} (1 - \log \alpha_{m1}) - \varepsilon \leq \int_{\tilde{U}'_{r_j}} |w_{r_j}(z)| dz \leq \varepsilon + 2 \left(\frac{1 + \varepsilon}{1 - \varepsilon} \right)^n C \|v\|_*.$$

Together with (3.7) this implies

$$1 + \log \lambda_1 \leq 4 C \|v\|_*$$

since $\varepsilon > 0$ was arbitrary. The inequality $\lambda_1^n \leq K \lambda_1 \dots \lambda_n$ therefore holds with $K = e^{(n-1)(4C\|v\|_*-1)}$.

4. A Local Version

The object of this section is a local version of the Theorems 2 and 3. If G is a domain in \mathbf{R}^n we denote by $\text{BMO}(G)$ the subspace of BMO consisting of all functions $u \in \text{BMO}$ with support in G (the support of a locally integrable function $u \in L^1_{\text{loc}}(\mathbf{R}^n)$ is the complement of the largest open set $O \subset \mathbf{R}^n$ with $u(x) = 0$ a.e. on O).

LEMMA 9. *If $u \in L^1_{\text{loc}}(\mathbf{R}^n)$ and if $\text{supp } u \subset G$, then*

$$\|u\|_* \leq 4 \sup_P \int_P |u(x) - u_P| dx,$$

where the supremum is extended over all cubes P with $\text{dia } P \leq 4n^{1/2} \text{ dia } G$.

G is contained in a ball B with radius $\text{dia } G$. If $Q \cap G \neq \emptyset$ for some cube with $\text{dia } Q \geq 4n^{1/2} \text{ dia } G$, then there exists a cube $P \subset Q$ with side length $n^{-1/2} \text{ dia } P = 2 \text{ dia } B$ such that $P \cap B = Q \cap B \supset Q \cap G$. With $N = \{x \in P : u(x) = 0\}$, $|N| \geq |P| - |G| \geq (1 - 2^{-n})|P|$, we obtain

$$\int_P |u - u_P| dx = |P|^{-1} \int_N |u_P| dx + |P|^{-1} \int_{P \setminus N} |u - u_P| dx$$

which shows that

$$|u_P| \leq (1 - 2^{-n})^{-1} \int_P |u - u_P| dx.$$

Finally

$$\begin{aligned} \int_Q |u - u_Q| dx &\leq 2 \int_Q |u - u_P| dx \\ &\leq 2|Q|^{-1} \int_{Q \setminus P} |u_P| dx + 2|Q|^{-1} \int_P |u - u_P| dx \\ &\leq 2(1 - 2^{-n})^{-1} \int_P |u - u_P| dx \end{aligned}$$

and the proof is complete.

THEOREM 4. *Assume that $f: G \rightarrow \mathbb{R}^n$ is a (orientation preserving) homeomorphism, $f \in \text{ACL}$, and that f is differentiable a.e. Then f is a quasiconformal mapping if and only if every point $x \in G$ has a neighbourhood U such that $\varphi: u \rightarrow u' = u \circ f^{-1}$ is an isomorphism of $\text{BMO}(U)$ onto $\text{BMO}(fU)$ which satisfies*

$$\|u'\|_* \leq C_0 \|u\|_*$$

for all $u \in \text{BMO}(U)$ with a fixed constant C_0 independent of U .

The proof for the quasiconformality of a homeomorphism $f: G \rightarrow \mathbb{R}^n$ satisfying all the above hypotheses is contained in the proof of Theorem 3. In order to show that a quasiconformal mapping gives rise to local isomorphisms of $\text{BMO}(U)$ onto $\text{BMO}(U')$, $U' = fU$, the full strength of the Lemmata 2 and 4 has to be used.

For $x \in G$ we choose a neighbourhood U in such a way that

$$4n^{1/2}(1+2k) \text{dia } U' < \text{dist}(U', \partial G'),$$

where k is the constant of Lemma 4. If P' is a cube with $\text{dia } P' \leq 4n^{1/2} \text{dia } U'$ and with $P' \cap U' \neq \emptyset$, then

$$\text{dist}(P', \partial G') \geq \text{dist}(U', \partial G') - \text{dia } P' > \text{dia } P'(1+2k) - \text{dia } P' = 2k \text{dia } P'.$$

Hence by Lemma 4 there exists a cube $Q \subset G$ with $fQ = Q' \supset P'$, $|Q'| \leq k^n n^{n/2} |P'|$ and with

$$\text{dist}(Q', \partial G') > \text{dia } Q'. \quad (4.1)$$

For a given function $u \in \text{BMO}(U)$ with $u' = u \circ f^{-1}$ we can then proceed as in the proof of Theorem 2. Condition (4.1) ensures the validity of Lemma 2. It follows (see (2.11)) that

$$\int_{P'} |u'(z) - u'_{P'}| dz \leq C \|u\|_*$$

and this inequality holds for all P' with $\text{dia } P' \leq 4n^{1/2} \text{dia } U'$. Therefore by Lemma 9

$$\|u'\|_* \leq 4C \|u\|_* = C_0 \|u\|_*.$$

REFERENCES

- [1] BOJARSKI, B. V., *Homeomorphic solutions of Beltrami systems*, (Russian), Dokl. Akad. Nauk S.S.S.R. 102 (1955), 661–664.
- [2] CARAMAN, P., *Homeomorfisme cvasiconforme n-dimensionale*, Bucuresti, 1968.
- [3] FEFFERMAN, C. and STEIN, E. M., *H^p spaces of several variables*, Acta Math. 129 (1972), 137–193.
- [4] GEHRING, F. W., *Symmetrization of rings in space*, Trans. Amer. Math. Soc. 101 (1961), 499–519.
- [5] —, *The L^p -integrability of the partial derivatives of a quasi-conformal mapping*, Acta Math. 130 (1973), 265–277.
- [6] GEHRING, F. W. and REICH, E., *Area distortion under quasiconformal mappings*, Ann. Ac. Sc. Fenn. 388 (1966), 1–15.
- [7] JOHN, F., *Rotation and strain*, Comm. Pure Appl. Math. 14 (1961), 319–413.
- [8] JOHN, F. and NIRENBERG, L., *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. 14 (1961), 415–426.
- [9] MOSER, J., *On Harnack's theorem of elliptic differential equations*, Comm. Pure Appl. Math. 14 (1961), 577–591.
- [10] MOSTOW, G. D., *Quasi-conformal mappings in n -space and the rigidity of hyperbolic space forms*, IHES, Publ. Math. 34 (1968), 53–104.
- [11] STAMPACCHIA, G., *The spaces $\mathcal{L}^{(p, \lambda)}$, $N^{(p, \lambda)}$ and interpolation*, Ann. Sc. Norm. Sup. Pisa 19 (1965), 443–462.
- [12] STEIN, E. M., *Singular integrals, harmonic functions and differentiability properties of functions of several variables*, Proc. Symp. Pure Math. 10 (1967), 316–335.
- [13] STEIN, E. M. and ZYGMUND, A., *Boundedness of translation invariant operators on Hölder spaces and L^p -spaces*, Ann. Math. 85 (1967), 337–349.
- [14] VÄISÄLÄ, J., *Two new characterizations for quasiconformality*, Ann. Ac. Sc. Fenn. 362 (1965), 1–12.

University of Bern
Switzerland

Received December 22, 1973