

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 49 (1974)

**Artikel:** On the genus of finite CW-H-spaces  
**Autor:** Zabrodsky, A.  
**DOI:** <https://doi.org/10.5169/seals-37978>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 06.02.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# On the genus of finite CW–H-spaces

A. ZABRODSKY

The study of the Grothendieck groups of certain stable categories (initiated in [1] and [2] and pursued in [3]) led to the study of two phenomena: That of non cancellation [4] and that of the genus of a space [7]. Their inter-relations were observed and their study led to the discovery of new finite CW–H-spaces. Our main reference in this study of the genus of an H-space should be [7] but many of the facts studied here are closely related to those studied in [1], [4] and [6]. Here we use the notion of genus with regard to  $p$ -equivalence in the sense of Serre ([5]) rather than that of localization which are equivalent for H-spaces (see [7]).

## 0. Notations, Definitions and Summary of Results

As usual we denote by  $PA$  and  $QA$  the modules of primitives and indecomposables in a Hopf algebra  $A$ .

We denote by  $\mathbf{P}$  the set of all primes. All spaces considered are of the homotopy type of simply connected CW complexes of finite type, all graded modules, accordingly, are finitely generated in each dimension.

Let  $p \in \mathbf{P}$ . A homomorphism  $\varphi: G \rightarrow G'$  between two finitely generated abelian groups is said to be a  $p$ -epimorphism ( $p$ -monomorphism) if  $\text{coker } \varphi$  ( $\ker \varphi$ ) is a finite group with order prime to  $p$ . If  $\varphi$  is both  $p$ -epimorphism and  $p$ -monomorphism then it is said to be  $p$ -isomorphism. If  $\mathbf{P}_1 \subset \mathbf{P}$  then  $\varphi$  is said to be a  $\mathbf{P}_1$ -epimorphism ( $\mathbf{P}_1$ -monomorphism or  $\mathbf{P}_1$ -isomorphism) if it is a  $p$ -epimorphism ( $p$ -monomorphism or  $p$ -isomorphism) for every  $p \in \mathbf{P}_1$ .

A map  $\varphi: Y \rightarrow X$  is said to be a  $\mathbf{P}_1$  equivalence iff  $\pi_*(\varphi)$  (equivalently  $H_*(\varphi, Z)$ ,  $H^*(\varphi, Z)$ ) is a  $\mathbf{P}_1$  isomorphism. By the genus of the space  $X$  we mean the set  $G(X)$  of homotopy classes of spaces  $Y$  which are  $p$ -equivalent to  $X$  for every  $p \in \mathbf{P}$ .

If  $X$  is a CW complex the homotopy (or Postnikov) approximation of  $X$  in  $\dim \leq m$  is a pair  $(Ht_m(X), \tau_m)$  where  $\tau_m: X \rightarrow Ht_m(X)$  is such that  $\pi_k(\tau_m)$  is an isomorphism for  $k \leq m$  and  $\pi_k(Ht_m(X)) = 0$  for  $k > m$ . Similarly, the homology (or Moore) approximation of  $X$  is a pair  $(Hl_m(X), \tau'_m)$  where  $\tau'_m: Hl_m(X) \rightarrow X$  yields an isomorphism  $H_k(Hl_m(X), Z) \rightarrow H_k(X, Z)$  for  $k \leq m$  and  $H_k(Hl_m(X), Z) = 0$  for  $k > m$ .

Though we work in the category of CW complexes and continuous maps by a commutative diagram we mean commutative up to homotopy.

A commutative diagram

$$\begin{array}{ccc}
 X & \xrightarrow{f_2} & Y_2 \\
 \downarrow h_2 & & \downarrow h_1 \\
 Y_1 & \xrightarrow{f_1} & X_0
 \end{array} \tag{0.1}$$

is said to be a fiber square induced by  $f_1, h_1$  if up to homotopy one has

$$X = \{y_1, y_2, \varphi \in Y_1 \times Y_2 \times PX_0 \mid f_1(y_1) = \varphi(0), h_1(y_2) = \varphi(1)\}.$$

$$h_2 = p_1 \mid X, \quad f_2 = p_2 \mid X$$

where

$$p_i: Y_1 \times Y_2 \times PX_0 \rightarrow Y_i \quad \text{are the projections.}$$

(0.1) is said to be a fiber square in  $\dim \leq m$  iff it becomes a fiber square after applying  $Ht_m$  to the entire diagram. If  $X$  is a finite dimensional H-space then  $H^*(X, Q) = \Lambda(\tilde{x}_{n_1}, \tilde{x}_{n_2}, \dots, \tilde{x}_{n_r})$ ,  $\tilde{x}_{n_i} \in H^{n_i}(X, Q)$ ,  $n_i \leq n_{i+1}$ .  $X$  is then said to be of type  $(n_1, n_2, \dots, n_r)$ . Given a vector  $\bar{n} = (n_1, n_2, \dots, n_r)$  of natural numbers we write  $K(Z, \bar{n}) = \prod_{i=1}^r K(Z, n_i)$ ,  $S^{\bar{n}} = \prod_{i=1}^r S^{n_i}$ .

In this study we consider  $G(X)$  for a finite dimensional H-space. Using exactly the same (or dual) methods one can obtain similar (or dual) results for H-spaces with finitely many non-vanishing homotopy groups (or for finite dimensional co-H-spaces).

To state results of this study concerning the structure of  $G(X)$  one needs few notations: If  $A$  and  $B$  are matrices (over any ring  $R$ ) denote by  $A * B$  the matrix

$$\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$$

Let  $X$  be an H-space of type  $(n_1, n_2, \dots, n_r)$ .  $\text{Hom } \mathcal{Z}(QH^*(X, Z)/\text{torsion}, QH^*(X, Z)/\text{torsion})$  can be identified with the set of all matrices (over  $Z$ ) of the form  $A_{s_1} * A_{s_2} * \dots * A_{s_l}$ ,  $A_{s_i}$  being an  $s_i \times s_i$  matrix,  $s_1, s_2, \dots, s_l$  uniquely determined by  $n_1, n_2, \dots, n_r$ . Denote all such matrices by  $\mathcal{M}(s_1, s_2, \dots, s_l; Z)$ . Given an integer  $t$  let  $\mathcal{A} = \mathcal{A}_t$  be the set of all matrices in  $\mathcal{M}(s_1, s_2, \dots, s_l; Z)$  which are invertible mod  $Z_t$ . Then one has:

*Theorem (essentially 2.4).* There exists an integer  $t$  ( $= t_k$  in 2.4) and a correspondence  $\xi: \mathcal{A} = \mathcal{A}_t \rightarrow G(X)$  described as follows: Fix a map  $h_0: X \rightarrow K(Z, \bar{n})$  so that  $QH^*(h_0, Z)/\text{torsion}$  is an isomorphism. Given  $A \in \mathcal{A}$  let  $f_0 = f_0(A): K(Z, \bar{n}) \rightarrow K(Z, \bar{n})$

be given by  $H^*(f_0, Z)$   $\iota_{n_i} = \sum_j a_{ij} \iota_{n_j}$   $A = (a_{ij})$ . If

$$\begin{array}{ccc} \tilde{Y} & \xrightarrow{f_1} & X \\ \downarrow h_1 & & \downarrow h_0 \\ K(Z, \bar{n}) & \xrightarrow{f_0} & K(Z, \bar{n}) \end{array}$$

is a fiber square then

- (a)  $[Hl_k(\tilde{Y})] = [Y] = \xi(A) \in G(X)$  ( $k = \dim X$ )
- (b)  $QH^*(h_1, Z)$ /torsion is an isomorphism where  $h_1: Y \rightarrow K(Z, \bar{n})$  is induced by  $\tilde{h}: h_1 = \tilde{h}_1 \tau'$ .

*Corollary (2.6)*  $\xi$  is onto.

If  $\tilde{\alpha}: \mathcal{M}(s_1, s_2, \dots, s_l; Z) \rightarrow \mathcal{M}(s_1, s_2, \dots, s_l; Z_t)$  is the reduction it induces homomorphism

$$\bar{\alpha}: GL(s_1, \dots, s_l; Z) \rightarrow GL(s_1, s_2, \dots, s_l, Z_t)$$

of invertible matrices and  $(\text{coker } \bar{\alpha}) = [(Z_t^*)/\{\pm 1\}]^l$  where  $Z_t^*$  are the units in  $Z_t$ . Note that  $\mathcal{A} = \tilde{\alpha}^{-1} GL(s_1, s_2, \dots, s_l; Z_t)$ .

*Structure Theorem (2.7)*  $\xi$  factors through  $\text{coker } \bar{\alpha}$  to obtain a correspondence  $\xi: [Z_t^*/\{\pm 1\}]^l \rightarrow G(X)$  which is onto.

The correspondence is given by

$$\xi(d_1, d_2, \dots, d_l) = \xi(\xi_{d_1} * I_{d_2} * \dots * I_{d_l})$$

where  $I_{d_i}$  is the  $s_i \times s_i$  matrix given by

$$I_{d_i} = \begin{pmatrix} I & 0 \\ 0 & d_i \end{pmatrix}$$

One also obtains:

*H-structure theorem (2.10, compare with [6] and [7] lemma 1.4):*

Let  $X$  be a finite CW complex. If for every prime  $p$  there exists an H-space  $X(p)$  and a  $p$ -equivalence  $f_p: X \rightarrow X(p)$  and if  $H^*(X(p), Q)$  are all isomorphic as Hopf algebras then  $X$  is an H-space.

*A Product Theorem (2.11).* Let  $A, A' \in \mathcal{A}$ . Then

$$\xi(A) \times \xi(A') = \xi(I) \times \xi(A \cdot A') = X \times \xi(A \cdot A').$$

Consequently if  $A_i, B_i \in \mathcal{A}$  and

$$A_1 \cdot A_2 \cdots \cdot A_m = B_1 \cdot B_2 \cdots \cdot B_m$$

then

$$\xi(A_1) \times \xi(A_2) \cdots \times \xi(A_m) \approx \xi(B_1) \times \xi(B_2) \cdots \times \xi(B_m).$$

*Theorem (2.12).* If  $[Y] \in G(X)$  then

$$Y^{\varphi(t)/2} \approx X^{\varphi(t)/2}$$

where  $\varphi$  is the Euler function. Finally, one obtains

*A non cancellation theorem (3.5 and compare with [1], [2] and [7], p. 83).* If  $[Y] \in G(X)$  then  $Y \times S^{\bar{n}} \approx X \times S^{\bar{n}}$ .

## 1. Some Algebraic Lemmas

The following are well known simple facts concerning finitely generated abelian groups:

1.1. LEMMA: *Let  $G$  and  $G'$  be finitely generated abelian groups. If for every prime  $p$  there exists a  $p$ -isomorphism  $\varphi_p: G \rightarrow G'$  then  $G$  and  $G'$  are isomorphic.*

1.2. LEMMA: *Suppose  $G \approx G'$ . Then  $\varphi: G \rightarrow G'$  is a  $p$ -isomorphism if and only if  $\varphi \otimes \mathbb{Z}_p$  is an isomorphism.*

1.3. COROLLARY. *Let  $\{\varphi_p \mid p \in \mathbf{P}_1\}$  be a finite family,  $\varphi_p: G \rightarrow G' \approx G$  a  $p$ -isomorphism. Put  $a = \prod_{p \in \mathbf{P}_1} p$ . Then  $\varphi = \sum_{p \in \mathbf{P}_1} (a \cdot p^{-1}) \varphi_p$  is a  $\mathbf{P}_1$  isomorphism.*

*Proof.* For every  $p_0 \in \mathbf{P}_1$   $\varphi \otimes \mathbb{Z}_{p_0} = a p_0^{-1} (\varphi_{p_0} \otimes \mathbb{Z}_{p_0})$  is an isomorphism. Hence  $\varphi$  is a  $\mathbf{P}_1$  isomorphism.

1.4. COROLLARY. *If  $[Y] \in G(X)$  then*

(a)  $H^*(Y) \approx H^*(X)$  and  $\pi(X) \approx \pi(Y)$ .

(b) *Let  $f: Y \rightarrow X$ . If  $\pi(f) \otimes \mathbb{Z}_p$  is an isomorphism for all  $p \in \mathbf{P}_1$  then  $f$  is a  $\mathbf{P}_1$  equivalence.*

Throughout this chapter let  $X$  be a finite CW-H-space.

1.5. PROPOSITION. *If  $[Y] \in G(X)$  then there exists a partition  $\mathbf{P} = \mathbf{P}_1 \cup \mathbf{P}_2$  and two maps  $f_i: Y \rightarrow X$   $i = 1, 2$  so that  $f_i$  is a  $\mathbf{P}_i$  equivalence.*

*Proof.* Choose an arbitrary prime  $q \in \mathbf{P}$ . Let  $f_2: Y \rightarrow X$  be a  $q$ -equivalence. Then there exists a set  $\mathbf{P}_2$  of primes so that  $\mathbf{P}_1 = \mathbf{P} - \mathbf{P}_2$  is finite and  $f_2$  is a  $\mathbf{P}_2$  equivalence. For every  $p \in \mathbf{P}_1$  let  $f_p$  be a  $p$ -equivalence. Let  $a$  be the product of all primes in  $\mathbf{P}_1$ . If  $f_1: Y \rightarrow X$  is given by  $[f_1] = \prod_{p \in \mathbf{P}_1} [f_p]^{a/p}$  (where the product represents a product in

the algebraic loop  $[Y, X]$  with an arbitrary bracketing) then by 1.3  $\pi_n(f_1) = \sum_{p \in \mathbf{P}_1} (ap^{-1}) \pi_n(f_p)$  is a  $\mathbf{P}_1$  isomorphism and  $f_1$  is a  $\mathbf{P}_1$  equivalence.

One notices that  $\mathbf{P}_1$  may be replaced by any larger (but finite) set of primes.

1.6. LEMMA. *Given a commutative diagram*

$$\begin{array}{ccc} T & \xrightarrow{h_2} & Y_1 \\ \downarrow h_1 & & \downarrow f_1 \\ Y_2 & \xrightarrow{f_2} & X_0 \end{array} \quad (1.6.1)$$

If for  $i=1, 2$   $h_i$  and  $f_i$  are  $\mathbf{P}_i$  equivalences in  $\dim \leq m$ ,  $\mathbf{P} = \mathbf{P}_1 \cup \mathbf{P}_2$  then (1.6.1) is a fiber square in  $\dim \leq m$ .

*Proof.* Form the fiber square of  $f_1$  and  $f_2$

$$\begin{array}{ccc} Z' & \xrightarrow{h'_2} & X \\ \downarrow h'_1 & & \downarrow f_1 \\ Y & \xrightarrow{f_2} & X_0 \end{array}$$

then there exists  $\alpha: Z \rightarrow Z'$  with  $h_i \sim h'_i \alpha$   $i=1, 2$ .  $f_i$  being  $\mathbf{P}_i$  equivalence in  $\dim \leq m$  implies that  $h'_i$  is a  $\mathbf{P}_i$  equivalence and as  $h_i$  is a  $\mathbf{P}_i$  equivalence so is  $\alpha$ . Now,  $\alpha$  being  $\mathbf{P}_1$  and  $\mathbf{P}_2$  equivalence in  $\dim \leq m$  is a homotopy equivalence in that range.

Let  $R$  be a ring. If  $M = \{M_n \mid 0 \leq n \leq l\}$  is a graded module over  $R$ ,  $M_n$ -free of rank  $s_n$  then  $\mathcal{M}(R) = \text{Hom}_R^\circ(M, M)$  can be represented by the set of matrices of the form  $A_1 * A_2 * \dots * A_l$  where  $A_n$  is  $s_n \times s_n$  matrix over  $R$  and  $B_1 * B_2$  is the matrix of the form

$$\begin{pmatrix} B_1 & 0 \\ 0 & B_2 \end{pmatrix}$$

Denote this set by  $\mathcal{M}(s_1, s_2, \dots, s_l; R)$ . Similarly define  $GL(s_1, s_2, \dots, s_l; R)$  and  $SL(s_1, s_2, \dots, s_l; R)$  as the invertible and determinant one matrices in  $\mathcal{M}(s_1, s_2, \dots, s_l; R)$ . The product  $*g$  in  $M(s_1, s_2, \dots, s_l; R)$  given by

$$(A_1 * A_2 * \dots * A_l) * g (\hat{A}_1 * \hat{A}_2 * \dots * \hat{A}_l) = (A_1 * \hat{A}_1) * (A_2 * \hat{A}_2) * \dots * (A_l * \hat{A}_l)$$

corresponds to the natural homeomorphism  $\text{Hom}^\circ(M, M) \otimes \text{Hom}^\circ(M, M) \rightarrow \text{Hom}^\circ(M \otimes M, M \otimes M)$ . Let  $X$  be of type  $(n_1, n_2, \dots, n_r)$ . Define  $s_1, s_2, \dots, s_l$  by the relation

$$n_{s_1+s_2+\dots+s_m} < n_{s_1+s_2+\dots+s_m+1} = n_{s_1+s_2+\dots+s_m+s_{m+1}} \quad 0 \leq m < l \quad (1.7)$$

Put  $M_m = QH_{s_1+s_2+\dots+s_m}^n(X, Z)/\text{torsion}$  and one has

$$\text{Hom}^\circ[QH^*(X, Z)/\text{torsion}, QH^*(X, Z)/\text{torsion}] = \mathcal{M}(s_1, s_2, \dots, s_l; Z).$$

Let  $t$  be an integer. Put

$$\alpha: GL(n, \mathbb{Z}) \rightarrow GL(n, \mathbb{Z}_t)$$

Then one has

1.8. LEMMA.  $GL(n, \mathbb{Z}_t)/\text{im } \alpha = \mathbb{Z}_t^*/\{\pm 1\}$  where  $\mathbb{Z}_t^*$  are the units in  $\mathbb{Z}_t$ . The homomorphism  $GL(n, \mathbb{Z}_t) \rightarrow GL(n, \mathbb{Z}_t)/\text{im } \alpha$  is given by the determinant.

Let  $\tilde{\alpha}: \mathcal{M}(s_1, s_2, \dots, s_l; \mathbb{Z}) \rightarrow \mathcal{M}(s_1, \dots, s_l; \mathbb{Z}_t)$  induce

$$\bar{\alpha}: GL(s_1, s_2, \dots, s_l; \mathbb{Z}) \rightarrow GL(s_1, s_2, \dots, s_l; \mathbb{Z}_t)$$

then

$$GL(s_1, s_2, \dots, s_l; \mathbb{Z}_t)/\text{im } \bar{\alpha} \approx [\mathbb{Z}_t^*/\{\pm 1\}]^l.$$

## 2. The Structure of $G(X)$

Throughout this chapter let  $(X, \mu)$  be an H-space of type  $\bar{n} = (n_1, n_2, \dots, n_r)$ . To this one associate the sequence  $(s_1, s_2, \dots, s_l)$  defined by (1.7). Let  $\dim X \leq k$ . Given an integer  $m$ ,  $m \geq k$  let  $t_m = t_m(G(X))$  be the order of the finite group

$$\sum_{n \leq m} (\ker \sigma_n + \text{coker } \sigma_n)$$

where  $\sigma_n: \pi_n(X) \rightarrow PH_n(X)/\text{torsion}$  is the Hurewicz-Serre homomorphism.

Another interpretation for  $t_m$  can be given by the following

2.1. LEMMA. Let  $h_0: X \rightarrow K(Z, \bar{n}) = \prod_i K(Z, n_i)$  satisfy:  $x_{n_i} = H^*(h_0, Z)_{n_i}$ ,  $1 \leq i \leq r$ , represent a basis for  $QH^*(X, Z)/\text{torsion}$ . Let  $F = \text{fiber } h_0$ . Then  $\text{order } \pi_n(F) = \text{order } (\ker \sigma_n + \text{coker } \sigma_{n+1})$ . Hence,  $t_m = \text{order } \sum_{n \leq m} \pi_n(F)$ .

(Note that  $\text{coker } \sigma_{m+1} = PH_{m+1}(X, Z)/\text{torsion} = 0$ ).

*Proof.* One has the following commutative diagram

$$\begin{array}{ccc} \pi_n(X) & \xrightarrow{\pi_n(h_0)} & \pi_n(K(Z, \bar{n})) \\ \downarrow \sigma_n(X) & & \approx \downarrow \sigma_n(K) \\ PH_n(X, Z)/\text{torsion} & \xrightarrow[\approx]{h_0*} & PH_n(K(Z, \bar{n}), Z)/\text{torsion} \end{array}$$

As  $QH^*(h_0, Z)/\text{torsion}$  is an isomorphism so is its dual  $h_0*$ . Hence,  $\ker \sigma_n = \ker \pi_n(h_0)$  and  $\text{coker } \sigma_n = \text{coker } \pi_n(h_0)$  and obviously  $\text{order } \pi_n(F) = \text{order } [\ker \pi_n(h_0) + \text{coker } \pi_{n+1}(h_0)]$ .

2.2. PROPOSITION. Given a set  $\{x_{n_1}, x_{n_2}, \dots, x_{n_r}\}$ ,  $x_{n_i} \in H^{n_i}(X, Z)/\text{torsion}$  which

reduces to a basis for  $QH^*(X, Z)/\text{torsion}$  and given maps  $g: Y \rightarrow \tilde{Y}, f: Y \rightarrow X$ , ( $Y$  and  $\tilde{Y}$  arbitrary CW complexes of  $\dim \leq m$ ) so that  $\text{im}[\tilde{H}(g, Z)/\text{torsion}]$  is an ideal in  $\tilde{H}(Y, Z)/\text{torsion}$ . For any set  $\{z_{n_1}, \dots, z_{n_r}\}$ ,  $z_{n_i} \in H^{n_i}(\tilde{Y}, Z)/\text{torsion}$  there exist maps  $\tilde{f}_i: \tilde{Y} \rightarrow X$ ,  $i = 1, 2, \dots, r$  with torsion  $\pi_n(\tilde{Y}) \subset \ker \pi_n(\tilde{f}_i)$  for  $n \leq m$  so that

$$[H^*(\tilde{f}, Z)/\text{torsion} - H^*(f, Z)/\text{torsion}] x_{n_i} = t_m [(H^*(g, Z)/\text{torsion}) z_{n_i}] \quad (2.2.1)$$

where  $\tilde{f}$  is given by

$$[\tilde{f}] = [\tilde{f}_r \circ g] ([f_{r-1} \circ g] \dots ([\tilde{f}_2 \circ g] ([\tilde{f}_1 \circ g] \cdot [f]) \dots))$$

(the products taken in  $[Y, X]$ ).

*Proof.* Let  $X_m = Ht_m(X)$ . As  $\dim Y \leq m$ ,  $\dim \tilde{Y} \leq m$  and  $m \geq k$  one can replace all maps in  $[Y, X]$  and  $[\tilde{Y}, X]$  by the corresponding maps in  $[Y, X_m]$  and  $[\tilde{Y}, X_m]$ . Consider  $x_{n_i}$  being in  $H^{n_i}(X_m, Z)/\text{torsion}$  and it suffices to prove relation (2.2.1) for  $X_m$  instead of  $X$ . By 2.1  $t_m = \text{order } \pi_*(F_m)$  ( $F_m$  the fiber of  $X_m \rightarrow K(Z, \bar{n})$ ). Hence,  $t_m[1] ([1] \in [K(Z, \bar{n}), K(Z, \bar{n})])$  lifts to  $h_1: K(Z, \bar{n}) \rightarrow X$ ,  $[h_0 \circ h_1] = t_m[1]$ . We construct  $\tilde{f}_i$  inductively. Suppose  $\tilde{f}_1, \tilde{f}_2, \dots, \tilde{f}_{i-1}$  where constructed so that if

$$[\tilde{f}_{i-1}] = [\cdot_{i-1} \circ g] ([\tilde{f}_{i-2} \circ g] (\dots ([\tilde{f}_1 \circ g] \cdot [f])))$$

then

$$[H^*(\tilde{f}_{i-1}, Z)/\text{torsion} - H^*(f, Z)/\text{torsion}] x_{n_j} = \begin{cases} t_m(H^*(g, Z)/\text{torsion}) z_{n_j} & j < i \\ t_m(H^*(g, Z)/\text{torsion}) \tilde{z}_{n_j} & j \geq i \end{cases}$$

(there is no problem in starting the induction by starting with  $\tilde{f}_0: \tilde{Y} \rightarrow X$  being the constant map and  $\tilde{z}_{n_j} = 0$ ). Let  $\tilde{f}_i$  be the composition

$$\tilde{Y} \xrightarrow{z_{n_i} - \tilde{z}_{n_i}} K(Z, n_i) \xrightarrow{\text{inj}} K(Z, \bar{n}) \xrightarrow{h_1} X.$$

Then torsion  $\pi(\tilde{Y}) \subset \ker \pi(\tilde{f}_i)$ . Let  $(\ )^*$  denote  $H^*(\ , Z)/\text{torsion}$ . If  $[\tilde{f}_i] = [\tilde{f}_i \circ g] [\tilde{f}_{i-1}]$  then

$$\tilde{f}_i^* x_{n_j} = \Lambda^*(\tilde{f}_i \circ g \times \tilde{f}_{i-1})^* \mu^* x_{n_j} = \sum_{s=0}^{s_0} (\tilde{f}_i \circ g)^* x'_{n_j, s} \cdot \tilde{f}_{i-1}^* x''_{n_j, s}$$

where  $x'_{n_j, s}$  and  $x''_{n_j, s}$  are given by  $\mu^* x_{n_j} = \sum_{s=0}^{s_0} x'_{n_j, s} \otimes x''_{n_j, s}$  ( $x'_{n_j, 0} = x_{n_j} = x''_{n_j, s_0}$ ,  $x''_{n_j, 0} = 1 = x'_{n_j, s_0}$  and for  $0 < s < s_0$ ,  $0 < \dim x'_{n_j, s} < n_j$ ,  $0 < \dim x''_{n_j, s} < n_j$ ).

As  $H^n(\tilde{f}_i \circ g, Z)/\text{torsion} = 0$  for  $n < n_i$  and  $\tilde{f}_i^* x_{n_i} = \tilde{f}_i^* h_0^* \iota_{n_i} = t_m(z_{n_i} - \tilde{z}_{n_i})$  one has  $\tilde{f}_i^* x_{n_j} = \tilde{f}_{i-1}^* x_{n_j}$  if  $j < i$ ,

$$\tilde{f}_i^* x_{n_i} = \tilde{f}_{i-1}^* x_{n_i} + t_m g^*(z_{n_i} - \tilde{z}_{n_i})$$

and hence for  $j \leq i$

$$(\hat{f}_i^* - f^*) x_{n_j} = (\hat{f}_i^* - \hat{f}_{i-1}^*) x_{n_j} + (\hat{f}_{i-1}^* - f^*) x_{n_j} = t_m g^* z_n$$

For  $j > i$  (as  $\tilde{f}_i^* x_{n_j} = 0$  for  $j \neq i$ ) one has

$$\alpha_j = (\hat{f}_i^* - \hat{f}_{i-1}^*) x_{n_j} = \sum_{s=1}^{s_0-1} \hat{f}_{i-1}^* x_{n_j, s} (\tilde{f}_i \circ g)^* x'_{n_j, s}$$

and as  $\text{im}(\tilde{f}_i \circ g)^* \subset \overline{t_m(\text{im} g^*)}$  and as  $\overline{\text{im} g^*}$  is an ideal  $\alpha_j \in t_m \overline{\text{im} g^*}$  and hence for  $i > j$   $(\hat{f}_i^* - f^*) x_{n_j} = (\hat{f}_i^* - \hat{f}_{i-1}^*) x_{n_j} + t_m g^* \tilde{z}_{n_j} = t_m g^* \tilde{z}'_{n_j}$  and 2.2 follows.

Let  $\tilde{t}_m$  be the smallest integer divisible by  $t_m$  and by every torsion prime in  $H^*(X, Z)$ . Let  $\mathbf{P}_m = \{p \mid p \mid \tilde{t}_m\}$  and  $\tilde{\mathbf{P}}_m = \mathbf{P} - \mathbf{P}_m$ . Applying 2.2 one obtains:

**2.3. PROPOSITION.** *Let  $[Y] \in G(X)$ . Given a  $\mathbf{P}_k$  equivalence  $f: Y \rightarrow X$ , a set of integral classes  $x_{n_1}, x_{n_2}, \dots, x_{n_r}$  representing a basis for  $QH^*(X, Z)/\text{torsion}$  and a set  $\tilde{z}_{n_1}, \tilde{z}_{n_2}, \dots, \tilde{z}_{n_r}, \tilde{z}_{n_t} \in H^{n_i}(Y, Z)$ , there exists a  $\mathbf{P}_k$  equivalence  $\hat{f}: Y \rightarrow X$  with*

$$[H^*(\hat{f}, Z) - H^*(f, Z)] x_{n_j} = \tilde{t}_k \tilde{z}_{n_j} + \theta_j$$

where  $\theta_j$  is a torsion element.

*Proof.* Apply 2.2 for  $Y = \tilde{Y}$ ,  $g = 1$ ,  $z_{n_j} = \tilde{t}_k / t_k \tilde{z}_{n_j}$ . Then one has only to show that  $\tilde{f} = \hat{f}$  is a  $\mathbf{P}_k$  equivalence. (Note that if  $H^*(X, Z)$  is a torsion free  $t_k = \tilde{t}_k$  and then  $H^*(\hat{f}, Z) \otimes Z_{t_k} = H^*(f, Z) \otimes Z_{t_k}$  implies that  $\hat{f}$  is a  $\mathbf{P}_k$  equivalence). By 2.2 torsion  $\pi(\tilde{f}_i) = 0$  and hence torsion  $\pi(\hat{f}) = \text{torsion } \pi(f)$ .

Consider the following diagram:

$$\begin{array}{ccc} \pi(Y)/\text{torsion} & \xrightarrow{\pi(\hat{f})/\text{torsion}} & \pi(X)/\text{torsion} \\ \downarrow \tilde{\sigma}(Y) & \xrightarrow{\pi(f)/\text{torsion}} & \downarrow \tilde{\sigma}(X) \\ PH_*(Y)/\text{torsion} & \xrightarrow[\hat{f}_* = f^*]{f_*} & PH_*(X)/\text{torsion} \end{array}$$

( $\tilde{\sigma}$  – the Hurewicz-Serre monomorphism).

Now all groups are isomorphic and  $\det[\tilde{\sigma}(Y)] = \det[\tilde{\sigma}(X)]$

$$H^*(\hat{f}, Z)/\text{torsion} \otimes Z_{t_k} = H^*(f, Z)/\text{torsion} \otimes Z_{t_k}$$

It follows that  $H^*(\hat{f}, Z)$  and its dual  $\hat{f}_*$  are  $\mathbf{P}_k$  isomorphisms. Now if  $G$  is free abelian a homomorphism  $\alpha: G \rightarrow G$  is a  $\mathbf{P}_k$  isomorphism if and only if  $(\det \alpha, t_k) = 1$ . Now,

$$(\det \hat{f}_*) (\det \tilde{\sigma}(Y)) = (\det \tilde{\sigma}(X)) (\det \pi(\hat{f})/\text{torsion})$$

implies  $(\det \hat{f}_*) = \det(\pi(\hat{f})/\text{torsion})$  and hence  $\pi(\hat{f})/\text{torsion}$  is a  $\mathbf{P}_k$  isomorphism.

As torsion  $\pi(\tilde{f}) = \text{torsion } \pi(f)$  and  $\pi(\tilde{f})/\text{torsion}$  are  $\mathbf{P}_k$  isomorphisms so is  $\pi(\tilde{f})$  and 2.3 follows. One can construct elements in  $G(X)$  as follows:

**2.4. PROPOSITION.** *Let  $h_0: X \rightarrow K(Z, \bar{n})$  realize a basis for  $QH^*(X, Z)/\text{torsion}$  i.e.:  $x_{n_i} = H^*(h_0, Z) \iota_{n_i}$  reduces to a basis for  $QH^*(X, Z)/\text{torsion}$ . Given any matrix  $A = (a_{ij}) \in \mathcal{M}(s_1, s_2, \dots, s_l; Z)$  with  $(\det A, \bar{t}_k) = 1$  let  $f_0: K(Z, \bar{n}) \rightarrow K(Z, \bar{n})$  be given by  $H^*(f_0, Z) \iota_{n_i} = \sum_j a_{ij} \iota_{n_j}$ . Let  $Y = Y(A) = Hl_k(\tilde{Y})$  where*

$$\begin{array}{ccc} \tilde{Y} & \xrightarrow{\tilde{f}_1} & X \\ \downarrow h_1 & & \downarrow h_0 \\ K(Z, \bar{n}) & \xrightarrow{f_0} & K(Z, \bar{n}) \end{array}$$

is a fiber square induced by  $h_0, f_0$ . Then:

- (a)  $\xi(A) = [Y] \in G(X)$
- (b) The set  $\{y_{n_i} = H^{n_i}(h_1, Z) \iota_{n_i}\}$  reduces to a basis for  $QH^*(Y, Z)/\text{torsion}$ , where  $h_1: Y \rightarrow K(Z, \bar{n})$  is the composition

$$H_{l_k}(\tilde{Y}) = Y \xrightarrow{\tau'_{l_k}} \tilde{Y} \xrightarrow{h_1} K(Z, \bar{n}).$$

*Proof.* By 2.1  $h_0$  is a  $\tilde{\mathbf{P}}_k = \mathbf{P} - \mathbf{P}_k$  equivalence in  $\dim \leq k$  hence so is  $\tilde{h}_1$ . It follows that in  $\dim \leq k$   $Y \approx \tilde{Y} \approx_{\tilde{\mathbf{P}}_k} K(Z, \bar{n}) \approx_{\tilde{\mathbf{P}}_k} X$ . Consequently,  $Y \approx_{\tilde{\mathbf{P}}_k} X$  in  $\dim \leq k$  but both being of  $\dim \leq k$  it follows that  $Y \approx_{\tilde{\mathbf{P}}_k} X$ . Now,  $(\det A, \bar{t}_k) = 1$  implies that  $A = \pi(f_0)$  is a  $\mathbf{P}_k$  isomorphism and hence  $f_0$  (and consequently  $\tilde{f}_1$ ) are  $\mathbf{P}_k$  equivalences. Again in  $\dim \leq k$   $Y \approx \tilde{Y}$  and  $Y \approx_{\tilde{\mathbf{P}}_k} X$  and (a) follows. To prove (b) one notices first that in  $\dim \leq k$  one can consider

$$\begin{array}{ccc} Y & \xrightarrow{f_1} & X \\ \downarrow h_1 & & \downarrow h_0 \\ K(Z, \bar{n}) & \xrightarrow{f_0} & K(Z, \bar{n}) \end{array}$$

as a fiber square. Further, if  $[Y] \in G(X)$  then  $\text{coker } \sigma_n(Y) \approx \text{coker } \sigma_n(X)$  and  $\ker \sigma_n(Y) = \ker \sigma_n(X)$  (as all these groups are finite they involve only a finite set  $\mathbf{P}_1$  of primes and by the procedure of 1.5 there exists a  $\mathbf{P}_1$  equivalence  $Y \rightarrow X$  which will yield the desired isomorphisms). One obtains the following commutative diagrams for  $n \leq k$

$$\begin{array}{ccccccc} & & \pi_n(F) & & & & \\ & & \swarrow \pi_n(f_1) & & \searrow & & \\ \pi_n(\tilde{F}) & \xrightarrow{j_1} & \pi_n(Y) & \xrightarrow{\pi_n(f_1)} & \pi_n(X) & & \pi_{n-1}(\tilde{F}) \\ & \downarrow \pi_n(h_1) & & & \downarrow \pi_n(h_0) & & \\ & \pi_n(K(Z, \bar{n})) & \xrightarrow{\pi_n(f_0)} & \pi_n(K(Z, \bar{n})) & & & \\ & & \searrow \varphi_0 & \swarrow \varphi_1 & & & \\ & & & \pi_{n-1}(F) & & & \end{array}$$

where  $\tilde{F} = \text{fiber } f_0 = \text{fiber } f_1 \approx \prod_i K(Z_{\lambda_i}, n_i - 1)$  where the matrix  $\text{diag}(\lambda_1, \lambda_2, \dots, \lambda_r)$  is equivalent to  $A$ . As  $j_0 = 0$  and as the order of  $\pi_n(\tilde{F})$  is prime to that of  $\pi_n(F)$   $\pi_n(j_1) = 0$  and  $\pi_n(f_1)$  is a monomorphism. As  $\varphi_1$  factors through a free group  $\varphi_1(\text{tor } \pi_n(X)) = 0$  and one has a commutative diagram with exact rows

$$\begin{array}{ccccccc}
 0 \rightarrow \pi_n(Y)/\text{torsion} & \xrightarrow{f_1 \#} & \pi_n(X)/\text{torsion} & \xrightarrow{\tilde{\phi}_1} & \pi_{n-1}(\tilde{F}) \rightarrow 0 \\
 \downarrow \sigma_n(Y) & & \downarrow \sigma_n(X) & & \downarrow \sigma' & & \\
 0 \rightarrow PH_n(Y)/\text{torsion} & \xrightarrow{f_{1*}} & PH_n(X)/\text{torsion} & \xrightarrow{\varphi} & \text{coker } f_1^* & & \\
 \downarrow h_{1*} & & \downarrow h_{0*} & & \downarrow h' & & \\
 0 \rightarrow PH_n(K(Z, \bar{n}), Z)/\text{torsion} & \xrightarrow{f_{0*}} & PH_n(K(Z, \bar{n}), Z)/\text{torsion} & \xrightarrow{\tilde{\phi}_0} & \pi_{n-1}(\tilde{F}) & & \\
 \sigma_n(K) \uparrow \approx & & \approx \uparrow \sigma_n(K) & & & & \\
 0 \rightarrow \pi_n(K(Z, \bar{n})) & \xrightarrow{\pi_n(f_0)} & \pi_n(K(Z, \bar{n})) & \xrightarrow{\varphi_0} & & & 
 \end{array}$$

$\tilde{\sigma}_n(Y)$  and  $\tilde{\sigma}_n(X)$  are monomorphisms.

$\text{Coker } \tilde{\sigma}_n(Y) \approx \text{coker } \sigma_n(Y) \approx \text{coker } \sigma_n(X) \approx \text{coker } \tilde{\sigma}_n(X)$  are  $\mathbf{P}_k$  torsion groups and as  $f_1$  is a  $\mathbf{P}_k$  equivalence  $f_{1\#}$  and  $f_{1*}$  are  $\mathbf{P}_k$  isomorphisms and they induce a  $\mathbf{P}_k$  isomorphism  $\text{coker } \tilde{\sigma}_n(Y) \rightarrow \text{coker } \tilde{\sigma}_n(X)$  and hence an isomorphism. It follows that the left upper square in the last diagram is a push out diagram and  $\sigma'$  and consequently  $h'$  are isomorphisms. As  $h_{0*}$  is an isomorphism so is  $h_{1*}$  and so is its dual  $QH^*(h_1, Z)/\text{torsion}$ .

Again let  $\tilde{\alpha}: \mathcal{M}(s_1, s_2, \dots, s_l; Z) \rightarrow \mathcal{M}(s_1, s_2, \dots, s; Z_{\bar{t}_k})$  and let  $\mathcal{A} = \tilde{\alpha}^{-1} \times GL(s_1, s_2, \dots, s_l; Z_{\bar{t}_k})$ . 2.4(a) defines a correspondence

$$\xi: \mathcal{A} \rightarrow G(X). \quad \xi(I) = [X].$$

**2.5. PROPOSITION.** *Let  $[Y] \in G(X)$ . Let  $f: Y \rightarrow X$  be a  $\mathbf{P}_k$  equivalence and let  $A \in \mathcal{A}$  represent  $QH^*(f, Z)/\text{torsion}$ . Then*

$$[Y] = \xi(A).$$

*Proof.* Suppose  $A$  represents  $QH^*(f, Z)/\text{torsion}$  with respect to bases represented by  $x_{n_1}, \dots, x_{n_r}$ ,  $x_{n_i} \in H^{n_i}(X, Z)$  and  $y'_{n_1}, \dots, y'_{n_r}, y'_{n_i} \in H^{n_i}(Y, Z)$ . We shall replace  $f$  by a  $\mathbf{P}_k$  equivalence  $\hat{f}$ ,  $QH^*(\hat{f}, Z)/\text{torsion} = QH^*(f, Z)/\text{torsion}$  and  $y'_{n_1}, \dots, y'_{n_2}$  by  $y_{n_1}, \dots, y_{n_r}$  representing the same basis as  $y'_{n_1}, \dots, y'_{n_r}$  so that if  $h_0: X \rightarrow K(Z, \bar{n})$ ,  $h_1: Y \rightarrow K(Z, \bar{n})$ ,  $f_0: K(Z, \bar{n}) \rightarrow K(Z, \bar{n})$  are given by

$$\begin{aligned}
 H^*(h_0, Z) \iota_{n_i} &= x_{n_i}, \quad H^*(h_1, Z) \iota_{n_i} = y_{n_i} \\
 H^*(f_0, Z) \iota_{n_i} &= \sum_j a_{ij} \cdot \iota_{n_j} \quad A = (a_{ij}),
 \end{aligned}$$

then

$$h_0 \circ \hat{f} \sim f_0 \circ h_1.$$

In the diagram

$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \downarrow h_1 & & \downarrow h_0 \\ K(Z, \bar{n}) & \xrightarrow{f_0} & K(Z, \bar{n}) \end{array}$$

$f_0$  and  $\tilde{f}$  are  $\tilde{\mathbf{P}}_k$  equivalences and as  $QH(h_i, Z)/\text{torsion } i=0, 1$  are isomorphisms by 2.1 they are  $\mathbf{P}_k$  equivalences in  $\dim \leq k$ . Hence, by 1.6 the last diagram represents a fiber square in  $\dim \leq k$  and hence  $[Y] = \xi(A)$ . Equivalently one has to construct  $\tilde{f}$  and find  $y_{n_i}$  so that

$$H^*(\tilde{f}, Z) x_{n_i} = \sum a_{ij} y_{n_j}.$$

First note that without a loss of generality one may assume  $A$  to be diagonal for otherwise  $D = E_1 A E_2$  where  $E_i$  are invertible and  $D$ -diagonal,  $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_r)$ . Then if  $E_2^{-1}$  applied on  $y_{n_1}, \dots, y_{n_r}$  yields the basis  $\bar{y}_{n_1}, \bar{y}_{n_2}, \dots, \bar{y}_{n_r}$  and  $E_1$  applied on  $x_{n_1}, \dots, x_{n_r}$  yields a basis  $\bar{x}_{n_1}, \dots, \bar{x}_{n_r}$  then

$$H^*(f, Z)(\bar{x}_{n_1}, \dots, \bar{x}_{n_r}) = E_1 A E_2 E_2^{-1}(y_{n_1}, y_{n_2}, \dots, y_{n_r}) = D(\bar{y}_{n_1}, \dots, \bar{y}_{n_r})$$

So  $H^*(f, Z) x_{n_i} = \sum_j a_{ij} y_{n_j}$  if and only if  $H^*(f, Z) \bar{x}_{n_i} = \lambda_i \bar{y}_{n_i}$ . So assume  $A = D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_r)$ . Hence, it is given that

$$H^*(f, Z) x_{n_i} = \lambda_i y'_{n_i} + d_i + \theta_i,$$

$d_i$ -decomposable and  $\theta_i$  a torsion element. Now  $(\det A, \bar{t}_k) = 1$  implies  $(\lambda_i, \bar{t}_k) = 1$ . Let  $1 + a_i \cdot \bar{t}_k = b \cdot \lambda_i$ . Now apply 2.3 for  $\tilde{z}_{n_i} = a_i \cdot d_i$ .  $\tilde{f}$  then will satisfy

$$H^*(\tilde{f}, Z) x_{n_i} = H^*(f, Z) x_{n_i} + \bar{t}_k a_i d_i + \hat{\theta}_i = \lambda_i y'_{n_i} + (1 + \bar{t}_k a_i) d_i + \hat{\theta}_i + \theta_i.$$

As (order  $(\theta_i + \hat{\theta}_i, \lambda_i) = 1$ )  $\theta_i + \hat{\theta}_i = \lambda_i \cdot \theta'_i$  and  $H^*(f_1, Z) x_{n_i} = \lambda_i (y'_{n_i} + b_i \cdot d_i + \theta'_i)$ .  $y_{n_i} = y'_{n_i} + b_i d_i + \theta'_i$  are the desired classes.

## 2.6. COROLLARY. $\xi: A \rightarrow G(X)$ of 2.4(a) is onto.

*Proof.* If  $[Y] \in G(X)$  then by 1.5 there exists a  $\mathbf{P}_k$  equivalence  $f: Y \rightarrow X$  and by 2.5  $[Y] = \xi(A)$  for some  $A \in \mathcal{A}$ .

It is quite obvious from the definition of  $\xi$  in 2.4 that if  $B \in GL(s_1, s_2, \dots, s_l; Z)$  then  $\xi(BA) = \xi(A)$ . A simple application of 2.3 and 2.5 shows also that for any  $C \in \mathcal{M}(s_1, s_2, \dots, s_l; Z)$   $\xi(A + \bar{t}_k C) = \xi(A)$ . Hence if,

$$\bar{\alpha}: GL(s_1, s_2, \dots, s_l; Z) \rightarrow GL(s_1, s_2, \dots, s_l; Z_{\bar{t}_k})$$

then  $\xi$  factors through  $\text{coker } \bar{\alpha} = (Z_{\bar{t}_k}^*/\{\pm 1\})^l$  (see 1.8). Hence, one obtains the following

2.7. A STRUCTURE THEOREM for  $G(X)$ . *There exists a correspondence  $\xi:(Z_{t_k}^*/\{\pm 1\})^l \rightarrow G(X)$  which is onto.  $\xi$  is given by*

$$\xi(d_1, d_2, \dots, d_l) = \xi(I_{d_1} * I_{d_2} * \dots * I_{d_l}),$$

$d_i \in Z_{t_k}^*$ , where  $I_{d_i}$  is the  $s_i \times s_i$  matrix given by

$$I(d_i, s_i) = I_{d_i} = \begin{pmatrix} I & 0 \\ 0 & d_i \end{pmatrix}$$

By (2.4) it follows that  $\tilde{Y} \approx \text{fiber}(\tilde{g}: X \rightarrow K_\lambda)$  for some map  $\tilde{g}$  where  $K_\lambda = \prod_{i=1}^r K(Z_{\lambda_i}, n_i)$  and the  $\lambda_i$ 's are defined by the fact that  $A$  is equivalent to  $\text{diag}(\lambda_1, \lambda_2, \dots, \lambda_r)$ . Moreover, if  $\varrho_{\lambda_i}: H^*(\ , Z) \rightarrow H^*(\ , Z_{\lambda_i})$  is the reduction then  $H^*(\tilde{g}, Z_{\lambda_i}) \tilde{t}_{n_i} = \varrho_{\lambda_i} x_{n_i}$  where  $\tilde{t}_{n_i} \in H_{n_i}(K(Z_{\lambda_i}, n_i), Z_{\lambda_i})$  is the fundamental class and  $x_{n_1}, x_{n_2}, \dots, x_{n_r}$  are integral classes representing a basis for  $QH^*(X, Z)/\text{torsion}$ . Note that given  $[Y] \in G(X)$  by 1.5 there exists a  $P_{2k}$  equivalence  $f: Y \rightarrow X$  and by 2.5  $[Y] = \xi(A)$ ,  $(\det A, \tilde{t}_{2k}) = 1$ .

2.8. PROPOSITION. *Let  $m=2k$ . Given integers  $\lambda_1, \lambda_2, \dots, \lambda_r$  prime to  $\tilde{t}_m$  and elements  $x_{n_1}, x_{n_2}, \dots, x_{n_r} \in H^*(X, Z)$  representing a basis for  $QH^*(X, Z)/\text{torsion}$ , there exists an H-structure  $\mu_0$  for  $X$  with respect to which  $\varrho_{\lambda_i} x_{n_i}$  are primitive.*

*Proof.* As all  $\lambda_i$  are prime to all torsion primes of  $H^*(X, Z)$   $H^*(X, Z_{\lambda_i}) \approx \approx H^*(X, Z) \otimes Z_{\lambda_i}$ . Replacing each  $\lambda_i$  by the product  $\lambda = \lambda_1 \lambda_2 \cdot \dots \cdot \lambda_r$  suffices to prove the theorem for  $\lambda_1 = \lambda_2 = \dots = \lambda_r = \lambda$ .

Denote  $(\ )^* = H^*(\ , Z)/\text{torsion}$ . Let  $\tilde{x}_{n_i}$  be the image of  $x_{n_i}$  in  $H^*(X, Z)/\text{torsion}$ . Given any H-structure  $\mu$  of  $X$  one has

$$\mu^* \tilde{x}_{n_i} = \tilde{x}_{n_i} \otimes 1 + 1 \otimes \tilde{x}_{n_i} + \sum_{s=1}^a \tilde{x}'_{n_i, s} \otimes \tilde{x}''_{n_i, s}.$$

Put  $\omega_i = \bar{\mu}^* \tilde{x}_{n_i} = \mu^* \tilde{x}_{n_i} - \tilde{x}_{n_i} \otimes 1 - 1 \otimes \tilde{x}_{n_i} = \sum_{s=1}^a x'_{n_i, s} \otimes x''_{n_i, s}$  then  $\omega_i = \Lambda^* \tilde{\omega}_i$  where  $\tilde{\omega}_i \in H^*(X \wedge X, Z)/\text{torsion}$  and  $\Lambda: X \times X \rightarrow X \wedge X$  is the reduction. As  $\dim X \times X \leq m$ ,  $\dim X \wedge X \leq m$  one can apply 2.2 for

$$Y = X \times X, \quad \tilde{Y} = X \wedge X, \quad f = \mu, \quad g = \Lambda.$$

(and note that  $\overline{\text{im } \Lambda^*}$  is an ideal) and  $z_{n_i} = k \tilde{\omega}_i$  where  $k$  is an integer satisfying  $1 + \tilde{t}_m k = b \lambda$ . If  $\tilde{f} = \mu_0$  then, as each  $\tilde{f}_i$  of 2.2 factors through  $X \wedge X$ ,  $\tilde{f}_i \mid X \vee X = *$  and  $\mu_0 \mid X \vee X = \mu \mid X \vee X$ . Hence  $\mu_0$  is an H-structure. Now,

$$\begin{aligned} \mu_0^* \tilde{x}_{n_i} &= \mu_0^* \tilde{x}_{n_i} + \tilde{t}_m k \omega_i = \tilde{x}_{n_i} \otimes 1 + 1 \otimes \tilde{x}_{n_i} + \omega_i + \tilde{t}_m k \omega_i \\ &= \tilde{x}_{n_i} \otimes 1 + 1 \otimes \tilde{x}_{n_i} + \lambda b \omega_i. \end{aligned}$$

As  $\varrho_\lambda x_{n_i} = \tilde{\varrho}_\lambda \tilde{x}_{n_i}$  ( $\tilde{\varrho}_\lambda: H^*(X, Z)/\text{torsion} \rightarrow H^*(X, Z_\lambda)$ )

$$\begin{aligned} H^*(\mu_0, Z_\lambda) \varrho_\lambda x_{n_i} &= H^*(\mu_0, Z_\lambda) \tilde{\varrho}_\lambda \tilde{x}_{n_i} \\ &= \tilde{\varrho}_\lambda \tilde{x}_{n_i} \otimes 1 + 1 \otimes \tilde{\varrho}_\lambda \tilde{x}_{n_i} = \varrho_\lambda x_{n_i} \otimes 1 + 1 \otimes \varrho_\lambda x_{n_i} \end{aligned}$$

and  $\varrho_\lambda x_{n_i}$  are primitive.

2.9. COROLLARY. *If  $[Y] \in G(X)$ . Then  $Y$  is an H-space and there exists a  $P_{2k}$  equivalence  $f_1: Y \rightarrow X$  which is an H-map with respect to some H-structure of  $X$ .*

*Proof.* By the remarks preceding 2.8 one may replace  $k$  by  $m=2k$  in 2.7 and then  $\tilde{f}_1: \tilde{Y} \rightarrow X$  is the fiber of a map  $\tilde{g}: X \rightarrow K_\lambda$   $K_\lambda = \prod_i K(\lambda_i, n_i)$  ( $\lambda_i, \tilde{\iota}_{2k} = 1$ ). By 2.8 choosing the H-structure of  $X$  properly  $\tilde{g}$  is an H-map and so  $\tilde{Y}$  and  $\tilde{f}_1$  become an H-space and an H-map. As  $Hl_{2k}(\tilde{Y}) = Hl_k(\tilde{Y}) = Y$   $Y$  and  $\tau': Y \rightarrow \tilde{Y}$  admit H-structure and so does  $f_1 = \tilde{f}_1 \circ \tau'$ .

2.10. COROLLARY. (See [6] and [7]). *Let  $X$  be a finite CW complex. If for every prime  $p$  there exist an H-space  $X(p)$  and a  $p$ -equivalence  $f_q: X \rightarrow X(p)$  and if all  $H^*(X(p), Q)$ ,  $p \in P$ , are isomorphic as Hopf algebras then  $X$  admits an H-structure.*

*Proof.* Let  $H^*(X, Q) \approx H^*(X(p), Q) \approx \Lambda(n_1, x_{n_2}, \dots, x_{n_r})$ . As  $X \approx (S^{\bar{n}} = S^{n_1} \times S^{n_2} \times \dots \times S^{n_r})$  for almost all primes one may assume that the number of different spaces  $X(p)$  and maps  $f_q$  is finite. Now, for every  $p, q \in P$   $H^*(X(p), Q) \approx H^*(X(q), Q)$ . Hence  $K(Z, \bar{n}) = \prod_{i=1}^r K(Z, n_i)$  admits an H-structure and there exist  $Q$  equivalences  $h_p: X(p) \rightarrow K(Z, \bar{n})$  which are H-maps. Moreover, using the procedure of [8] one may replace  $X(p)$  by  $X'(p)$  so that  $h_p$  decomposes into two H-maps  $X(p) \xrightarrow{h'_p} X'(p) \xrightarrow{h''_p} K(Z, \bar{n})$  where  $h'_p$  is a  $p$ -equivalence and  $h''_p$  is a  $P - \{p\}$  equivalence. The pullback  $X' = \prod_{h''_p} X'(p)$  is then an H-map, and  $X' \approx_p X'(p) \approx_p X$ . Hence  $[Y] \in G(X')$  and apply 2.9.

Let  $A, \hat{A} \in \mathcal{A} \subset M(s_1, s_2, \dots, s_l; Z)$ . Then

$$\bar{\alpha}A \equiv \bar{\alpha}(I(d_1, s_1) * I(d_2, s_2) * \dots * I(d_l, s_l)) \bmod \text{im } \bar{\alpha}.$$

$$\bar{\alpha}\hat{A} \equiv \bar{\alpha}(I(\hat{d}_1, s_1) * I(\hat{d}_2, s_2) * \dots * I(\hat{d}_l, s_l)) \bmod \text{im } \bar{\alpha}$$

(If  $A = A_1 * A_2 * \dots * A_l$  then  $d_i = \det A_i \in Z_{t_k}^*$  and similarly for  $\hat{A}_i$  and  $\hat{d}_i$ ). It can be easily seen that

$$\xi(A) \times \xi(\hat{A}) = \xi(A *_g \hat{A}) \in G(X \times X)$$

As  $I(d_i, s_i) * I(\hat{d}_i, s_i) \equiv I * I(d_i \cdot \hat{d}_i, s_i) = I(d_i \cdot \hat{d}_i, 2s_i) \bmod \text{im } \bar{\alpha}$  one has  $\tilde{\alpha}(A *_g A \hat{A}) \equiv \tilde{\alpha}(I *_g A \hat{A}) \bmod \text{im } \bar{\alpha}$  and therefore

$$\xi(A) \times \xi(\hat{A}) = \xi(A *_g \hat{A}) = \xi(I *_g A \hat{A}) = \xi(I) \times \xi(A \cdot \hat{A}).$$

We thus proved

2.11. A PRODUCT THEOREM. *Let  $A, \hat{A} \in \mathcal{A}$ . Then  $\xi(A) \times \xi(\hat{A}) = X \times \xi(A \cdot \hat{A})$  and consequently if  $A_i, B_i \in \mathcal{A}$  and if  $A_1 \cdot A_2 \cdot \dots \cdot A_n$  and  $B_1 \cdot B_2 \cdot \dots \cdot B_m$  have the same image in  $GL(s_1, s_2, \dots, s_l; \mathbb{Z}_{t_k})/\text{im } \bar{\alpha}$  then*

$$\xi(A_1) \times \xi(A_2) \times \dots \times \xi(A_m) = \xi(B_1) \times \xi(B_2) \times \dots \times \xi(B_m).$$

If  $\varphi$  is the Euler function then as  $I(d_i, s_i)^{\varphi(t_k)/2} \equiv I \pmod{\text{im } \bar{\alpha}}$  one obtains (see [7], p. 82):

$$2.12. \text{ COROLLARY. For every } [Y] \in G(X) \text{ } Y^{\varphi(t_k)/2} \approx X^{\varphi(t_k)/2}.$$

Similarly one obtains

2.13. COROLLARY. *Let  $Y$  be a finite CW H-space if  $QH^n(Y, Z)/\text{torsion} \neq 0$  whenever  $QH^n(X, Z)/\text{torsion} \neq 0$  then  $G(X \times Y) = \{X\} \times G(Y)$ .*

### 3. Genus and non-Cancellation

3.1. DEFINITION. (See [5]). Let  $K$  be a finite CW complex,  $\mathbf{P}_1$  – a set of primes.  $K$  is said to be  $\mathbf{P}_1$  universal if for every prime  $q \notin \mathbf{P}_1$  there exists a  $\mathbf{P}_1$  equivalence  $f_q: K \rightarrow K$  so that  $\bar{H}^*(f_q, \mathbb{Z}_q) = 0$ .

A space  $Y$  is said to be an  $H_0$  space if  $H^*(Y, Q)$  is a free (commutative and associative) algebra. If  $H^*(Y, Q) = \Lambda(x_{n_1}, x_{n_2}, \dots, x_{n_r})$  then  $Y$  is said to be of type  $n_1, n_2, \dots, n_r$ . The following is a generalization of [7] Lemma 1.5:

3.2. PROPOSITION. *Let  $\mathbf{P} = \mathbf{P}_1 \cup \mathbf{P}_2$ . Let  $X, Y_1, Y_2$  be finite CW complexes  $f: X \rightarrow Y_1 \times Y_2$ . If  $X$  is an  $H_0$ -space,  $Y_i - \mathbf{P}_i$  universal  $i = 1, 2$  and if  $p_i \circ f$  is a  $\mathbf{P}_i$  equivalence where  $p_i: Y_1 \times Y_2 \rightarrow Y_i$ ,  $i = 1, 2$ , is the projection then  $f$  admits a homotopy left inverse.*

*Proof.* We may assume that  $\mathbf{P}_1 \cap \mathbf{P}_2 = \emptyset$ . Let  $h_0: X \rightarrow X_0 = K(Z, \bar{n})$  be a rational equivalence. Put  $m = \max(n_r, \dim Y_1 \times Y_2)$ . Factor  $\hat{h}_0: Ht_m(X) \rightarrow X_0$  by maps  $h_{i, i-1}: X_i \rightarrow X_{i-1}$ ,  $0 < i \leq s$ ,  $X_s = Ht_m(X)$ ,  $h_{i, i-1}$  being the fiber of a map  $k_i: X_{i-1} \rightarrow K(Z_{p_i}, m_i)$ ,  $p_i \in \mathbf{P}$ . Let  $\alpha_i: Y_1 \times Y_2 \rightarrow Y_1 \times Y_2$  be given as follows: If  $p_i \notin \mathbf{P}_1$  let  $g'_i: Y_1 \rightarrow Y_1$  be a  $\mathbf{P}_1$  equivalence with  $\bar{H}^*(g'_i, \mathbb{Z}_{p_i}) = 0$ . If  $p_i \notin \mathbf{P}_2$  let  $g''_i: Y_2 \rightarrow Y_2$  be defined similarly. Let

$$\alpha_i = \begin{cases} g'_i \times 1 & \text{if } p_i \notin \mathbf{P}_1 \\ 1 \times g''_i & \text{if } p_i \notin \mathbf{P}_2 \end{cases}$$

suffices to find a left inverse for  $\hat{f} = \alpha_1 \circ \alpha_2 \circ \dots \circ \alpha_s \circ f$  and one has  $p_i \circ \hat{f}$  is a  $\mathbf{P}_i$  equivalence. Let  $y_{n_i} = H^*(h_0, Z)$ ,  $\iota_{n_1}, \dots, \iota_{n_r} \in H^*(K(Z, \bar{n}), Z)$  the “fundamental vector”. As  $p_1 \circ f$  is a  $\mathbf{P}_1$  equivalence  $\lambda_1 y_{n_i} \in \text{im } H^*(p_1 \circ \hat{f}, Z)$ ,  $\lambda_1$  prime to  $\mathbf{P}_1$ . Similarly

$\lambda_2 y_{n_i} \in \text{im } H^*(p_2 \circ f, Z)$ . As  $(\lambda_1, \lambda_2) = 1$   $\lambda_1 y_{n_i}, \lambda_2 y_{n_i} \in \text{im } H^*(\hat{f}, Z)$  implies  $y_{n_i} \in \text{im } H^*(\hat{f}, Z)$ . Hence,  $h_0$  factors through  $Y_1 \times Y_2$ ,  $h_0 = r_0 \circ \hat{f}$ ,  $r_0: Y_1 \times Y_2 \rightarrow K(Z, \bar{n}) = X_0$ . Suppose one obtains inductively the following commutative diagram:

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y_1 \times Y_2 & \xrightarrow{\alpha_s} & \cdots Y_1 \times Y_2 & \xrightarrow{\alpha_n} & Y_1 \times Y_2 \\ \tau_m \downarrow & & & & & \downarrow r_{n-1} & \\ Ht_m(X) = X_s & \xrightarrow{h_{s, s-1}} & X_{s-1} & \longrightarrow & \cdots X_n & \xrightarrow{h_{n, n-1}} & X_{n-1} \xrightarrow{k_n} K(Z_{p_n}, m_n) \end{array}$$

Without loss of generality suppose  $p_n \notin \mathbf{P}_1$ . Then  $H^*(\alpha_n, Z_{p_n}) = H^*(g'_n, Z_{p_n}) \otimes 1 = \varepsilon \otimes 1$  (where  $\varepsilon: H^* \rightarrow Z_p \rightarrow H^*$  is induced by augmentation). Hence, one obtains

$$\begin{array}{ccc} Y_1 \times Y_2 & \xrightarrow{p_2} & Y_2 \\ \downarrow \alpha_n & & \downarrow \tilde{r} \\ Y_1 \times Y_2 & \xrightarrow{r_{n-1}} & X_{n-1} \xrightarrow{k_n} K(Z_{p_n}, m_n) \end{array}$$

$p_2 \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f$  is a  $\mathbf{P}_2$ -equivalence, and as  $k_n \circ h_{n, n-1} \sim *$ ,  $* \sim k_n \circ r_{n-1} \circ \alpha_n \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f \sim \tilde{r} \circ p_2 \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f$ . It follows that  $\tilde{r} \sim *$ ,  $k_n \circ r_{n-1} \circ \alpha_n \sim *$  and  $r'_n: Y_1 \times Y_2 \rightarrow X_n$  exists so that  $h_{n, n-1} \circ r'_n \sim r_{n-1} \circ \alpha_n$ . Comparing  $\varphi_1 = r'_n \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f$  and  $\varphi_2 = h_{n+1, n} \circ h_{n+2, n+1} \circ \cdots \circ h_{s, s-1} \circ \tau_m$  there exists  $\omega: X \rightarrow K(Z_{p_n}, m_n - 1)$  so that  $\omega * \varphi_1 \sim \varphi_2$  (where  $*$  denotes the action of  $[X, K(Z_{p_n}, m_n - 1)]$  on  $[X, X_n]$  induced by the principal fibration  $K(Z_{p_n}, m_n - 1) \rightarrow X_n \rightarrow X_{n-1}$ ). But  $\alpha_{n+1} \circ \alpha_{n+2} \circ \cdots \circ \alpha_s \circ f$  is a  $\mathbf{P}_2$  equimorphism, hence,  $\omega$  factors as  $\tilde{\omega} \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f$ ,  $\tilde{\omega}: Y_1 \times Y_2 \rightarrow K(Z_{p_n}, m_n - 1)$ .

Replacing  $r'_n$  by  $\tilde{\omega} * r'_n = r_n$  one obtains  $r_n: Y_1 \times Y_2 \rightarrow X_n$  and

$$\begin{aligned} r_n \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f &= (\tilde{\omega} * r'_n) \circ \alpha_{n+1} \circ \cdots \circ \alpha_0 \sim \\ &\sim (\tilde{\omega} \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f) * (r'_n \circ \alpha_{n+1} \circ \cdots \circ \alpha_s \circ f) \sim \\ &\sim \omega * \varphi_1 \sim \varphi_2 = h_{n+1, n} \circ h_{n+2, n+1} \circ \cdots \circ h_{s, s-1} \circ \tau_m. \end{aligned}$$

The final step of lifting  $r_s: Y_1 \times Y_2 \rightarrow X_s = Ht_m(X)$  to  $r: Y_1 \times Y_2 \rightarrow X$  with  $r \circ f \sim 1$  is automatic as  $\dim X \leq m$ ,  $\dim Y_1 \times Y_2 \leq m$  and hence  $[X, \tau_m]: [X, X] \xrightarrow{\sim} [X, X_s]$ ,  $[Y_1 \times Y_2, \tau_m]: [Y_1 \times Y_2, X] \xrightarrow{\sim} [Y_1 \times Y_2, X_s]$  and  $\tau_m = [X, \tau_m][1]$ .

### 3.3. LEMMA. Let

$$\begin{array}{ccc} X & \xrightarrow{h_2} & Y_1 \\ h_1 \downarrow & & \downarrow f_1 \\ Y_2 & \xrightarrow{f_2} & X_0 \end{array}$$

be a fiber square. If  $X_0$  is an H-space then  $X \xrightarrow{(h_1 \times h_2) \circ \Delta} Y_1 \times Y_2 \xrightarrow{[f_2][f_1]^{-1}} X_0$  is a (quasi) fibration.

*Proof.* Denote  $\mu_{X_0}(x, x')$  by  $x \cdot x'$  and if  $c: X \rightarrow X$  is the homotopy inverse put

$c(x)=x^{-1}$ . Now,

$$X = \{y_1, y_2, \varphi \in Y_1 \times Y_2 \times PX_0 \mid f_1(y_1) = \varphi(0), \varphi(1) = f_2(y_2)\}$$

and

$$F = \{y_1, y_2, \varphi \in Y_1 \times Y_2 \times \mathcal{L}X_0 \mid f_2(y_2) \cdot f_1(y_1)^{-1} = \varphi(1)\}$$

is the fiber of  $[f][f_1]^{-1}$ . Let  $\alpha: X \rightarrow F$  be given by  $\alpha(y_1, y_2, \varphi) = y_1, y_2, a(f_1(y_1)) + \varphi \cdot f_1(y_1)^{-1}$  where  $a(z)$  is the homotopy connecting the base point with  $z \cdot z^{-1}$ . The obvious maps  $X \rightarrow Y_1 \times Y_2$  and  $F \rightarrow Y_1 \times Y_2$  are (Hurewicz) fibrations with the same fiber  $\Omega X_0$  and  $\alpha|_{\Omega X_0} = 1$ . Hence,  $\alpha$  is a homotopy equivalence.

Combining 1.6, 3.2 and 3.3 one gets

### 3.4. THEOREM. *Given a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{h_2} & Y_1 \\ \downarrow h_1 & & \downarrow f_1 \\ Y_2 & \longrightarrow & X_0 \end{array}$$

Suppose  $f_i, h_i$  are  $\mathbf{P}_i$  equivalences in  $\dim \leq n$ ,  $i=1, 2$ ,  $\mathbf{P}_1 \cup \mathbf{P}_2 = \mathbf{P}$ . If  $Y_1$  is  $\mathbf{P}_i$  universal  $i=1, 2$ ,  $X$  is an  $H_0$  space and  $X_0$  is an  $H$ -space then

$$Y_1 \times Y_2 \approx X \times X_0 \quad \text{in } \dim \leq n.$$

*Proof.* By 1.6 3.3.1 is a fiber square. By 3.3 one has a fibration

$$X \xrightarrow{(h_1 \times h_2)\Delta} Y_1 \times Y_2 \rightarrow X_0$$

where  $h_i$  are  $\mathbf{P}_i$  equivalences and one can apply 3.2 to find a left inverse for  $(h_1 \times h_2)\Delta$  and 3.3 follows.

Let  $X$  be again an  $H$ -space of type  $(n_1, n_2, \dots, n_r)$  and of dimension  $\leq k$ . The set of primes  $\mathbf{P}_{2k}$  contains all primes involved in  $H_*(X, Z)$ ,  $\pi_n(X)$   $n \leq 2k$  and  $\text{coker}(\pi_*(X)/\text{torsion} \rightarrow PH_*(X, Z)/\text{torsion})$ . One can easily see that every map  $S^n \rightarrow X$  yielding an isomorphism of  $\pi_*/\text{torsion}$  is a  $\tilde{\mathbf{P}}_k = \mathbf{P} - \mathbf{P}_k$  equivalence. (Hence  $\mathbf{P}_k$  contains only regular primes). Let  $[Y] \in G(X)$  and let  $f: Y \rightarrow X$  be a  $\mathbf{P}_{2k}$  equivalence which is an  $H$ -map (see 2.9). Diagonalizing  $\pi(f)/\text{torsion}$  one obtains bases  $a_1, \dots, a_r \in \pi_*(Y)/\text{torsion}$  and  $b_1, \dots, b_r \in \pi_*(X)/\text{torsion}$  so that  $(\pi(f)/\text{torsion}) a_i = \lambda_i b_i$ ,  $\lambda_i$  prime to  $\mathbf{P}_{2k}$ . If  $\tilde{a}_i \in \pi_{n_i}(Y)$ ,  $\tilde{b}_i \in \pi_{n_i}(X)$  represent  $a_i$  and  $b_i$  respectively then  $\pi(f) \tilde{a}_i = \lambda_i \tilde{b}_i + \theta_i$ ,  $\theta_i$  a torsion element. But  $\lambda_i$  is prime to the torsion primes in  $\pi_n(X)$   $n \leq 2k$  hence to the order of  $\theta_i$  and  $\theta_i = \lambda_i \tilde{\theta}_i$ . Replacing  $\tilde{b}_i$  by  $\tilde{b}_i + \tilde{\theta}_i$  if necessary one may assume  $\pi(f) \tilde{a}_i =$

$= \lambda_i \tilde{b}_i$ . Hence one obtains the following commutative diagram:

$$\begin{array}{ccccc}
 S^{n_1} \vee S^{n_2} \dots \vee S^{n_k} & \xrightarrow{\tilde{a}_1 \vee \tilde{a}_2 \dots \vee \tilde{a}_2} & Y & & \\
 \downarrow h_{\lambda_1} \vee h_{\lambda_2} \dots \vee h_{\lambda_2} & h = \prod_i h_{\lambda_i} & \downarrow & & \\
 S^{\tilde{n}} & \xrightarrow{\tilde{a}_1 \cdot \tilde{a}_2 \dots \cdot \tilde{a}_r = \tilde{a}} & & & \\
 \downarrow & & \downarrow f & & \\
 S^{\tilde{n}} & \xrightarrow{\tilde{b}_1 \cdot \tilde{b}_2 \dots \cdot \tilde{b}_2 = \tilde{b}} & & & \\
 \downarrow & & \downarrow & & \\
 S^{n_1} \vee S^{n_2} \vee \dots \vee S^{n_2} & \xrightarrow{\tilde{b}_1 \vee \tilde{b}_2 \dots \vee \tilde{b}_2} & X & & 
 \end{array}$$

As  $f$  is an H-map one can add  $h = \prod_i h_{\lambda_i}$  to the above diagram to obtain a commutative diagram:

$$\begin{array}{ccc}
 S^{\tilde{n}} & \xrightarrow{\tilde{a}} & Y \\
 \downarrow h & \downarrow f & \\
 S^{\tilde{n}} & \xrightarrow{\tilde{b}} & X
 \end{array}$$

Now  $f$  and  $h$  are  $\mathbf{P}_k$  equivalences while  $\tilde{a}$  and  $\tilde{b}$  are  $\mathbf{P}_k$ -equivalences and one can apply 3.4 to get the following:

**3.5. A NON CANCELATION THEOREM.** (Compare with [1], [2] and [7], p. 83). *If  $[Y] \in G(X)$  then  $Y \times S^{\tilde{n}} \approx X \times S^{\tilde{n}}$ .*

## REFERENCES

- [1] FREYD, P., *Stable homotopy*. Proceeding of the conference on categorical algebra (La Jolla, 1965). Springer 1966.
- [2] —, *Stable homotopy II*. AMS, Proceedings of symposia in pure mathematics, Vol. XVII, 1970, pp. 161–183.
- [3] HILTON, P. J., *On the Grothendieck group of compact polyhedra*, Fund. Math. 61 (1967), pp. 199–214.
- [4] HILTON, P. J. and ROITBERG, J., *On principal  $S^3$  bundles over spheres*, Ann. of Math. 90 (1969), pp. 91–107.
- [5] MIMURA, M. and TODA, H., *On  $p$ -equivalences and  $p$ -universal spaces*, Comment. Math. Helv. 46 (1971), pp. 87–97.
- [6] MISLIN, G., *H-spaces mod  $p$ . H-spaces conference*, Neuchâtel 1970, Springer lecture notes 196.
- [7] —, *The Genus of an H-space*, Symp. on algebraic topology, Seattle 1971, Springer lecture notes 249, pp. 75–83.
- [8] ZABRODSKY, A., *Homotopy associativity and finite CW complexes*, Topology 9 (1970), pp. 121–128.

*The Hebrew University of Jerusalem, Israel  
and Forschungsinstitut für Mathematik ETH Zurich, Switzerland*

Received July 16, 1973