

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 50 (1975)

Artikel: Spines of Topological Manifolds.
Autor: Pedersen, Erik Kjaer
DOI: <https://doi.org/10.5169/seals-38791>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 04.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Spines of Topological Manifolds

ERIK KJAER PEDERSEN

In this paper we prove that a closed 2-connected topological manifold has a PL-spine, i.e. there is a locally tamely embedded complex such that a regular neighborhood of this complex is the manifold with a disc deleted (dimension is assumed to be at least 6). This “spine method” together with the relative edition of regular neighborhoods of complexes in topological manifolds [5] makes it easy to use general position arguments in topological manifolds. This will be used in a forthcoming paper to extend various embedding theorems to the topological category.

The methods we use are *PL*-approximation theorems due to Cernavskii, Connally, Miller, Rushing... as quoted in [5] theorem 2 and blocktransversality for *PL* complexes and *PL* submanifolds as was first considered by C. Morlet [4] and later extended by D. Stone [6].

DEFINITION 1. A spine of a topological manifold M with $\partial M \neq \emptyset$ is a locally tamely embedded complex $K \subset M$ so that K is a strong deformation retract of M and $K \subset M$ is a simple homotopy equivalence. In case $\partial M = \emptyset$ by the spine of M we mean a spine of M with a disc deleted.

THEOREM 2. Let $(M, \partial_- M, \partial_+ M)$ be a triad of topological manifolds $\dim(M) = m$, and assume $m \geq 6$ and

$$\Pi_j(M, \partial_+ M) = 0 \quad \text{for } j < m - r, \quad r \leq m - 3.$$

Further assume there is a *PL*-complex P locally tamely embedded in the interior of $\partial_+ M$, $\dim(P) = p$ and $m - p \geq 4$. Then there is a complex K of dimension $\max(p + 1, r, 2)$, locally tamely embedded in M such that

$$K \cap \partial_+ M = P$$

$$K \cap \partial_- M = K'$$

K' a subcomplex of K , K' has a neighborhood of the form $K' \times I$ in K and

$$\partial_- M \cup K \subset M$$

is a strong deformation retract and a simple homotopy equivalence.

Theorem 2 has an immediate corollary:

COROLLARY 3. *Let M be a closed topological manifold, $\dim(M) \geq 6$ and $\pi_j(M) = 0$ for $j \leq r$, $r > 2$. Then M has a spine of dimension $m - r$.*

Proof. Let $\bar{M} = M - (\text{interior of a disc})$. Put $\partial_- M = \emptyset$, $\partial_+ M = S^{m-1}$, $P = \emptyset$, and apply Theorem 2.

Proof of Theorem 2. First let us consider the case where $P = \emptyset$. Put $k = \max(r, 2)$. According to Kirby and Siebenmann [3] M has a handlebodydecomposition relative to $\partial_- M$ with no handles of dimension greater than k : Kirby and Siebenmann prove that $(M, \partial_- M)$ has a handlebodydecomposition, and one can then cancel handles to get a minimal handledecomposition. Because of problems with torsion one needs at least 1- and 2-handles.

We filter M by the handlefiltration

$$\partial_- M \times I = M_0 \subset M_1 \subset \cdots \subset M_s = M$$

where M_{i+1} is obtained from M_i by adjoining a single handle, no handles of dimension greater than $m - 3$. The proof will be by downwards induction on the statement:

There is a locally tamely embedded complex

$$K_i \subset \overline{M - M_i} \quad \dim(K_i) \leq k$$

such that

$$K'_i = K_i \cap \partial_+ M_i$$

is contained in the interior of $\partial_+ M_i$, K'_i has a neighborhood in K_i of the form $K'_i \times [0, 1]$ and $M_i \cup K_i$ is a simple strong deformation retract of M .

It is easy to start the induction, we let K_{s-1} be the core of the last handle. Then clearly $M_{s-1} \cup K_{s-1}$ is a simple strong deformation retract of $M = M_s$, so assume the statement for $i + 1$. Now

$$M_{i+1} = M_i \cup_{S^{j-1} \times D^{m-j}} D^j \times D^{m-j}$$

for some $j \leq k$. Let

$$\bar{E} = D^j \times D^{m-j} \cap \partial M_{i+1} = D^j \times S^{m-j-1}$$

take an outside collar $\partial \bar{E} \times [0, 2]$ of $\partial \bar{E}$ in $\partial_+ M_{i+1}$ and let

$$E_1 = \bar{E} \cup \partial \bar{E} \times [0, 1], \quad E_2 = \bar{E} \cup \partial \bar{E} \times [0, 2].$$

\bar{E} has a *PL* structure being a codimension 0 submanifold of the boundary of $D^j \times D^{m-j}$, and we can extend this *PL* structure to E_1 and E_2 using the collar. K'_{i+1} is of codi-

mension more than 3 in $\partial_+ M_{i+1}$, so by [1], see e.g. [5] Theorem 2, since $\dim \partial_+ M_{i+1} \geq 5$, there is an ambient ε -isotopy of E_2 fixing ∂E_2 that moves K'_{i+1} to be *PL* embedded in E_2 except in a neighborhood of ∂E_2 which can be assumed small. So we may assume, since this can be taken to be the restriction of an ambient isotopy of M , that $K'_{i+1} \cap E_1 \subset E_1$ is *PL*. Using [4] we can isotop K'_{i+1} further by a small ambient isotopy so that K'_{i+1} intersects $\partial \bar{E} = S^{j-1} \times S^{m-j-1}$ blocktransversally. Assume this done, and denote

$$Z = K'_{i+1} \cap \partial \bar{E}.$$

Since the normal blockbundle of $\partial \bar{E}$ in E_1 is a trivial one dimensional bundle we obtain that $\partial \bar{E}$ has a neighborhood in E of the form $\partial \bar{E} \times (-1, 1)$ and

$$\partial \bar{E} \times (-1, 1) \cap K'_{i+1} = Z \times (-1, 1)$$

since it is the restriction of the trivial blockbundle to Z , by blocktransversality. Z is a *PL* subcomplex of $S^{j-1} \times S^{m-j-1}$, which is the boundary of $S^{j-1} \times D^{m-j}$, of dimension $m-1-r$, so of codimension at least 3. By [2] Theorem 5.2 there is a subcomplex Z' of $S^{j-1} \times D^{m-j}$ of dimension $\min(\dim(Z)+1, j)$ so that

$$Z' \cap S^{j-1} \times S^{m-j-1} = Z$$

and $S^{j-1} \times D^{m-j}$ simplicially collapses to Z' ($S^{j-1} \times D^{m-j}$ is the mapping cylinder of the projection $S^{j-1} \times S^{m-j-1} \rightarrow S^{j-1}$, so take Z' to be the mapping cylinder of the restriction to Z). Using [2] lemma 2.20 this implies that

$$S^{j-1} \times D^{m-j} \times I$$

simplicially collapses to

$$S^{j-1} \times D^{m-j} \times 0 \cup Z' \times I \cup S^{j-1} \times D^{m-j} \times 1$$

so taking $S^{j-1} \times D^{m-j} \times I$ to be a collar of $S^{j-1} \times D^{m-j}$ in $D^j \times D^{m-j}$ we see that if we define D to be

$$D = \overline{D^j \times D^{m-j} - S^{j-1} \times D^{m-j} \times I}$$

there is a simple strong deformation retract of $M_{i+1} \cup K_{i+1}$ to $M_i \cup Z' \times I \cup K_{i+1} \cup D$. However D is a disc, and $Z' \times 1 \cup K_{i+1} \cap \partial D$ is of codimension bigger than 3 in ∂D , so we may as before assume it is *PL*-embedded and D now simplicially collapses to the cone of $Z' \times 1 \cup K_{i+1} \cap \partial D$ thus finishing the induction step. It is clear by construction that K'_i has a product neighborhood in K_i .

In case $P \neq \emptyset$ the proof is the same except we have to go through the motions of the induction step in the initial step of the induction too.

REFERENCES

- [1] CONNELLY, R., *Unknotting close embeddings of polyhedra in codimensions other than two*, to be published, see also Proceedings of The University of Georgia Topology of Manifolds Institute 1969, 384–389.
- [2] HUDSON, J. F. P., *Piecewise Linear Topology* (Math. Lecture Note Ser.) Benjamin, New York, 1969.
- [3] KIRBY, R. C. and SIEBENMAN, L. C., *Some theorems on Topological Manifolds*. 'Manifolds', editor N. Kuiper, Springer Lecture Notes, 1971.
- [4] MORLET, C., *Topologie des variétés semi-linéaires*, Ann. Scient. Ec. Norm., 4^{ème} serie, t. 1, 1968, 313–394.
- [5] PEDERSEN, E. K., *Topological Concordances* (to appear).
- [6] STONE, D. A., *Stratified Polyhedra*, Lecture Notes in Mathematics vol. 252, Springer Verlag 1972.

*Dept. of Mathematics
University of Chicago
5734 University Avenue
Chicago, Ill. 60637
U.S.A.*

present address:
*Matematisk Institut
Odense Universitet
Niels Bohrs Allé
5000 Odense
Danmark*

Received March 1, 1974