

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 50 (1975)

Artikel: Regular Rational Homotopy Types.
Autor: Body, Richard
DOI: <https://doi.org/10.5169/seals-38795>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 04.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Regular Rational Homotopy Types

by RICHARD BODY

§1. Regular Rational Homotopy Types

In general, spaces sharing the same integral cohomology ring need not be homotopy-equivalent. However we shall see that cohomology rings which satisfy an algebraic regularity condition may be shared by only a finite number of distinct homotopy types, [1], [2], [6].

The class of rings we shall consider are those associative, graded-commutative rings which, when tensored with the rational field, have a regular set of relations (see Definition 2.1).

THEOREM (3.1). *Let A be an associative, graded-commutative ring such that $A \otimes Q$ has a regular set of relations. Then there are only a finite number of homotopy types of finite, simply-connected polyhedra X for which $H^*(X; Z)$ is isomorphic to A , as graded rings.*

Within the category of finite, simply-connected CW complexes, all H -spaces, Riemannian symmetric spaces and homogeneous spaces which are compact Lie groups modulo a closed subgroup of maximal rank, all have integral cohomology appropriate to the above theorem.

Finally, we may view the above theorem as a generalization of the results of [1], because the set of relations $\{x_1^{n_1}, x_2^{n_2}, \dots, x_m^{n_m}\}$ is regular.

§2. Regular Sequences of Relations, and the Construction of a Model Space

All rings under consideration will be associative and graded-commutative (i.e. $a \cdot b = (-1)^{(\dim a)(\dim b)} b \cdot a$).

A free algebra over the rational field Q will then be the tensor product of

- 1) polynomial algebras on each of the even-dimensional generators, and
- 2) exterior algebras on each of the odd-dimensional generators.

Denote such a free algebra as $F = F(\xi_1, \xi_2, \dots, \xi_m)$ where (ξ_1, \dots, ξ_m) denotes a choice of generators.

DEFINITION 2.1. A sequence of elements $(\lambda_1, \dots, \lambda_k)$ in F , a rational free algebra

is said to be *regular* if F is a free-module over the free algebra $F(\lambda_1, \dots, \lambda_k)$ (under the action of the inclusion of the generators λ_i in F).

From [5], under the condition that λ_i is of even degree, regularity is equivalent to the following condition on the sequence $(\lambda_1, \lambda_2, \dots, \lambda_k)$:

$$\lambda_i \text{ mod } (\lambda_1, \dots, \lambda_{i-1}) \text{ in } F/(\lambda_1, \dots, \lambda_{i-1}) \text{ is not a zero divisor.}$$

If B , a rational algebra, has a regular set of relations, i.e. $B \simeq F(\xi_1, \xi_2, \dots, \xi_m)/(\lambda_1, \dots, \lambda_k)$, we may construct a CW complex M_B which has rational cohomology isomorphic to B , by the following procedure.

Let K_X and K_A be generalized Eilenberg-MacLane spaces.

$$K_X = \prod_{i=1}^m K(Z, \dim \xi_i); \quad K_A = \prod_{i=1}^k K(Z, \dim \lambda_i).$$

The integral cohomology of K_X , modulo torsion, is generated by fundamental classes x_1, x_2, \dots, x_m ; that of K_A by generators l_1, l_2, \dots, l_k , where $\dim \xi_i = \dim x_i$ and $\dim \lambda_i = \dim l_i$.

Without loss of generality we may consider $\lambda_i \in F(\xi_1, \dots, \xi_m)$ to be a polynomial in ξ_1, \dots, ξ_m with coefficients which are integers, collectively having no common divisor.

Now define a map $\beta: K_X \rightarrow K_A$ (i.e. a sequence of integral cohomology classes of K_X) by requiring that $\beta^*(l_i) = \lambda_i(x_1, \dots, x_m)$.

LEMMA 2.2. *The Fibre of β , denoted M_B , has rational cohomology $H^*(M_B; Q) \simeq B$.*

Proof. By induction on k , the number of relations $\lambda_1, \dots, \lambda_k$. Suppose

$$B_k = \frac{F(x_1, \dots, x_m)}{(\lambda_1, \dots, \lambda_k)} \simeq H^*(\text{Fibre } \beta_k; Q).$$

We have a commutative diagram

$$\begin{array}{ccccc} & & M_{B_{k+1}} & & \\ & & \downarrow & & \\ M_{B_k} & \longrightarrow & K_X & \xrightarrow{\beta_k} & K_{A_k} \\ \downarrow l=l_{k+1} & & \downarrow \beta_{k+1} & & \parallel \\ K(Z, \dim \lambda_{k+1}) & \longrightarrow & K_{A_{k+1}} & \longrightarrow & K_{A_k} \end{array}$$

in which the left hand square may be considered a pullback square. Hence $M_{B_{k+1}}$ is homotopy-equivalent to the fibre of l , and we may calculate $H^*(M_{B_{k+1}}; Q)$ from the Serre spectral sequence with rational coefficients, of the fibration

$$\Omega K(Z, \dim \lambda_{k+1}) \rightarrow M_{B_{k+1}} \rightarrow M_{B_k}$$

Let $s = \dim \lambda_{k+1}$. Let us first consider the case in which s is even. For dimensional reasons, the only possibly non-zero differential is d_s , and indeed the transgression of the fundamental class of the fibre is

$$\lambda_{k+1} \bmod (\lambda_1, \dots, \lambda_k) \quad \text{in} \quad \frac{F(\xi_1, \dots, \xi_m)}{(\lambda_1, \dots, \lambda_k)} \simeq H^*(M_{B_k}; Q).$$

Because this element is not a zero-divisor, $d_s^{*, s-1}$ is a monomorphism and $E_\infty^{*, *} \simeq H^*(M_{B_{k+1}}; Q) \simeq F(\xi_1, \dots, \xi_m)/(\lambda_1, \dots, \lambda_{k+1})$ as required.

Now assume that s is odd. The first possibly non-zero differential is again d_s ; the transgression of the fundamental class of the fibre is $\tau(\mathbf{v}_{s-1}) = \lambda_{k+1} \bmod (\lambda_1, \dots, \lambda_k)$. But $H^*(M_{B_k}; Q)$ is a free $F(\tau(\mathbf{v}_{s-1}))$ -module and the kernel of $d_s^{*, p(s-1)}$ for $p > 0$ is exactly the $H^*(M_{B_k}; Q)$ -module with basis $\mathbf{v}_{s-1}^p \otimes \tau(\mathbf{v}_{s-1})$. This coincides with the image of $d_s^{*, (p+1)(s-1)}$. Hence $E_{s+1}^{*, p} \simeq 0$, $p > 0$ and again

$$E_{s+1}^{*, 0} \simeq E_\infty^{*, *} \simeq H^*(M_{B_{k+1}}; Q) \simeq \frac{F(\xi_1, \dots, \xi_m)}{(\lambda_1, \dots, \lambda_{k+1})}.$$

§3. Distance Between Homotopy Types

Let A be a graded-commutative, associative ring such that $A \otimes Q \simeq F(\xi_1, \dots, \xi_m)/(\lambda_1, \dots, \lambda_k) \simeq B$ is regular. If A is the integral cohomology ring of some finite CW complex, it has a highest non-vanishing dimension, say D . Every finite complex with cohomology A will then be homotopy-equivalent to some $(D+1)$ -dimensional CW complex.

Denote a $(D+2)$ -homology section of M_B by M_A [3]. Also let t denote the order of the torsion subgroup of A , considered as an abelian group. Finally let $q: H^*(; Z) \rightarrow H^*(; Q)$ be the coefficient homomorphism induced by the standard inclusion $Z \subset Q$.

Given any finite, simply-connected CW complex X with $H^*(X; Z) \simeq A$, we shall construct a map $\phi: X \rightarrow M_A$ such that $\phi^*: H^*(M_A; Z) \rightarrow A$ has kernel and cokernel of orders bounded by some integer-valued function of the isomorphism class of A . By appealing to the results of [1], this will then be sufficient to deduce that there are at most a finite number of such homotopy types X .

First choose indivisible elements $y_1, y_2, \dots, y_m \in H^*(X; Z)$ such that $q(y_i) = \xi_i$. Let $\alpha: X \rightarrow K_X$ be defined by requiring $\alpha^*(x_i) = t^{d_i} \cdot y_i$, where $d_i = \text{degree } y_i$. Then $\alpha^* \beta^*(l_i) = \lambda_i(t^{d_1} y_1, t^{d_2} y_2, \dots, t^{d_m} y_m) = t^{\text{deg } \lambda_i} \lambda_i(y_1, y_2, \dots, y_m)$ is a torsion element of $H^*(X; Z)$ and hence 0. $\beta \circ \alpha$ is null-homotopic and α lifts to the fibre of β , $\hat{\alpha}: X \rightarrow M_B$. Because X has dimension at most $(D+1)$, the cellular approximation theorem shows that $\hat{\alpha}$ factors through the $(D+1)$ -skeleton of M_B , hence through M_A . It is trivial to verify that this map $\phi: X \rightarrow M_A$ induces an isomorphism on rational cohomology.

The kernel of $\phi^* : H^*(M_A) \rightarrow A$ has order bounded by the order of the torsion subgroup of $H^*(M_A; Z)$, a function of the isomorphism class of A .

The cokernel of ϕ^* is a quotient of the cokernel of α^* . The cokernel of α^* in turn is a quotient of the group $A/(\text{Algebra generated by } \alpha^*x_i, i=1, 2, \dots, m)$, a finite group whose order is a function of the isomorphism class of A .

Thus, according to the terminology of [1], each homotopy type X , with $H^*(X; Z) \simeq \simeq A$ is of bounded distance from M_A , and with the help of Theorem 3.2 of [1] there are only a finite number of such homotopy types. We have demonstrated

THEOREM 3.1. *Let A be a graded-commutative, associative ring such that $A \otimes Q$ has a regular set of relations. Let $HT(A)$ be the class of all homotopy types of simply-connected, finite CW complexes X such that, as graded algebras $H^*(X; Z) \simeq A$. Then $HT(A)$ is a finite set.*

It may be remarked that all such spaces X are 0-universal in the sense of Serre [4]. This may be demonstrated by noting that, for each integer r , there exists endomorphisms $\varrho_1 : K_X \rightarrow K_X$ and $\varrho_2 : K_A \rightarrow K_A$ defined by $\varrho_1^*(x_i) = r^{\dim x_i} x_i$ and $\varrho_2^*(l_i) = r^{\dim l_i} l_i$ inducing an endomorphism of the fibre of β , M_B , hence of M_A which satisfy Mimura, Toda and O'Neill's condition (b') for 0-universality [4].

BIBLIOGRAPHY

- [1] BODY, R. A., and DOUGLAS, R. R., *Homotopy types within a rational homotopy type* (to appear in *Topology*)
- [2] BODY, R. A., *Homotopy types whose rational cohomology is nice*, Notices of the A.M.S. 21, no. 1 (1974), A-222.
- [3] HILTON, PETER, *Homotopy Theory and Duality*, Gordon and Breach, New York, (1965), 89–91.
- [4] MIMURA, O'NEILL and TODA, *On p-equivalence in the sense of Serre*, Japanese Journal of Mathematics, XL (1971), 1–10.
- [5] SAMUEL, O. and ZARISKI, P., *Commutative Algebra*, vol. 2 Appendix VI Van Nostrand, Princeton (1960).
- [6] STASHEFF, J. D., *H-space problems, H-spaces Neuchâtel Août 1970*, Lecture Notes in Mathematics, vol. 196 Springer-Verlag, Berlin (1971), problem 5, p. 123.

I.H.E.S.
 35, rte de Chartres
 Bures-sur-Yvette
 France

Received, February 12, 1974