

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 50 (1975)

**Erratum:** Addendum to the Paper Entitled: "Structural Theorems for Topological Actions of Z2-Torion Real, Complex and Quaternionic Projective Spaces".

**Autor:** Hsiang, Wu Yi

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 20.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**Addendum to the Paper Entitled:  
“Structural Theorems for Topological Actions of  
 $\mathbb{Z}_2$ -Torion Real, Complex and Quaternionic Projective Spaces”**

WU YI HSIANG

In the statement of Theorem 3, there is a missing case that all connected components of  $F(G, X)$  are of *RP*-type. The condition for case 2 should be:  $Sq^2(\zeta) = \gamma \cdot \zeta \neq 0$  and  $Sq^1(\gamma) = 0$ ; and there should be the following third case, namely,

(c) *Case 3,  $Sq^2(\zeta) = \gamma \cdot \zeta \neq 0$  and  $Sq^1(\gamma) \neq 0$ ; then all connected components of  $F(G, X)$  are of *RP*-type, i.e.,  $F^j \sim RP^{(k_j-1)}$  and there exist  $w_j, v_1, v_2 \in H^1(B_G)$  such that*

$$\gamma = v_1^2 + v_1 v_2 + v_2^2, \quad \alpha_j = w_j (w_j + v_1) (w_j + v_1 + v_2)$$

and

$$\iota_j^*(\zeta) = \xi_j^4 + \gamma \xi_j^2 + (Sq^1 \gamma) \xi_j + \alpha_j, \quad \iota_j^*: H_G^*(X) \rightarrow H_G^*(F^j).$$

And furthermore, the system of local weights at  $F^j, \Omega_j$ , is given by

$$\begin{aligned} \Omega_j = & \{(w_i + w_j), (v_1 + w_i + w_j), (v_2 + w_i + w_j), (v_1 + v_2 + w_i + w_j) \text{ multi. } k_i \ (i \neq j); \\ & v_1, v_2, (v_1 + v_2), 0 \text{ multi. } (k_j - 1)\}, \end{aligned}$$

and

$$F(x) \text{ is of } \begin{cases} \text{RP-type} & \text{if } v_1 \mid G_x \text{ and } v_2 \mid G_x \text{ are linearly independent,} \\ \text{CP-type} & \text{if } v_1 \mid G_x \text{ and } v_2 \mid G_x \text{ are linearly dependent but not all zero,} \\ \text{QP-type} & \text{if } v_1 \mid G_x = v_2 \mid G_x = 0. \end{cases}$$

Correspondingly, one should add the following proof to the proof of Theorem 3 for the above case 3:

In the case  $Sq^2(\zeta) = \gamma \cdot \zeta \neq 0$  and  $Sq^1(\gamma) = \delta \neq 0$ , it is not difficult to show that all connected components of  $F(G, X)$  are of *RP*-type. Let  $\xi_j$  be the generator of  $H^*(F^j)$ ,  $\iota_j^*$  be the restriction homomorphism of  $H_G^*(X)$  to  $H_G^*(F^j)$ . Then, it follows from the following equations:

$$Sq^1(\iota_j^*\zeta) = \iota_j^*(Sq^1\zeta) = 0 \quad \text{and} \quad Sq^2(\iota_j^*\zeta) = \iota_j^*(Sq^2\zeta) = \gamma \cdot (\iota_j^*\zeta)$$

that

$$\begin{aligned} i_j^* \zeta &= \xi_j^4 + \gamma \cdot \xi_j^2 + \delta \xi_j + \alpha_j; \\ Sq^1 \delta &= Sq^1 \alpha_j = 0, \quad Sq^2 \delta = \gamma \delta, \quad Sq^2 \alpha_j = \gamma \alpha_j. \end{aligned}$$

Let  $f_j = (0, \dots, 0, \xi_j^{(k_j-1)}, 0, \dots, 0)$  be the fundamental cohomology class of  $F_j$ . Then, simple computation will show that the ideal  $I_X(f_j)$  is generated by  $a(f_j) = \delta^{(k_j-1)} \cdot \prod_{i \neq j} (\alpha_i + \alpha_j)$ , (cf. the proof of Theorem 1). For simplicity, we may assume without loss of generality that  $\alpha_1 = 0$ .

(i) Suppose that at least one  $k_j > 1$ : Then it follows from Theorem B that  $\delta$  splits into product of linear factors, say  $\delta = v_1 \cdot v_2 \cdot v_3$ . Notice that

$$Sq^1 \delta = (v_1 + v_2 + v_3) \cdot \delta = 0 \quad \text{implies} \quad v_1 + v_2 + v_3 = 0,$$

i.e.,

$$\delta = v_1 v_2 (v_1 + v_2),$$

and

$$Sq^2 \delta = \gamma \cdot \delta \quad \text{implies} \quad \gamma = v_1^2 + v_1 v_2 + v_2^2.$$

Again, it follows from Theorem B that  $a(f_1) = \delta^{(k_1-1)} \cdot \prod_{j \neq 1} \alpha_j^{k_j}$  splits, and hence all  $\alpha_j, j \neq 1$ , split into product of linear factors, say

$$\alpha_j = w_j \cdot (w_j + \mu_{j,1}) (w_j + \mu_{j,2}) \cdot (w_j + \mu_{j,3}).$$

On the other hand,

$$\begin{aligned} Sq^1 \alpha_j &= (\mu_{j,1} + \mu_{j,2} + \mu_{j,3}) \cdot \alpha_j = 0 \quad \text{implies that} \quad \mu_{j1} + \mu_{j2} + \mu_{j3} = 0 \\ Sq^2 \alpha_j &= (\mu_{j1} \mu_{j2} + \mu_{j2} \mu_{j3} + \mu_{j3} \mu_{j1}) \cdot \alpha_j = \gamma \cdot \alpha_j \end{aligned}$$

implies that

$$\{\mu_{j1}, \mu_{j2}, \mu_{j3}\} = \{v_1, v_2, v_1 + v_2\}$$

as a set, namely

$$\alpha_j = w_j (w_j + v_1) (w_j + v_2) (w_j + v_1 + v_2).$$

(ii) Suppose all  $k_j = 1$ : Then  $s = (n+1) > 1$ ,  $a(f_1) = \prod_{j \neq 1} \alpha_j$ . Hence,  $\alpha_j$  again splits and the above proof will show that

$$\alpha_j = w_j (w_j + \mu_{j1}) (w_j + \mu_{j2}) (w_j + \mu_{j3}), \quad \mu_{j1} + \mu_{j2} + \mu_{j3} = 0,$$

and

$$(\mu_{j1}\mu_{j2} + \mu_{j2}\mu_{j3} + \mu_{j3}\mu_{j1}) = \gamma \quad \text{for all } j > 1,$$

and the assertion follows easily.

*Remark.* All actions of Case 3 are cohomologically modelled after the following type of linear actions on  $QP^n = Sp(n+1)/Sp(1) \times Sp(n)$ .

EXAMPLE 3c. Notice that the effective group of isometry on the symmetric space  $QP^n$  is  $Sp(n+1)/\{\pm 1\}$  and  $SO(3) = Sp(1)/\{\pm 1\}$  sits in  $Sp(n+1)/\{\pm 1\}$  as diagonal unit quaternions. Let  $G$  be a  $\mathbb{Z}_2$ -tori of  $Sp(n+1)/\{\pm 1\}$  containing a maximal  $\mathbb{Z}_2$ -tori of  $SO(3)$ . Then the restricted  $G$ -action on  $QP^n$  is of the type of Case 3.

### Acknowledgement

The author wishes to thank Mr. Ted Chang and Mr. Tor Skjelbred for pointing out the possibility of the above third case.

Received January 16, 1975.

