

Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	50 (1975)
Artikel:	Irregular Primes and Integrality Theorems for Manifolds.
Autor:	Frank, David
DOI:	https://doi.org/10.5169/seals-38812

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Irregular Primes and Integrality Theorems for Manifolds

DAVID FRANK¹⁾

1. Introduction

Number theory has long played an important part in topology. We examine here a topological problem whose solution depends on the distinction between regular and irregular primes. (Recall that a prime p is said to be irregular if p is odd and p divides the ideal class number of the p -th cyclotomic field; the first irregular prime is 37.)

Let M be a closed oriented topological manifold of dimension $4k$, smooth on the complement of a point. A necessary condition for M itself to be smoothable is that the Pontrjagin number $p_k[M]$ be an integer. If, moreover, M is a spin manifold ($w_2 M = 0$), the integrality theorem of Atiyah-Borel-Hirzebruch-Singer [2], [3], [4] says an additional necessary condition is that $(1/a_k)\hat{A}[M]$ be an integer, where \hat{A} is the \hat{A} -genus and $a_k = 1$ (k even) or 2 (k odd).

We wish to know to what extent these necessary conditions for the smoothability of M are sufficient. If $k > 1$, M is triangulable [11]; then the obstruction to extending the smoothing of (M -point) to M is an element Σ_M of the group of exotic spheres Γ^{4k-1} . There is a splitting [6], [9] $\Gamma^{4k-1} = bP(4k) \oplus \pi'_{4k-1}$, where $bP(4k)$ is the subgroup of exotic spheres bounding parallelizable manifolds and π'_{4k-1} is a certain complementary summand. The integrality of $p_k[M]$ and $\hat{A}[M]$ can provide no information on the component of Σ_M in π'_{4k-1} (Lemma 3.2 below). Hence we let Σ_M^b denote the component of Σ_M in $bP(4k)$, and pose the

Question. Let M be a closed oriented spin manifold of dimension $4k$, $k > 1$, smooth on the complement of a point. Suppose M satisfies the integrality condition: $p_k[M]$ and $(1/a_k)\hat{A}[M]$ are integers. Is $\Sigma_M^b = 0$ in $bP(4k)$?

THEOREM 1.1. *Suppose $\dim M \leq 200$, $\dim M \neq 136$. If M satisfies the integrality condition, then $\Sigma_M^b = 0$.*

THEOREM 1.2. *There is a manifold N of dimension 136 satisfying the integrality condition with Σ_N of order 37 in $bP(136)$.*

Since 37 is the first irregular prime, this suggests

¹⁾ Supported by NSF grant No. P029431000.

THEOREM 1.3. *Let p be a prime. The following are equivalent:*

- i) p is irregular
- ii) for some $k > 1$, there is a manifold M of dimension $4k$ satisfying the integrality condition with order Σ_M^b divisible by p .

In each dimension $4k$, it is possible to determine which primes p (if any) can divide the order of a Σ_M^b (for some M satisfying the integrality condition). Table 1.4 lists all such values of k and p which can occur in the range $k < 109$. If a value of k does not appear in the table, then for a manifold M of dimension $4k$, the integrality condition implies $\Sigma_M^b = 0$.

Table 1.4

$k = \frac{1}{4} \dim M$	34	51	52	62	63	70	76	80	84	88	95	106
$p = \text{order } \Sigma_M^b$	37	59	37	67	103	37	131	59	101	37	67	37

A reference for the number theory we use is [5]. However, we follow the notation of [10], which is standard in topology and differs from the notation of [5].

2. Proof of the Main Theorems

Let B_i be the i -th Bernoulli number [10]. For convenience, define B_0 to be 1. Kummer gave a numerical criterion for regularity.

ASSERTION 2.1. *A prime p is irregular if and only if p divides the numerator of B_n , for some $n < \frac{1}{2}(p-1)$.*

We give a table of all irregular primes less than 233. For each p , we give those $n < \frac{1}{2}(p-1)$ such that p divides num B_n . This information is in fact available for all $p \leq 4001$.

Table 2.2

p	37	59	67	101	103	131	149	157
n	16	22	29	34	12	11	65	31, 55

If k is an integer and p is a prime, let $c (= c(k, p))$ denote the residue class of k modulo $\frac{1}{2}(p-1)$. Thus $0 \leq c < \frac{1}{2}(p-1)$. If $p=2$, set $c(k, 2)=0$.

THEOREM 2.3. *Let $k > 1$ be an integer and p a prime. The following are equivalent:*

- i) $p \leq 2k-1$ and p divides num B_c , where c is the residue class of k modulo $\frac{1}{2}(p-1)$,
- ii) there is a manifold M of dimension $4k$ satisfying the integrality condition with order Σ_M^b divisible by p .

We show that Theorem 2.3, whose proof we defer to the next section, implies the results of the introduction. Note first that (2.3) implies Theorem 1.3. For if p is irregular, then p divides $\text{num } B_c$ for some $c < \frac{1}{2}(p-1)$. Let $k = a(\frac{1}{2})(p-1) + c$, for any integer $a \geq 1$. Then $p \leq 2k-1$ and c is the residue class of $k \pmod{\frac{1}{2}(p-1)}$, so by (2.3) there is a manifold M with the desired properties. Conversely, if M exists, then p divides $\text{num } B_c$, so p is irregular.

We now prove Theorems 1.1 and 1.2. Let $\dim M = 4k$, $k \leq 50$. Suppose M satisfies the integrality condition and p divides the order of Σ_M^b . Then $p \leq 99$ and p is irregular, so $p = 37, 59$, or 67 . Suppose $p = 59$. Then $k \geq 30$. Let c be the residue class of $k \pmod{29}$. Then 59 divides $\text{num } B_c$, and from Table 2.2 we find that $c = 22$. Thus $k \equiv 22 \pmod{29}$ and $30 \leq k \leq 50$. Since no such k exists, we conclude that $p \neq 59$. Similarly, we may exclude $p = 67$.

If $p = 37$, then $k \geq 19$ and, from Table 2.2, $k \equiv 16 \pmod{18}$. Thus $k = 34$ is the only possibility, proving Theorems 1.1 and 1.2.

We leave the further verification of (1.4) to the reader.

3. Proof of Theorem 2.3

We will need

THEOREM (von Staudt). *Let p be a prime. Then p divides $\text{denom } B_k$ if and only if $p-1$ divides $2k$.*

THEOREM (Kummer's Congruence). *Suppose $p-1$ does not divide $2k$, p a prime. Let $r = \frac{1}{2}(p-1)$. Then B_k/k is a p -integer (p does not divide its denominator), and in the ring of p -integers*

$$B_k/k \equiv \pm B_{k+r}/(k+r) \pmod{p}.$$

LEMMA 3.1. *The prime p divides $\text{num}(B_k/k)$ if and only if p divides $\text{num } B_c$, where c is the residue class of $k \pmod{\frac{1}{2}(p-1)}$.*

Proof. Suppose p divides $\text{num}(B_k/k)$. Then by von Staudt's Theorem, $p-1$ does not divide $2k$. (In particular, p is odd.) Write $k = a(\frac{1}{2})(p-1) + c$, where $0 \leq c < \frac{1}{2}(p-1)$. Then Kummer's Congruence shows p divides $\text{num}(B_c/c)$. Hence p divides $\text{num } B_c$.

Conversely, suppose p divides $\text{num } B_c$, where $0 \leq c < \frac{1}{2}(p-1)$. Since $2c < p-1$, we see that p divides $\text{num}(B_c/c)$ and $p-1$ does not divide $2c$. Hence by Kummer, p divides $\text{num}(B_k/k)$.

LEMMA 3.2. *Let M and N be manifolds of dimension $4k$, $k > 1$. If $\Sigma_M^b = \Sigma_N^b$, then $p_k[M] = p_k[N] \pmod{1}$ and $\hat{A}[M] = \hat{A}[N] \pmod{a_k}$.*

Proof. Let $\Sigma = \Sigma_N - \Sigma_M$. Then the $bP(4k)$ -component of Σ is zero. It follows from [6] or [9] that there is a spin manifold V with $\Sigma_V = \Sigma$ such that all Pontrjagin numbers of V are zero. Let X be the connected sum of M , $-N$, and V . Then $\Sigma_X = \Sigma_M - \Sigma_N + \Sigma_V = 0$, so X is smoothable. Thus $p_k[M] - p_k[N] = p_k[M] - p_k[N] + p_k[V] = p_k[X]$ is an integer.

A similar argument applies to the \hat{A} -genus.

Proof of Theorem 2.3. Let W be the closed Milnor manifold of dimension $4k$ and signature 8. Thus W -(point) is smooth and parallelizable. Then $\Sigma_W \in bP(4k)$. In fact, Σ_W generates $bP(4k)$, which is a cyclic group of order $N_k = a_k 2^{2k-2} (2^{2k-1} - 1)$ num (B_k/k).

Now let qW be the connected sum of q copies of W . Then qW is smoothable on the complement of a point and $\Sigma_{qW} = q\Sigma_W$. Note that $\Sigma_{qW} = 0$ if and only if N_k divides q . Suppose M is a manifold of dimension $4k$ which satisfies the integrality condition but $\Sigma_M^b \neq 0$. Then $\Sigma_M^b = \Sigma_{qW}$ for some q . Therefore by Lemma 3.2, the manifold qW satisfies the integrality condition. Using the Hirzebruch signature theorem for the L -genus and the definition of the \hat{A} -genus [10], we have

$$8 = \text{signature } W = L_k[W] = \frac{2^{2k}(2^{2k-1}-1) B_k}{(2k)!} p_k[W]$$

$$\hat{A}[W] = \frac{-B_k}{2(2k)!} p_k[W].$$

We easily compute

$$p_k[qW] = q \frac{(2k)!}{2^{2k-3} (2^{2k-1}-1) B_k}$$

$$\hat{A}[qW] = \frac{-q}{2^{2k-2} (2^{2k-1}-1)}.$$

Now suppose the order of Σ_{qW} is ps , where p is prime and s is an integer. Then we can write

$$q = \frac{rN_k}{ps}, \quad r \text{ prime to } p.$$

Thus

$$p_k[qW] = \frac{4a_k r (2k-1)! \text{ denom}(B_k/k)}{ps}$$

$$\hat{A}[qW] = \frac{(-a_k r) \text{ num}(B_k/k)}{ps}.$$

Then the integrality condition for qW says that p divides $\text{num}(B_k/k)$ and $p \leq 2k-1$. But if p divides $\text{num}(B_k/k)$, then by Lemma 3.1, p divides $\text{num} B_c$, which proves half of Theorem 2.3.

Conversely, if p divides $\text{num} B_c$, then p divides $\text{num}(B_k/k)$. Let $q = N_k/p$. Thus $\text{order } \Sigma_{qW} = p$. If $p \leq 2k-1$, then qW satisfies the integrality condition, concluding the proof of Theorem 2.3.

Remarks. 1. The manifold $N = qW$ ($q = N_k/p$, $p \leq 2k-1$) in fact satisfies the stronger integrality theorem of [2], [3], [4]: $(1/a_k) \langle \hat{A}(N) ph(\gamma), [N] \rangle$ is an integer for every real vector bundle γ on N . Here ph is the Pontrjagin character. Nevertheless qW is not oriented cobordant to a smooth manifold. This does not contradict [12].

2. Note that $(1/a_k) \hat{A}[M] \bmod 1$ is the Eells-Kuiper invariant μ of Σ_M [7]. Similarly $p_k[M] \bmod 1$ is the Milnor invariant $\lambda(\Sigma_M)$ [13]. Thus our results say, for example, that if Σ is an exotic sphere with $\mu(\Sigma) = \lambda(\Sigma) = 0$, then Σ^b has order a product of irregular primes and $\dim \Sigma \geq 135$, or else $\Sigma^b = 0$.

3. There is a manifold N of dimension 436 satisfying the integrality condition with order Σ_N^b equal to the product of distinct primes (59)(157). This is the smallest dimension of such an example.

4. Information on the component of Σ_M in π'_{4k-1} can be obtained using the invariant of [8].

5. For the 136-dimensional manifold of Theorem 1.2 we may take the connected sum of $2^{66}(2^{67}-1)$ (1/37) $\text{num}(B_{34}/34)$ copies of W . Knowing B_{34} (Adams [1]) we find (1/37) $\text{num}(B_{34}/34) =$

$$125, 235, 502, 160, 125, 163, 977, 598, 011, 460, 214, 000, 388, 469.$$

REFERENCES

- [1] ADAMS, J. C., *Table of the first sixty-two numbers of Bernoulli*, J. reine angew. Math. (Crelle) 85 (1878), 269–272.
- [2] ATIYAH, M. F. and HIRZEBRUCH, F., *Riemann-Roch theorems for differentiable manifolds*, Bull. Amer. Math. Soc. 65 (1959), 276–281.
- [3] ATIYAH, M. F. and SINGER, I. M., *The index of elliptic operators III*, Ann. of Math. 87 (1968), 546–604.
- [4] BOREL, A. and HIRZEBRUCH, F., *Characteristic classes and homogeneous spaces III*, Amer. J. Math. 82 (1960), 491–504.
- [5] BOREVICH, Z. I. and SHAFAREVICH, I. R., *Number Theory*, Academic Press, 1966.
- [6] BRUMFIEL, G., *On the homotopy groups of BPL and PL/O*, Ann. of Math. 88 (1968), 291–311.
- [7] EELLS, J. and KUIPER, N., *An invariant for certain smooth manifolds*, Annali di Matematica 60 (1962), 93–110.
- [8] FRANK, D., *An invariant for almost-closed manifolds*, Bull. Amer. Math. Soc. 74 (1968), 562–567.
- [9] ———, *The signature defect and the homotopy of BPL and PL/O*, Comment. Math. Helv. 48 (1973), 525–530.
- [10] HIRZEBRUCH, F., *Topological Methods in Algebraic Geometry*, Springer-Verlag, 1966.

- [11] KIRBY, R. and SIEBENMANN, L., *Some theorems on topological manifolds*, Manifolds-Amsterdam 1970. Lecture Notes in Math. 197, Springer, 1971.
- [12] STONG, R. E., *Relations among characteristic numbers II*, Topology 5 (1966), 133–148.
- [13] MILNOR, J., *Differential structures on spheres*, Amer. J. Math. 81 (1959), 962–972.

*State University of New York
at Stony Brook*

Current address:
University of Minnesota

Received July 24, 1974