

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 51 (1976)

Artikel: Aspherical Manifolds and Higher-Dimensional Knots
Autor: Eckmann, Beno
DOI: <https://doi.org/10.5169/seals-39430>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 25.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Aspherical Manifolds and Higher-Dimensional Knots

BENO ECKMANN

E. Dyer and R. Vasquez [3] proved that the complement of a higher-dimensional knot $S^{n-2} \subset S^n$, $n \geq 4$, is never aspherical unless the knot group is infinite cyclic (and hence, for $n \geq 5$, the imbedding is unknotted¹⁾). In the present note we give a simple proof of this fact based on some remarks concerning compact ∂ -manifolds. By the same method we show that the complement of a link in S^n , $n \geq 4$, is never aspherical.

Let X be a compact n -dimensional ∂ -manifold, $G = \pi_1(X)$ its fundamental group. If ∂X is connected, let G_0 be the image of $\pi_1(\partial X)$ in G . Using the connection between G_0 and the boundary $\partial \tilde{X}$ of the universal cover of X we first note that $H^{n-1}(X; \mathbf{Z}G) = 0$ if and only if $G_0 = G$. If, moreover, X is aspherical, we show that $\text{cd } G_0 < n - 1$ implies $G_0 = G$ (and vice-versa). Since in the case of a knot-complement in S^n , $n \geq 4$, the image G_0 is infinite cyclic, the Dyer-Vasquez result follows. Actually the asphericity is used here in a weak form only, cf. 3.2 below. – In the case where ∂X is not connected, and if X is aspherical, then for at least one of the components of ∂X one has $\text{cd } G_0 = n - 1$. This immediately implies the sphericity of higher-dimensional links.

1. The Fundamental Group of a ∂ -Manifold

1.1. Let X be a ∂ -manifold; by this we mean a connected cellular manifold with non-empty boundary ∂X . We write i for the inclusion map $\partial X \rightarrow X$, and G for the fundamental group $\pi_1(X)$.

We will always assume X to be compact. The universal cover \tilde{X} of X is a ∂ -manifold which may be compact or not; its boundary $\partial \tilde{X}$ is the inverse image $p^{-1}(\partial X)$ under the covering map $p: \tilde{X} \rightarrow X$. We want to get information on the number of connected components of $\partial \tilde{X}$; i.e., on the integral homology group $H_0(\partial \tilde{X})$. The exact homology sequence

$$\dots \rightarrow H_1(\tilde{X}) \rightarrow H_1(\tilde{X} \text{ mod } \partial \tilde{X}) \rightarrow H_0(\partial \tilde{X}) \rightarrow H_0(\tilde{X}) \rightarrow 0$$

yields

$$H_0^{\text{red}}(\partial \tilde{X}) = H_1(\tilde{X} \text{ mod } \partial \tilde{X}),$$

¹⁾ In [3], only $n \geq 6$ is mentioned (Levine, Stallings), but the result holds for $n = 5$ as well (C. T. C. Wall, Shaneson).

where H_0^{red} is the reduced homology group. Poincaré duality in \tilde{X} further yields

$$H_0^{\text{red}}(\partial\tilde{X}) = \bar{H}^{n-1}(\tilde{X}),$$

n being the dimension of X , and \bar{H} denoting cohomology with compact supports (i.e., if we use a cell decomposition, cohomology based on finite cellular cochains). If $C(\tilde{X})$ denotes the chain complex of \tilde{X} corresponding to a finite cell decomposition of X , one may replace (cf. [2], p. 359) the finite cochain group $\text{Hom}_{\text{fin}}(C(\tilde{X}), \mathbf{Z})$ by the equivariant group $\text{Hom}_G(C(\tilde{X}), \mathbf{Z}G)$. It follows that

$$\bar{H}^{n-1}(X) = H^{n-1}(X; \mathbf{Z}G),$$

the last group being cohomology with local coefficients given by the left G -module $\mathbf{Z}G$. We thus obtain

$$H_0^{\text{red}}(\partial\tilde{X}) = H^{n-1}(X; \mathbf{Z}G). \quad (1)$$

This yields the following results:

PROPOSITION 1.1. *$\partial\tilde{X}$ is connected if and only if $H^{n-1}(X; \mathbf{Z}G) = 0$.*

PROPOSITION 1.2. *If ∂X is not connected, then $H^{n-1}(X; \mathbf{Z}G) \neq 0$.*

1.2. We now assume the boundary manifold ∂X to be connected and write G_0 for $i_*\pi_1(\partial X)$, the image of $\pi_1(\partial X)$ under the inclusion map $i: \partial X \rightarrow X$. The connected components of $\partial\tilde{X} = p^{-1}(\partial X)$ correspond bijectively to the cosets of G modulo G_0 . Proposition 1.1 can therefore be reformulated as follows.

PROPOSITION 1.3. *Let X be a compact manifold of dimension n with connected boundary ∂X . Then $H^{n-1}(X; \mathbf{Z}G) = 0$ if and only if $G_0 = G$; i.e., if $\pi_1(\partial X) \rightarrow \pi_1(X)$ is surjective.*

Let $K(G_0, 1)$ denote an Eilenberg-MacLane complex of the group G_0 . There is a map $j: \partial X \rightarrow K(G_0, 1)$, determined up to homotopy, which induces the surjection $\pi_1(\partial X) \rightarrow G_0$. We now further assume that the inclusion $i: \partial X \rightarrow X$ can be factored up to homotopy through j :

$$i = hj: \partial X \xrightarrow{j} K(G_0, 1) \xrightarrow{h} X. \quad (2)$$

Then $i^*: H^{n-1}(X; \mathbf{Z}G) \rightarrow H^{n-1}(\partial X; \mathbf{Z}G)$ is factored as $i^* = j^*h^*$ through the cohomology group $H^{n-1}(G_0; \mathbf{Z}G)$ and will thus be 0 if we assume this group to be 0

(in particular, if the cohomology dimension $\text{cd } G_0$ is $< n-1$). The homomorphism i^* appears in the exact sequence with local coefficients

$$\dots \rightarrow H^{n-1}(X \text{ mod } \partial X; \mathbf{Z}G) \rightarrow H^{n-1}(X; \mathbf{Z}G) \xrightarrow{i^*} H^{n-1}(\partial X; \mathbf{Z}G) \rightarrow \dots \quad (3)$$

By Poincaré duality $H^{n-1}(X \text{ mod } \partial X; \mathbf{Z}G) = H_1(X; \mathbf{Z}G)$; the latter group is computed from $\mathbf{Z}G \otimes_G C(\tilde{X}) = C(\tilde{X})$, i.e., it is equal to $H_1(\tilde{X})$ and hence 0.

Note that the argument is valid both in the orientable and non-orientable case: in the non-orientable case the duality yields $H^{n-1}(X \text{ mod } \partial X; \mathbf{Z}G) = H_1(X; \check{\mathbf{Z}} \otimes \mathbf{Z}G)$, where $\check{\mathbf{Z}}$ is the group of twisted integers. But $\check{\mathbf{Z}} \otimes \mathbf{Z}G$ (with diagonal action) is easily seen to be isomorphic to $\mathbf{Z}G$.

Thus by (3) i^* is always injective. Under the factorization assumption (2), and if $H^{n-1}(G_0; \mathbf{Z}G) = 0$, we have seen that $i^* = 0$, and therefore $H^{n-1}(X; \mathbf{Z}G) = 0$. Combining this with Prop. 1.3 we get

THEOREM 1.4 *Let X be a compact manifold of dimension n with connected boundary ∂X , and let $i: \partial X \rightarrow X$ be the inclusion, $G = \pi_1(X)$, $G_0 = i_* \pi_1(\partial X)$. If i can be factored up to homotopy as $i = hj: \partial X \rightarrow K(G_0, 1) \rightarrow X$ and if $H^{n-1}(G_0; \mathbf{Z}G) = 0$, then $G_0 = G$.*

1.3. In Theorem 1.4 the condition $H^{n-1}(G_0; \mathbf{Z}G) = 0$ can be replaced by $H_{n-1}(G_0) = 0$.

To prove this, let $e \in H_{n-1}(\partial X)$ be the fundamental class of ∂X [$e \in H_{n-1}(\partial X; \check{\mathbf{Z}}$) in the non-orientable case]. For any $z \in H^{n-1}(G_0; \mathbf{Z}G)$, the cap-product formula

$$j_*(e \cap j^*z) = j_*e \cap z$$

together with $j_*e = 0$ yields $j^*z = 0$, since $j_*: H_0(\partial X; \mathbf{Z}G) \rightarrow H_0(G_0; \mathbf{Z}G)$ and $e \cap -$ are both isomorphisms. Now $j^* = 0$ implies $i^* = 0$.

1.4. If we do not assume that ∂X is connected, Theorem 1.4 has to be restated in a slightly different form.

Let $\partial_\nu X$, $\nu = 0, 1, \dots, k$ be the connected components of ∂X , and $G_\nu = i_{\nu*} \pi_1(\partial_\nu X)$ the images in G of their fundamental groups under the inclusions $i_\nu: \partial_\nu X \rightarrow X$ (determined up to conjugacy only). Let K be the disjoint union of the $K(G_\nu, 1)$ and $j: \partial X \rightarrow K$ the union of the maps $j_\nu: \partial_\nu X \rightarrow K(G_\nu, 1)$ inducing $i_{\nu*}$. If i can be factored up to homotopy as $i = hj: \partial X \rightarrow K \rightarrow X$ and if $H^{n-1}(G_\nu; \mathbf{Z}G) = 0$ for all ν (or: if $H_{n-1}(G_\nu) = 0$ for all ν) then it follows as above that $H^{n-1}(X; \mathbf{Z}G) = 0$; i.e., ∂X must be connected, $k = 0$, $G = G_0$.

THEOREM 1.5. *Let X be a compact ∂ -manifold, $G = \pi_1(X)$ and $G_v = i_{v*}\pi_1(\partial_v X)$, $v = 0, \dots, k$. If $i: \partial X \rightarrow X$ can be factored as $i = hj: \partial X \rightarrow K \rightarrow X$ and if $H^{n-1}(G_v; \mathbf{Z}G) = 0$ (or: $H_{n-1}(G_v) = 0$) for $v = 0, \dots, k$, then ∂X is connected and $G_0 = G$.*

2. Aspherical Manifolds

2.1. The notations being as in 1.1, we now assume the manifold X to be aspherical; in other words, an Eilenberg-MacLane complex $K(G, 1)$ for its fundamental group $G = \pi_1(X)$. Since cohomology of X with local coefficients vanishes in dimensions $k \geq n$, the cohomology dimension $\text{cd} G$ is $\leq n - 1$. Note that the chain complex $C(\tilde{X})$ constitutes a finitely generated free resolution for G ; therefore $H^{n-1}(G; \mathbf{Z}G) = H^{n-1}(X; \mathbf{Z}G) = 0$ implies $H^{n-1}(G; A) = 0$ for all free G -modules A and hence (cf. [1] p. 105) for all G -modules A , and thus is equivalent to $\text{cd} G < n - 1$.

The results of Section 1 can now be applied as follows.

PROPOSITION 2.1. *Let G be a group admitting a $K(G, 1) = X$ which is a compact manifold of dimension n with non-empty boundary ∂X . Then $\text{cd} G < n - 1$ if and only if $\partial \tilde{X}$ is connected; in particular, if ∂X is not connected then $\text{cd} G = n - 1$.*

Note that any group admitting a $K(G, 1)$ which is a finite cell-complex admits a compact manifold with non-empty boundary as Eilenberg-MacLane space (imbed $K(G, 1)$ in some \mathbf{R}^N and take a regular neighborhood).

2.2. For aspherical X , assuming ∂X connected, the factorization (2) of $i: \partial X \rightarrow X$ is always possible. Hence Theorem 1.4 yields

THEOREM 2.2. *Let G be a group admitting a $K(G, 1) = X$ which is a compact manifold of dimension n with connected boundary ∂X , and $G_0 = i_*\pi_1(\partial X)$. Then $G_0 = G$ if and only if $\text{cd} G_0 < n - 1$. In other words, one always has $\text{cd} G = \text{cd} G_0$; namely, $< n - 1$ if $G_0 = G$ and $= n - 1$ if $G_0 \neq G$.*

From Theorem 1.5 we immediately get

THEOREM 2.3. *Let G be a group admitting a $K(G, 1) = X$ which is a compact ∂ -manifold of dimension n . If ∂X is not connected, then $\text{cd} G_v = n - 1$ for at least one component $\partial_v X$ of ∂X , G_v being the image of $\pi_1(\partial_v X)$ under the inclusion.*

3. Higher-dimensional Knots

3.1. Let throughout this section $S^{n-2} \subset S^n$, $n \geq 4$, be a knot, i.e., a differentiable imbedding of S^{n-2} in S^n , $C = S^n - S^{n-2}$ its complement, and X its closed comple-

ment $S^n - V^n$ where V^n is an open tubular neighborhood of S^{n-2} in S^n . Then X and C have the same homotopy type, and $G = \pi_1(X)$ is the corresponding knot group. ∂X is a product $S^1 \times S^{n-2}$, and $\pi_1(\partial X) \cong \mathbf{Z}$ imbeds injectively into G .

THEOREM 3.1. *If the knot complement is aspherical, then $G \cong \mathbf{Z}$.*

Proof. Since $i_*\pi_1(\partial X) = G_0 \cong \mathbf{Z}$, we have $\text{cd} G_0 = 1$, and hence Theorem 2.2 applies: $G_0 = G \cong \mathbf{Z}$.

3.2. Note that the asphericity of X is not used in full here. The factorization (2) of i can be obtained under weaker assumptions, as follows.

Let $j: \partial X \rightarrow S^1 = K(G_0, 1)$ be the projection of $\partial X = S^1 \times S^{n-2}$ onto $S^1 \times pt$, and h the imbedding $S^1 \times pt \rightarrow \partial X \rightarrow X$. If we assume

(a) a sphere $pt \times S^{n-2} \subset \partial X$ is nullhomotopic in X

then hj and i can be made, by a homotopy, to agree on $S^1 \times pt \vee pt \times S^{n-2}$. If we further assume

(b) $\pi_{n-1}(X) = 0$

then i and hj are homotopic, and thus, by Theorem 1.4, $G = G_0 \cong \mathbf{Z}$.

THEOREM 3.1'. *If $pt \times S^{n-2}$ is nullhomotopic in the knot complement C and if $\pi_{n-1}(C) = 0$ then $G \cong \mathbf{Z}$.*

3.3. G. A. Swarup [4] has proved that Theorem 3.1' holds without the assumption that $\pi_{n-1}(C) = 0$ provided G is *accessible*. Since it is conjectured that all finitely generated groups are accessible, it is possible that the nullhomotopy of S^{n-2} in C alone is sufficient to conclude that $G \cong \mathbf{Z}$.

3.4. HIGHER-DIMENSIONAL LINKS. If X is the closed complement of a link

$$\bigcup_{v=0, \dots, k} S_v^{n-2} \subset S^n, \quad n \geq 4, \quad k > 0,$$

then ∂X is not connected. The images G_v of $\pi_1(\partial_v X)$ are all $\cong \mathbf{Z}$. By Theorem 2.3 X can not be aspherical.

THEOREM 3.2. *The complement of a link in S^n , $n \geq 4$, is never aspherical.*

REFERENCES

- [1] R. BIERI and B. ECKMANN, *Groups with homological duality generalizing Poincaré duality*, *Inv. Math.* 20 (1973), 103–124.
- [2] H. CARTAN and S. EILENBERG, *Homological Algebra*, Princeton University Press, 1956.
- [3] E. DYER and A. T. VASQUEZ, *The sphericity of higher dimensional knots*, *Can. J. Math.* 25 (1973), 1132–1136.
- [4] G. A. SWARUP, *Accessible groups and an unknotting criterion* (to appear in *J. of Pure and Applied Algebra*).

E.T.H. Zürich

Received September, 1975