

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 51 (1976)

Artikel: A New Bound for the Genus of a Nilpotent Group
Autor: Lemaire, Claude
DOI: <https://doi.org/10.5169/seals-39434>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 21.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A New Bound for the Genus of a Nilpotent Group

CLAUDE LEMAIRE

Introduction

Gr is the category of groups with the usual morphisms; $[\text{Gr}]$ the class of isomorphism classes of Gr ; η_0 the full subcategory of Gr consisting of all finitely generated infinite nilpotent groups with finite commutator subgroups; N_p and \hat{N}_p are the p -localization and the p -completion of the nilpotent group N .

Assume that N is a finitely generated nilpotent group. In the sense of Pickel ([1]) the genus set of N , denoted by $G_p(N)$, is the set of isomorphism classes of the finitely generated nilpotent groups M with $\hat{N}_p \simeq \hat{M}_p$ for each prime p , and $N_0 \simeq M_0$ (N_0 is the rationalization of N). In the sense of Mislin, the genus set of N , $G(N)$, is the set of the finitely generated nilpotent groups M with $M_p \simeq N_p$ for each prime p (see [2]).

In this note, we are only concerned with groups N in η_0 ; in this case, it has just been proved by Warfield in [3] that $G_p(N) = G(N)$.⁽¹⁾ This result is also an easy consequence of our proof.

This paper provides a bound for $G_p(N)$, when N is in η_0 , thus a bound for $G(N)$ (see the theorem); examples in section 3 show that this bound can be an improvement of Mislin's one ([2]).

1. Preliminaries

For a nilpotent group N , we denote its maximal torsion subgroup by TN , the p -components by T_pN and the quotient N/TN by SN .

LEMMA 1. *Let N be in η_0 , M a finitely generated nilpotent group with $\hat{M}_p \simeq \hat{N}_p$ for each prime p . Then*

$$(a) \quad TN \simeq TM$$

$$(b) \quad SN \simeq SM$$

This research was supported by the National Research Council of Canada.

⁽¹⁾ I am very grateful to Professor Mislin for having reported me this fact before the publication of [3].

- (c) M belongs to η_0 (d) $N_0 \simeq M_0$
 (e) the class of M belongs to $G_p(N)$.

Proof. (a) is proved by Pickel ([1], proposition 3.5).

(b) SN is abelian (since $[N, N]$ is included in TN) thus free abelian; then SM is also abelian as a subgroup of \widehat{SM}_p which is isomorphic to \widehat{SN}_p and SM is free abelian with the same rank since SN and SM have isomorphic finite quotients ([1], proposition 3.5).

(c) Since SM is abelian, $[M, M]$ is included in TM which is finite.

(d) $N_0 \simeq (SN)_0 \simeq (SM)_0 \simeq M_0$.

(e) obvious, by hypothesis and d.

Among the invariants in $G_p(N)$, we have thus

- the groups TN and SN , which we shall denote simply by T and S .
- the rank k of S .
- the set Q of primes p for which T_p is non-trivial.

The lemma shows us that the class of M belongs to $G_p(N)$ if and only if $\hat{N}_p \simeq \hat{M}_p$ for each prime p and that each element in $G_p(N)$ can be described as the class of the central term M of an extension:

$$T \twoheadrightarrow M \twoheadrightarrow S$$

With this description, we have only to consider primes in Q , since, for $p \notin Q$, $\hat{M}_p \simeq \hat{S}_p \simeq \hat{N}_p$. This is the basis of the proof of the theorem; before stating it, we need two further notions.

From [4] we recall the definition of Blackburn's function for a prime p .

It is the function defined recursively by

$$d_p(0) = 0$$

$$d_p(n) = d_p(n-1) + m$$

where p^m is the highest power of p that does not exceed n .

With the aid of d_p , we introduce

$$u(p) = t(p) + d_p(\text{nil } N) \quad \text{where} \quad \exp T = \prod_{p \in Q} p^{t(p)}$$

and

$$u = \prod_{p \in Q} p^{u(p)}$$

2. The Main Result

THEOREM. *Let N be in η_0 . Then $|G_p(N)| \leq \phi(u)/2$ (ϕ is Euler's function and we assume $u > 2$).*

Proof. Since all the products have Q as their index set, we shall omit the subscripts. Define $E(X, Y)$ to be the class of equivalence classes of all the extensions

$$Y \twoheadrightarrow K \rightarrow X$$

θ is the map from $E(S, T)$ to $[Gr]$ defined by "take the isomorphism class of the central term."

We know that $G_p(N)$ is included in $\text{Im } \theta$. In particular, we can choose x_0 in $E(S, T)$ such that $\theta(x_0)$ is the class of N .

We intend to build a diagram Δ :

$$\begin{array}{ccccc} \hat{\Omega} = \prod GL(k, \hat{\mathbb{Z}}_p) & & & & \\ \downarrow \delta & \searrow \lambda & & & \\ GL(k, \mathbb{Z}/u) & \xrightarrow{\mu} & E(S, T) & \xrightarrow{\theta} & [Gr] \\ \downarrow \gamma & & \nearrow \nu & & \\ (\mathbb{Z}/u)^*/\{1, -1\} & & & & \end{array}$$

where $(\mathbb{Z}/u)^*/\{1, -1\}$ is the cokernel of $(GL(k, \mathbb{Z}) \rightarrow GL(k, \mathbb{Z}/u))$; γ, δ are the canonical epimorphisms; the two triangles are commutative; $G_p(N) = \text{Im } \nu = \text{Im } (\theta \circ \mu)$.

The result follows from $|(\mathbb{Z}/u)^*/\{1, -1\}| = \phi(u)/2$.

First step: Definition of λ

For each p in Q , there exists a map C_p from $E(S, T)$ to $E(\hat{S}_p, T_p)$ defined by completion (we know that the completion is exact). Let us denote by \hat{C} the product map. C_p can be factorized into a map τ_p from $E(S, T)$ to $E(S, T_p)$:

$$\tau_p(T \twoheadrightarrow E \rightarrow S) = (T_p \twoheadrightarrow E/T_{p'}, E \rightarrow S)$$

(p' is the complement of p in the set of prime numbers) and the p -completion \bar{C}_p , applied now to $E(S, T_p)$ see [] p. 339). We denote by τ and \bar{C} the product maps.

On the other hand, we can define:

a map pb from $\prod E(\hat{S}_p, T_p)$ to $\prod E(S, T_p)$ by pull-back along the canonical injections of S in \hat{S}_p
 and a map PB from $\prod E(S, T_p)$ to $E(S, T)$ defined by

$$\left(\begin{array}{ccc} T_p \longrightarrow E_p \xrightarrow{\sigma_p} S \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ T_q \longrightarrow E_q \xrightarrow{\sigma_q} S \end{array} \right) \mapsto (T \longrightarrow E \longrightarrow S)$$

where E is the pull-back of the groups E_p along the morphisms σ_p .

It is easy to check that \bar{C} and pb , τ and PB , are inverses of one another. Thus:

$$E(S, T) \simeq \prod E(\hat{S}_p, T_p)$$

Now, each ω_p belonging to $GL(k, \hat{\mathbb{Z}}_p)$ acts on $E(\hat{S}_p, T_p)$ by pull-back along ω_p^{-1} . We define $\hat{\Omega} = \prod GL(k, \hat{\mathbb{Z}}_p)$ and

$$\lambda: \hat{\Omega} \rightarrow E(S, T): (\omega_p) \mapsto \hat{C}^{-1}(\omega_p C_p(x_0))$$

Second step: Definition of μ

Consider δ to be the canonical epimorphism from $\hat{\Omega}$ onto $GL(k, \mathbb{Z}/u) \simeq \prod GL(k, \mathbb{Z}/p^{u(p)})$.

The existence of μ and the commutativity of the upper triangle of Δ are proved by the

LEMMA 2. *If ω belongs to $\ker \delta$, then $\lambda(\omega) = x_0$.*

Proof. For simplicity, if \hat{X}_p is a p -complete nilpotent group, we denote by $\hat{X}_p^{(u)}$ the subgroup of \hat{X}_p generated by the $u(p)$ -powers and by $\hat{X}_p^{/u}$ the quotient of \hat{X}_p modulo $\hat{X}_p^{(u)}$. If ω belongs to $\ker \delta$, the map induced by ω_p on $\hat{S}_p^{/u}$ is the identity. On the other hand, by our choice of $u(p)$, each element of $\hat{N}_p^{(u)}$ is a $p^{(p)}$ -power ([4], Lemma 2), thus $\text{Im } T_p \cap \hat{N}_p^{(u)} = \{1\}$. We can

consider the commutative diagrams

$$\begin{array}{ccccc} T_p & \longrightarrow & \hat{N}_p & \twoheadrightarrow & \hat{S}_p \\ \parallel & & \downarrow & & \downarrow \\ T_p & \longrightarrow & \hat{N}_p^{/u} & \twoheadrightarrow & \hat{S}_p^{/u} \end{array}$$

where the right squares are pull-backs. The action of the ω_p 's does not modify the bottoms, neither, consequently, the tops, and thus $\lambda(\omega) = x_0$.

Third Step: $\text{Im}(\mu \circ \theta) = G_p(N)$

If the class of M belongs to $\text{Im}(\mu \circ \theta)$, M is clearly finitely generated and nilpotent, and \hat{M}_p is isomorphic to \hat{N}_p for each p ; Lemma 1 shows that the class of M belongs to $G_p(N)$.

Suppose now that the class of M belongs to $G_p(N)$. We know that there exists

$$x: T \longrightarrow M \longrightarrow S$$

and, for each p , a commutative diagram

$$\begin{array}{ccccc} \hat{C}_p x_0: T_p & \longrightarrow & \hat{N}_p & \longrightarrow & \hat{S}_p \\ \xi_p \downarrow & & \epsilon_p \downarrow & & \omega_p \downarrow \\ \hat{C}_p x: T_p & \longrightarrow & \hat{M}_p & \longrightarrow & \hat{S}_p \end{array}$$

We state that M is isomorphic to the central term of $\mu(\delta(\omega)) = \lambda(\omega)$, where $\omega = (\omega_p)$. Indeed, there exists, for each p , a commutative diagram

$$\begin{array}{ccccc} T_p & \longrightarrow & N_{\omega_p} & \longrightarrow & \hat{S}_p \\ \xi_p \downarrow & & \epsilon_p \downarrow & & \parallel \\ T_p & \longrightarrow & \hat{M}_p & & \hat{S}_p \end{array}$$

the first row being $\omega_p C_p x_0$. Applying pb and then PB , we obtain the required isomorphism.

Fourth step: Definition of ν

We can identify $\text{GL}(k, \mathbb{Z})$ with a subgroup of $\hat{\Omega}$. Now we prove:

LEMMA 3. *Let ω_1, ω_2 be in $\hat{\Omega}$ and suppose there exists η in $\text{GL}(k, \mathbb{Z})$ such that $\omega_2 = \eta \omega_1$. Then $(\theta \circ \lambda) \omega_1 = (\theta \circ \lambda) \omega_2$.*

Proof. For each p , there exists a commutative diagram

$$\begin{array}{ccccc} T_p & \longrightarrow & N_{\omega_{1,p}} & \longrightarrow & \hat{S}_p \\ || & & \downarrow \wr & & \downarrow \wr \\ T_p & \longrightarrow & N_{\omega_{2,p}} & \longrightarrow & \hat{S}_p \end{array}$$

where the rows are $\omega_{1,p}C_px_0$ and $\omega_{2,p}C_px_0$ and η_p is the extension of η to \hat{S}_p . Such a diagram still exists after applying pb , with the same η for each p and PB produces isomorphic central terms.

In order to define ν and assure the commutativity of the lower triangle of Δ , we have to prove that $(\theta \circ \mu) \delta(\omega)$ only depends on the class of $\delta(\omega)$ modulo $\delta(\text{GL}(k, \mathbb{Z}))$. Suppose $\delta(\omega_2) = \delta(\eta) \cdot \delta(\omega_1)$. Then: $(\theta \circ \mu) \delta(\omega_2) = (\theta \circ \mu) \delta(\eta\omega_1) = (\theta \circ \lambda)(\eta\omega_1) = (\theta \circ \lambda)\omega_1$ (by Lemma 3) $= (\theta \circ \mu) \delta(\omega_1)$. The proof of the theorem is completed.

Remark. The proof can be rewritten with p -localization instead of p -completions. All differences disappear after the second step, and the third implies the equality, in η_0 , of $G_p(N)$ and $G(N)$ (see [3] for another proof.)

3. Comparison with Mislin's Result

Since the theorem gives a bound for $|G(N)|$, it is interesting to compare this bound to that established by Mislin in [2]. We adopt the improvement of Mislin's bound introduced at the end of [5] and we follow the notation of this paper as far as possible. We have to compare $u(p) = t(p) + d_p(\text{nil } N)$ to the exponents $v(p)$ of $\exp(QN)_{ab}$ where $QN = N/(ZN)^{\exp TZN}$. First, two remarks.

If N is abelian, $N = ZN$, $TN = TZN$, $\exp(QN)_{ab} = \exp TN$ and $d_p(1) = 0$ for any p : the two bounds are always equal. More generally, if we assume that TN is central (and $2 \notin Q$), then $\text{nil } N \leq 2$ and $d_p(\text{nil } N) = 0$; $(QB)_{ab}$ contains at least a copy of $\mathbb{Z}/\exp TN$, and $u(p) \leq v(p)$. In this case, we can have $u(p) < v(p)$ (see below).

Our bound can give new information, as it is shown by the following class of examples (a modified version of examples elaborated by Professor G. Frei, whom I thank here). The proof is standard and we omit it.

Consider groups N with generators $a_1, a_2, a_3, \dots, a_{2n+1}$, relations $[a_i, a_j] = 1$ except if i is odd and $j = i+1$; in this case, $[a_i, a_{i+1}] = a_{i+2}$; we assume also that the a_i 's are of finite order for $i \in J = \{3, 5, \dots, 2n+1\}$. From $[a_i^m, a_{i+1}] = a_{i+2}^m$ ($i = 3, 5, \dots, 2n-1$), we conclude that the orders r_i of a_i ($i \in J$) are related by $r_{2n+1} | r_{2n-1} \cdots | r_3$. We assume $r_{2n+1} \neq 1$, and we write s_i for r_i/r_{i+2} ($i \in J$) with $r_{2n+3} = 1$. For $i \geq 2$, $\Gamma_i(N)$ is generated by $\{a_j | j \in J, j \geq 2i-1\}$. Thus

N is nilpotent of class $n+1$ and its commutator subgroup is finite. ZN is generated by $a_1^{r_3}, a_2^{r_3}, a_3^{r_5}, \dots, a_{2n+1}$ and $TZN \cong \prod_{i \in J} \mathbb{Z}/s_i$; its exponent is the LCM of the s_i 's, denoted by s . r_{2n+1} divides s such that s is not 1. $ZN^{\exp TZN}$ is the free abelian subgroup generated by $a_1^{sr_3}, a_2^{sr_3}, a_4^{sr_5}, \dots$. Finally, $(QN)_{ab}$ is the abelian group with $a_1, a_2, a_4, \dots, a_{2n}$ as generators and relations $a_1^{sr_3} = 1, a_2^{sr_3} = 1, \dots, a_{2n}^{sr_{2n+1}} = 1$; its exponent is sr_3 . On the other hand, TN is generated by $\{a_i \mid i \in J\}$. It is abelian and its exponent is r_3 .

If $x(p)$ represents the exponent of the p -component of the natural x ,

$$v(p) - u(p) = s(p) - d_p(n+1)$$

where

$$s(p) = \max (r_i(p) - r_{i+2}(p))$$

$$s(p) \geq r_{2n+1}(p);$$

$$s(p) \geq 1 \quad \text{for each } p \text{ in } Q$$

We can conclude:

- for any choice of the nilpotency class except 1 (n fixed) and any choice of $Q(2 \notin Q)$ there exists a group in η_0 for which our bound is strictly smaller than Mislin's one: it is sufficient to choose r_{2n+1} adequately.
- on the other hand, if we fix a bound for the exponent of TN , it is always possible (now choosing n) to find a group in η_0 for which Mislin's bound is better.

REFERENCES

- [1] P. F. PICKEL, *Finitely generated nilpotent groups with isomorphic finite quotients*, Trans. AMS, 160 (1971), 327–341.
- [2] G. MISLIN, *Nilpotent groups with finite commutator subgroups*, in Lecture notes in Mathematics, no. 148, Springer-Verlag, (1974), 103–120.
- [3] R. B. WARFIELD, *Genus and cancellation for groups with finite commutator subgroup*, to appear in the Journal of Pure and Applied Algebra (1975).
- [4] N. BLACKBURN, *Conjugacy in nilpotent groups*, Proc. AMS 16 (1965) 143–148, MR 30 #3140.
- [5] P. HILTON and G. MISLIN, *On the genus of a nilpotent group with finite commutator subgroup* (to appear).

Département de Mathématiques
Université Laval
Québec, Canada, G1K 7P4

Received August 13, 1975

