

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 51 (1976)

**Artikel:** A New Bound for the Genus of a Nilpotent Group  
**Autor:** Lemaire, Claude  
**DOI:** <https://doi.org/10.5169/seals-39434>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 30.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# A New Bound for the Genus of a Nilpotent Group

CLAUDE LEMAIRE

## Introduction

$\text{Gr}$  is the category of groups with the usual morphisms;  $[\text{Gr}]$  the class of isomorphism classes of  $\text{Gr}$ ;  $\eta_0$  the full subcategory of  $\text{Gr}$  consisting of all finitely generated infinite nilpotent groups with finite commutator subgroups;  $N_p$  and  $\hat{N}_p$  are the  $p$ -localization and the  $p$ -completion of the nilpotent group  $N$ .

Assume that  $N$  is a finitely generated nilpotent group. In the sense of Pickel ([1]) the genus set of  $N$ , denoted by  $G_p(N)$ , is the set of isomorphism classes of the finitely generated nilpotent groups  $M$  with  $\hat{N}_p \simeq \hat{M}_p$  for each prime  $p$ , and  $N_0 \simeq M_0$  ( $N_0$  is the rationalization of  $N$ ). In the sense of Mislin, the genus set of  $N$ ,  $G(N)$ , is the set of the finitely generated nilpotent groups  $M$  with  $M_p \simeq N_p$  for each prime  $p$  (see [2]).

In this note, we are only concerned with groups  $N$  in  $\eta_0$ ; in this case, it has just been proved by Warfield in [3] that  $G_p(N) = G(N)$ .<sup>(1)</sup> This result is also an easy consequence of our proof.

This paper provides a bound for  $G_p(N)$ , when  $N$  is in  $\eta_0$ , thus a bound for  $G(N)$  (see the theorem); examples in section 3 show that this bound can be an improvement of Mislin's one ([2]).

## 1. Preliminaries

For a nilpotent group  $N$ , we denote its maximal torsion subgroup by  $TN$ , the  $p$ -components by  $T_p N$  and the quotient  $N/TN$  by  $SN$ .

LEMMA 1. *Let  $N$  be in  $\eta_0$ ,  $M$  a finitely generated nilpotent group with  $\hat{M}_p \simeq \hat{N}_p$  for each prime  $p$ . Then*

$$(a) \quad TN \simeq TM$$

$$(b) \quad SN \simeq SM$$

---

This research was supported by the National Research Council of Canada.

<sup>(1)</sup> I am very grateful to Professor Mislin for having reported me this fact before the publication of [3].

*Proof.* (a) is proved by Pickel ([1], proposition 3.5).

(b)  $SN$  is abelian (since  $[N, N]$  is included in  $TN$ ) thus free abelian; then  $SM$  is also abelian as a subgroup of  $\widehat{SM}_p$  which is isomorphic to  $\widehat{SN}_p$  and  $SM$  is free abelian with the same rank since  $SN$  and  $SM$  have isomorphic finite quotients ([1], proposition 3.5).

- (c) Since  $SM$  is abelian,  $[M, M]$  is included in  $TM$  which is finite.
- (d)  $N_0 \simeq (SN)_0 \simeq (SM)_0 \simeq M_0$ .
- (e) obvious, by hypothesis and *d*.

Among the invariants in  $G_p(N)$ , we have thus

- the groups  $TN$  and  $SN$ , which we shall denote simply by  $T$  and  $S$ .
- the rank  $k$  of  $S$ .
- the set  $Q$  of primes  $p$  for which  $T_p$  is non-trivial.

The lemma shows us that the class of  $M$  belongs to  $G_p(N)$  if and only if  $\hat{N}_p \simeq \hat{M}_p$  for each prime  $p$  and that each element in  $G_p(N)$  can be described as the class of the central term  $M$  of an extension:

$$T \rightarrowtail M \twoheadrightarrow S$$

With this description, we have only to consider primes in  $Q$ , since, for  $p \notin Q$ ,  $\hat{M}_p \simeq \hat{S}_p \simeq \hat{N}_p$ . This is the basis of the proof of the theorem; before stating it, we need two further notions.

From [4] we recall the definition of Blackburn's function for a prime  $p$ . It is the function defined recursively by

$$d_p(0) = 0$$

$$d_p(n) = d_p(n-1) + m$$

where  $p^m$  is the highest power of  $p$  that does not exceed  $n$ .

With the aid of  $d_p$ , we introduce

$$u(p) = t(p) + d_p(\text{nil } N) \quad \text{where} \quad \exp T = \prod_{p \in Q} p^{t(p)}$$

and

$$u = \prod_{p \in Q} p^{u(p)}$$

## 2. The Main Result

**THEOREM.** *Let  $N$  be in  $\eta_0$ . Then  $|G_p(N)| \leq \phi(u)/2$  ( $\phi$  is Euler's function and we assume  $u > 2$ ).*

*Proof.* Since all the products have  $Q$  as their index set, we shall omit the subscripts. Define  $E(X, Y)$  to be the class of equivalence classes of all the extensions

$$Y \longrightarrow K \longrightarrow X$$

$\theta$  is the map from  $E(S, T)$  to  $[\text{Gr}]$  defined by “take the isomorphism class of the central term.”

We know that  $G_p(N)$  is included in  $\text{Im } \theta$ . In particular, we can choose  $x_0$  in  $E(S, T)$  such that  $\theta(x_0)$  is the class of  $N$ .

We intend to build a diagram  $\Delta$ :

$$\begin{array}{ccccc} \hat{\Omega} = \prod & \text{GL}(k, \hat{\mathbb{Z}}_p) & & & \\ \downarrow \delta & & \searrow \lambda & & \\ \text{GL}(k, \mathbb{Z}/u) & \xrightarrow{\mu} & E(S, T) & \xrightarrow{\theta} & [\text{Gr}] \\ \downarrow \gamma & & \nearrow \nu & & \\ (\mathbb{Z}/u)^*/\{1, -1\} & & & & \end{array}$$

where  $(\mathbb{Z}/u)^*/\{1, -1\}$  is the cokernel of  $(\text{GL}(k, \mathbb{Z}) \rightarrow \text{GL}(k, \mathbb{Z}/u))$ ;  $\gamma, \delta$  are the canonical epimorphism; the two triangles are commutative;  $G_p(N) = \text{Im } \nu = \text{Im } (\theta \circ \mu)$ .

The result follows from  $|(\mathbb{Z}/u)^*/\{1, -1\}| = \phi(u)/2$ .

*First step: Definition of  $\lambda$*

For each  $p$  in  $Q$ , there exists a map  $C_p$  from  $E(S, T)$  to  $E(\hat{S}_p, T_p)$  defined by completion (we know that the completion is exact). Let us denote by  $\hat{C}$  the product map.  $C_p$  can be factorized into a map  $\tau_p$  from  $E(S, T)$  to  $E(S, T_p)$ :

$$\tau_p(T \longrightarrow E \rightarrow S) = (T_p \longrightarrow E/T_{p'}, E \rightarrow S)$$

( $p'$  is the complement of  $p$  in the set of prime numbers) and the  $p$ -completion  $\bar{C}_p$ , applied now to  $E(S, T_p)$  see [ ] p. 339). We denote by  $\tau$  and  $\bar{C}$  the product maps.

On the other hand, we can define:

a map  $pb$  from  $\prod E(\hat{S}_p, T_p)$  to  $\prod E(S, T_p)$  by pull-back along the canonical injections of  $S$  in  $\hat{S}_p$

and a map  $PB$  from  $\prod E(S, T_p)$  to  $E(S, T)$  defined by

$$\left( \begin{array}{ccc} T_p & \longrightarrow & E_p \xrightarrow{\sigma_p} S \\ \cdot & & \cdot \\ T_q & \longrightarrow & E_q \xrightarrow{\sigma_q} S \end{array} \right) \mapsto (T \longrightarrow E \longrightarrow S)$$

where  $E$  is the pull-back of the groups  $E_p$  along the morphisms  $\sigma_p$ .

It is easy to check that  $\bar{C}$  and  $pb$ ,  $\tau$  and  $PB$ , are inverses of one another. Thus:

$$E(S, T) \simeq \prod E(\hat{S}_p, T_p)$$

Now, each  $\omega_p$  belonging to  $GL(k, \hat{\mathbb{Z}}_p)$  acts on  $E(\hat{S}_p, T_p)$  by pull-back along  $\omega_p^{-1}$ . We define  $\hat{\Omega} = \prod GL(k, \hat{\mathbb{Z}}_p)$  and

$$\lambda : \hat{\Omega} \rightarrow E(S, T) : (\omega_p) \mapsto \hat{C}^{-1}(\omega_p C_p(x_0))$$

*Second step: Definition of  $\mu$*

Consider  $\delta$  to be the canonical epimorphism from  $\hat{\Omega}$  onto  $GL(k, \mathbb{Z}/u) \simeq \prod GL(k, \mathbb{Z}/p^{u(p)})$ .

The existence of  $\mu$  and the commutativity of the upper triangle of  $\Delta$  are proved by the

**LEMMA 2.** *If  $\omega$  belongs to  $\ker \delta$ , then  $\lambda(\omega) = x_0$ .*

*Proof.* For simplicity, if  $\hat{X}_p$  is a  $p$ -complete nilpotent group, we denote by  $\hat{X}_p^{(u)}$  the subgroup of  $\hat{X}_p$  generated by the  $u(p)$ -powers and by  $\hat{X}_p^{(u)}$  the quotient of  $\hat{X}_p$  modulo  $\hat{X}_p^{(u)}$ . If  $\omega$  belongs to  $\ker \delta$ , the map induced by  $\omega_p$  on  $\hat{S}_p^{(u)}$  is the identity. On the other hand, by our choice of  $u(p)$ , each element of  $\hat{N}_p^{(u)}$  is a  $p^{u(p)}$ -power ([4], Lemma 2), thus  $\text{Im } T_p \cap \hat{N}_p^{(u)} = \{1\}$ . We can

consider the commutative diagrams

$$\begin{array}{ccccc} T_p & \longrightarrow & \hat{N}_p & \longrightarrow & \hat{S}_p \\ \parallel & & \downarrow & & \downarrow \\ T_p & \longrightarrow & \hat{N}'^u_p & \longrightarrow & \hat{S}'^u_p \end{array}$$

where the right squares are pull-backs. The action of the  $\omega_p$ 's does not modify the bottoms, neither, consequently, the tops, and thus  $\lambda(\omega) = x_0$ .

*Third Step:  $\text{Im}(\mu \circ \theta) = G_p(N)$*

If the class of  $M$  belongs to  $\text{Im}(\mu \circ \theta)$ ,  $M$  is clearly finitely generated and nilpotent, and  $\hat{M}_p$  is isomorphic to  $\hat{N}_p$  for each  $p$ ; Lemma 1 shows that the class of  $M$  belongs to  $G_p(N)$ .

Suppose now that the class of  $M$  belongs to  $G_p(N)$ . We know that there exists

$$x : T \longrightarrow M \longrightarrow S$$

and, for each  $p$ , a commutative diagram

$$\begin{array}{ccccc} \hat{C}_p x_0 : T_p & \longrightarrow & \hat{N}_p & \longrightarrow & \hat{S}_p \\ \xi_p \downarrow & & \epsilon_p \downarrow & & \omega_p \downarrow \\ \hat{C}_p x : T_p & \longrightarrow & \hat{M}_p & \longrightarrow & \hat{S}_p \end{array}$$

We state that  $M$  is isomorphic to the central term of  $\mu(\delta(\omega)) = \lambda(\omega)$ , where  $\omega = (\omega_p)$ . Indeed, there exists, for each  $p$ , a commutative diagram

$$\begin{array}{ccccc} T_p & \longrightarrow & N_{\omega_p} & \longrightarrow & \hat{S}_p \\ \xi_p \downarrow & & \epsilon_p \downarrow & & \parallel \\ T_p & \longrightarrow & \hat{M}_p & \longrightarrow & \hat{S}_p \end{array}$$

the first row being  $\omega_p C_p x_0$ . Applying  $pb$  and then  $PB$ , we obtain the required isomorphism.

*Fourth step: Definition of  $\nu$*

We can identify  $\text{GL}(k, \mathbb{Z})$  with a subgroup of  $\hat{\Omega}$ . Now we prove:

LEMMA 3. *Let  $\omega_1, \omega_2$  be in  $\hat{\Omega}$  and suppose there exists  $\eta$  in  $\text{GL}(k, \mathbb{Z})$  such that  $\omega_2 = \eta \omega_1$ . Then  $(\theta \circ \lambda)\omega_1 = (\theta \circ \lambda)\omega_2$ .*

*Proof.* For each  $p$ , there exists a commutative diagram

$$\begin{array}{ccccc} T_p & \longrightarrow & N_{\omega_{1,p}} & \longrightarrow & \hat{S}_p \\ \parallel & & \downarrow \wr & & \downarrow \wr \\ T_p & \longrightarrow & N_{\omega_{2,p}} & \longrightarrow & \hat{S}_p \end{array}$$

where the rows are  $\omega_{1,p}C_p x_0$  and  $\omega_{2,p}C_p x_0$  and  $\eta_p$  is the extension of  $\eta$  to  $\hat{S}_p$ . Such a diagram still exists after applying  $pb$ , with the same  $\eta$  for each  $p$  and  $PB$  produces isomorphic central terms.

In order to define  $\nu$  and assure the commutativity of the lower triangle of  $\Delta$ , we have to prove that  $(\theta \circ \mu) \delta(\omega)$  only depends on the class of  $\delta(\omega)$  modulo  $\delta(\text{GL}(k, \mathbb{Z}))$ . Suppose  $\delta(\omega_2) = \delta(\eta) \cdot \delta(\omega_1)$ . Then:  $(\theta \circ \mu) \delta(\omega_2) = (\theta \circ \mu) \delta(\eta \omega_1) = (\theta \circ \lambda)(\eta \omega_1) = (\theta \circ \lambda) \omega_1$  (by Lemma 3)  $= (\theta \circ \mu) \delta(\omega_1)$ . The proof of the theorem is completed.

*Remark.* The proof can be rewritten with  $p$ -localization instead of  $p$ -completions. All differences disappear after the second step, and the third implies the equality, in  $\eta_0$ , of  $G_p(N)$  and  $G(N)$  (see [3] for another proof.)

### 3. Comparison with Mislin's Result

Since the theorem gives a bound for  $|G(N)|$ , it is interesting to compare this bound to that established by Mislin in [2]. We adopt the improvement of Mislin's bound introduced at the end of [5] and we follow the notation of this paper as far as possible. We have to compare  $u(p) = t(p) + d_p(\text{nil } N)$  to the exponents  $v(p)$  of  $\exp(QN)_{ab}$  where  $QN = N/(ZN)^{\exp TN}$ . First, two remarks.

If  $N$  is abelian,  $N = ZN$ ,  $TN = TZN$ ,  $\exp(QN)_{ab} = \exp TN$  and  $d_p(1) = 0$  for any  $p$ : the two bounds are always equal. More generally, if we assume that  $TN$  is central (and  $2 \notin Q$ ), then  $\text{nil } N \leq 2$  and  $d_p(\text{nil } N) = 0$ ;  $(QB)_{ab}$  contains at least a copy of  $\mathbb{Z}/\exp TN$ , and  $u(p) \leq v(p)$ . In this case, we can have  $u(p) < v(p)$  (see below).

Our bound can give new information, as it is shown by the following class of examples (a modified version of examples elaborated by Professor G. Frei, whom I thank here). The proof is standard and we omit it.

Consider groups  $N$  with generators  $a_1, a_2, a_3, \dots, a_{2n+1}$ , relations  $[a_i, a_j] = 1$  except if  $i$  is odd and  $j = i+1$ ; in this case,  $[a_i, a_{i+1}] = a_{i+2}$ ; we assume also that the  $a_i$ 's are of finite order for  $i \in J = \{3, 5, \dots, 2n+1\}$ . From  $[a_i^m, a_{i+1}] = a_{i+2}^m$  ( $i = 3, 5, \dots, 2n-1$ ), we conclude that the orders  $r_i$  of  $a_i$  ( $i \in J$ ) are related by  $r_{2n+1} | r_{2n-1} \cdots | r_3$ . We assume  $r_{2n+1} \neq 1$ , and we write  $s_i$  for  $r_i/r_{i+2}$  ( $i \in J$ ) with  $r_{2n+3} = 1$ . For  $i \geq 2$ ,  $\Gamma_i(N)$  is generated by  $\{a_j \mid j \in J, j \geq 2i-1\}$ . Thus

$N$  is nilpotent of class  $n+1$  and its commutator subgroup is finite.  $ZN$  is generated by  $a_1^{r_3}, a_2^{r_3}, a_3^{r_3}, \dots, a_{2n+1}$  and  $TZN \simeq \prod_{i \in J} \mathbb{Z}/s_i$ ; its exponent is the *LCM* of the  $s_i$ 's, denoted by  $s$ .  $r_{2n+1}$  divides  $s$  such that  $s$  is not 1.  $ZN^{\exp TZN}$  is the free abelian subgroup generated by  $a_1^{sr_3}, a_2^{sr_3}, a_4^{sr_3}, \dots$ . Finally,  $(QN)_{ab}$  is the abelian group with  $a_1, a_2, a_4, \dots, a_{2n}$  as generators and relations  $a_1^{sr_3} = 1, a_2^{sr_3} = 1, \dots, a_{2n}^{sr_{2n+1}} = 1$ ; its exponent is  $sr_3$ . On the other hand,  $TN$  is generated by  $\{a_i \mid i \in J\}$ . It is abelian and its exponent is  $r_3$ .

If  $v(p)$  represents the exponent of the  $p$ -component of the natural  $x$ ,

$$v(p) - u(p) = s(p) - d_p(n+1)$$

where

$$s(p) = \max (r_i(p) - r_{i+2}(p))$$

$$s(p) \geq r_{2n+1}(p);$$

$$s(p) \geq 1 \text{ for each } p \text{ in } Q$$

We can conclude:

- for any choice of the nilpotency class except 1 ( $n$  fixed) and any choice of  $Q (2 \notin Q)$  there exists a group in  $\eta_0$  for which our bound is strictly smaller than Mislin's one: it is sufficient to choose  $r_{2n+1}$  adequately.
- on the other hand, if we fix a bound for the exponent of  $TN$ , it is always possible (now choosing  $n$ ) to find a group in  $\eta_0$  for which Mislin's bound is better.

## REFERENCES

- [1] P. F. PICKEL, *Finitely generated nilpotent groups with isomorphic finite quotients*, Trans. AMS, 160 (1971), 327–341.
- [2] G. MISLIN, *Nilpotent groups with finite commutator subgroups*, in Lecture notes in Mathematics, no. 148, Springer-Verlag, (1974), 103–120.
- [3] R. B. WARFIELD, *Genus and cancellation for groups with finite commutator subgroup*, to appear in the Journal of Pure and Applied Algebra (1975).
- [4] N. BLACKBURN, *Conjugacy in nilpotent groups*, Proc. AMS 16 (1965) 143–148, MR 30 #3140.
- [5] P. HILTON and G. MISLIN, *On the genus of a nilpotent group with finite commutator subgroup* (to appear).

Département de Mathématiques  
Université Laval  
Québec, Canada, G1K 7P4

Received August 13, 1975

