

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 52 (1977)

Artikel: On the inverse limit of free nilpotent groups.
Autor: Baumslag, G. / Stambach, U.
DOI: <https://doi.org/10.5169/seals-39996>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On the inverse limit of free nilpotent groups

G. BAUMSLAG and U. STAMMBACH

0. Introduction

A group P is called *parafree* (see [B1], [B2]) if it is residually nilpotent and if there exists a free group F and a homomorphism $\varphi: F \rightarrow P$ such that φ induces isomorphisms $\varphi_i: F/F_i \xrightarrow{\sim} P/P_i$, $i \geq 2$ modulo the terms F_i , P_i of the lower central series. Since F is residually nilpotent, the map φ is injective, so that F may be thought of as a subgroup of P . If $F = F(X)$ is free on the set X , then P is called parafree on X . It is plain (see [B2]) that a parafree group on X can be embedded in $\hat{F} = \varprojlim F/F_i$. The group \hat{F} thus certainly merits some interest. This paper is a contribution to the study of the group \hat{F} .

In section 1 we introduce some notation and prove a basic lemma which enables us to identify the subgroups of $\hat{F} = \hat{F}(X)$ which are parafree on X . In section 2 we deal with the case of finitely generated free groups $F = F(X)$. It turns out (Corollary 2.2) that in this case the group \hat{F} is parafree on X . This result contrasts with the case where X is infinite. In section 3 we deal in detail with the case where X is countably infinite. We prove the following results: \hat{F} is *not* parafree on X (Corollary 3.5; see also [BK]); \hat{F}_{ab} contains uncountably many linearly independent divisible elements (Theorem 3.9); \hat{F} contains a free subgroup of uncountable rank with a generating set which is linearly independent mod \hat{F}_2 (Theorem 3.6); the 2-generator subgroups of \hat{F} are free (Theorem 3.11). We note that the restriction to the countable case in all of the main results of this section is not decisive. The conclusions remain true if X is allowed to be uncountable.

In section 4 we define two subgroups \bar{F} , \tilde{F} of $\hat{F} = \hat{F}(X)$. The group \bar{F} is the union of the subgroups of \hat{F} which are parafree on X . It is shown that \bar{F} is parafree on X , so that \bar{F} is the universal parafree group on X in the sense that it contains all groups which are parafree on X (Theorem 4.1). The subgroup \tilde{F} of \hat{F} consists of all elements of \hat{F} which can be expressed by finitely many elements of X . It is shown that \tilde{F} too is parafree on X (Proposition 4.4). Hence clearly $\tilde{F} \subseteq \bar{F}$. But we show that $\tilde{F} \neq \bar{F}$ if X is countably infinite (Proposition 4.5).

We also show that the group \bar{F} is freely indecomposable (Corollary 4.3). We were however unable to settle the question whether \tilde{F} and \hat{F} are freely indecomposable in case X is infinite.

1. The inverse limit

Let G be a group and let $\{G_i\}$ denote its lower central series, i.e.

$$G_1 = G, \quad G_i = [G, G_{i-1}], \quad i \geq 2. \quad (1.1)$$

As usual we shall denote G/G_2 by G_{ab} . We consider the inverse system of the canonical projections

$$\{G/G_i \rightarrow G/G_{i-1}\}. \quad (1.2)$$

Its inverse limit is denoted by

$$\hat{G} = \varprojlim G/G_i \quad (1.3)$$

and its canonical maps by $\tau_i: \hat{G} \rightarrow G/G_i$. We may regard \hat{G} as the subgroup of the (categorical) product $\prod_{i=2}^{\infty} G/G_i$ consisting of the elements

$$\Lambda = (\lambda_1 G_2, \lambda_2 G_3, \dots) \quad (1.4)$$

with $\lambda_i \in G$ and $\lambda_{i+1} \equiv \lambda_i$ modulo G_{i+1} . Then clearly $\tau_i(\Lambda) = \lambda_{i-1} G_i \in G/G_i$. By universality of the inverse limit the family $\pi_i: G \rightarrow G/G_i$ of canonical projections induces a homomorphism $h: G \rightarrow \hat{G}$ such that $\pi_i = \tau_i h$. Plainly it is given by

$$h(x) = (xG_2, xG_3, \dots), \quad x \in G. \quad (1.5)$$

The homomorphism h is injective if and only if G is residually nilpotent (i.e. if $G_\omega = \bigcap_{i \geq 2} G_i = e$).

In the sequel we shall be interested in subgroups P of \hat{G} with $hG \subseteq P \subseteq \hat{G}$ and with the property that h induces isomorphisms $h_i: G/G_i \xrightarrow{\sim} P/P_i$, $i \geq 2$. The following lemma characterizes these subgroups.

LEMMA 1.1. *Let P be a group with $hG \subseteq P \subseteq \hat{G}$. Then the following statements are equivalent*

- (i) $h: G \rightarrow P$ induces isomorphisms $h_i: G/G_i \xrightarrow{\sim} P/P_i$, $i \geq 2$;
- (ii) $\tau_i: P \rightarrow \hat{G} \rightarrow G/G_i$ induces isomorphisms $\sigma_i: P/P_i \xrightarrow{\sim} G/G_i$, $i \geq 2$;
- (iii) $h: G \rightarrow P$ induces an epimorphism $h_2: G_{ab} \twoheadrightarrow P_{ab}$;
- (iv) $\tau_2: P \rightarrow G_{ab}$ induces a monomorphism $\sigma_2: P_{ab} \hookrightarrow G_{ab}$.

Proof. We consider the map

$$\pi_i = \tau_i h: G \rightarrow P \subseteq \hat{G} \rightarrow G/G_i, \quad i \geq 2. \quad (1.6)$$

It induces the identity $G/G_i \xrightarrow{h_i} P/P_i \xrightarrow{\sigma_i} G/G_i$, $i \geq 2$. Hence h_i is always injective and σ_i surjective. Moreover h_i is surjective if and only if σ_i is injective. This proves the equivalence of (i) and (ii). Also, it is well known that h_i is surjective if and only if h_2 is, proving the equivalence of (i) and (iii). Finally h_2 is surjective if and only if σ_2 is injective, proving the equivalence of (iii) and (iv).

COROLLARY 1.2. *Let $F = F(X)$ be the free group on the set X . Then a group P with $F \subseteq P \subseteq \hat{F}$ is parafree on X if and only if $h_2: F_{ab} \rightarrow P_{ab}$ is surjective.*

Proof. This is immediate from Lemma 1.1, since the groups F and \hat{F} are clearly residually nilpotent.

2. The case of finitely generated free groups

Let $F = F(X)$ be the free group on the set $X = \{x_1, x_2, \dots, x_n\}$. The following result is due to Bousfield–Kan [BK]. Since its proof to be found in [BK] uses topological methods we shall include, for completeness, a purely algebraic proof; it is also due to Bousfield–Kan.

THEOREM 2.1. *Let F be a finitely generated free group. Then $h: F \rightarrow \hat{F}$ induces isomorphisms $h_i: F/F_i \xrightarrow{\sim} \hat{F}/\hat{F}_i$, $i \geq 2$.*

COROLLARY 2.2. *If $F = F(x_1, \dots, x_n)$, then \hat{F} is parafree on $X = \{x_1, \dots, x_n\}$.*

Proof. By Lemma 1.1 we only have to show that $\sigma_2: \hat{F}_{ab} \rightarrow F_{ab}$ induced by $\tau_2: \hat{F} \rightarrow F/F_2$ is injective. We thus have to show that an element

$$\Lambda^* = (\lambda_1 F_2, \lambda_2 F_3, \dots) \in \hat{F} \quad (2.1)$$

with $\lambda_1 \in F_2$ is in \hat{F}_2 . In the course of the proof we shall need the following two wellknown results which we mention without proof.

LEMMA 2.3.

$$[ab, c] = [a, c]^b [b, c],$$

$$[c, ab] = [c, b][c, a]^b.$$

LEMMA 2.4. *Let $F = F(x_1, \dots, x_n)$. Then given $a \in F_k$, $k \geq 2$ there exist*

$u_1, \dots, u_n \in F_{k-1}$ such that

$$a \equiv [u_1, x_1][u_2, x_2] \cdots [u_n, x_n] \pmod{F_{k+1}}.$$

We shall construct elements

$$\Gamma^{(i)} = (\gamma_1^{(i)} F_2, \gamma_2^{(i)} F_3, \dots) \in \hat{F}, \quad 1 \leq i \leq n \quad (2.2)$$

such that in \hat{F}

$$\Lambda^* = [\Gamma^{(1)}, h(x_1)][\Gamma^{(2)}, h(x_2)] \cdots [\Gamma^{(n)}, h(x_n)]. \quad (2.3)$$

In order to find $\gamma_k^{(i)}$ we proceed by induction on k . Since $\lambda_1 \in F_2$ and hence $\lambda_2 \in F_2$ there are, by Lemma 2.4, elements $u_1, \dots, u_n \in F_1$ such that

$$\lambda_2 \equiv [u_1, x_1] \cdots [u_n, x_n] \pmod{F_3}. \quad (2.4)$$

Set $\gamma_1^{(i)} = u_i$. We then have

$$\lambda_1 \equiv [\gamma_1^{(1)}, x_1] \cdots [\gamma_1^{(n)}, x_n] \pmod{F_2} \quad (2.5)$$

and also

$$\lambda_2 \equiv [\gamma_1^{(1)}, x_1] \cdots [\gamma_1^{(n)}, x_n] \pmod{F_3}. \quad (2.6)$$

Suppose now that $\gamma_1^{(i)}, \dots, \gamma_k^{(i)}, 1 \leq i \leq n$ have already been determined such that

$$\gamma_{l+1}^{(i)} \equiv \gamma_l^{(i)} \pmod{F_{l+1}}, \quad 1 \leq l \leq k-1, \quad (2.7)$$

$$\lambda_l \equiv [\gamma_l^{(1)}, x_1] \cdots [\gamma_l^{(n)}, x_n] \pmod{F_{l+1}}, \quad 1 \leq l \leq k, \quad (2.8)$$

and in addition

$$\lambda_{k+1} \equiv [\gamma_k^{(1)}, x_1] \cdots [\gamma_k^{(n)}, x_n] \pmod{F_{k+2}}. \quad (2.9)$$

Since $\lambda_{k+2} \equiv \lambda_{k+1} \pmod{F_{k+2}}$ there exists $r_{k+2} \in F_{k+2}$ such that

$$\lambda_{k+2} = ([\gamma_k^{(1)}, x_1] \cdots [\gamma_k^{(n)}, x_n])r_{k+2}. \quad (2.10)$$

By Lemma 2.4 we can find $v_i \in F_{k+1}$ such that

$$r_{k+2} \equiv [v_1, x_1] \cdots [v_n, x_n] \pmod{F_{k+3}}. \quad (2.11)$$

We may thus set

$$\gamma_{k+1}^{(i)} = \gamma_k^{(i)} \cdot v_i, \quad 1 \leq i \leq n. \quad (2.12)$$

We then clearly have

$$\gamma_{k+1}^{(i)} \equiv \gamma_k^{(i)} \pmod{F_{k+1}}. \quad (2.13)$$

Moreover

$$\begin{aligned} [\gamma_{k+1}^{(1)}, x_1] \cdots [\gamma_{k+1}^{(n)}, x_n] &= [\gamma_k^{(1)} v_1, x_1] \cdots [\gamma_k^{(n)} v_n, x_n] \\ &= [\gamma_k^{(1)}, x_1]^{v_1} [v_1, x_1] \cdots [\gamma_k^{(n)}, x_n]^{v_n} [v_n, x_n], \text{ by Lemma 2.3} \\ &\equiv ([\gamma_k^{(1)}, x_1] \cdots [\gamma_k^{(n)}, x_n]) ([v_1, x_1] \cdots [v_n, x_n]) \pmod{F_{k+3}} \\ &\equiv \lambda_{k+2} \pmod{F_{k+3}}, \text{ by (2.10), (2.11)}. \end{aligned} \quad (2.14)$$

A fortiori we have

$$[\gamma_{k+1}^{(1)}, x_1] \cdots [\gamma_{k+1}^{(n)}, x_n] \equiv \lambda_{k+1} \pmod{F_{k+2}}. \quad (2.15)$$

This completes the proof of Theorem 2.1.

For the proof of Corollary 2.2 we only have to remark that \hat{F} , being a subgroup of $\prod_{i \geq 2} F/F_i$, is residually nilpotent. Next we recall that the group \hat{F} contains any parafree group on X (see [B2]). Since, by Corollary 2.2, the group \hat{F} is itself parafree on X if X is finite, we see that \hat{F} is the biggest parafree group on the (finite) set X .

We note that the proof of Theorem 2.1 works equally well if the free group F is replaced by an arbitrary group G generated by the set $X = \{x_1, \dots, x_n\}$. We may thus state

COROLLARY 2.5. *If G is finitely generated, then $\hat{G} = \hat{G}$.*

Finally we note

COROLLARY 2.6. *If F is finitely generated, then \hat{F} is freely indecomposable.*

Proof. Suppose $\hat{F} = A * B$ with $A \neq \{e\} \neq B$. Then there are surjective maps $A \twoheadrightarrow C_1$, $B \twoheadrightarrow C_2$ with C_i , $i = 1, 2$ infinite cyclic. Thus we obtain an epimorphism $f: \hat{F} \rightarrow C_1 * C_2 = F(x, y)$ onto the free group on two generators. Take elements $a, b \in \hat{F}$ with $fa = x$, $fb = y$. Since $F(x, y)$ is free there exists $g: F(x, y) \rightarrow \hat{F}$ with $gx = a$, $gy = b$ and $fg = Id_F$. Using Corollary 2.5 we obtain an epimorphism $p: \hat{F}(x, y) \rightarrow \hat{F} \cong \hat{F} \rightarrow F(x, y)$. But, by Corollary 2.2 the groups $\hat{F}(x, y)$ and $F(x, y)$

are parafree of the same rank, so that by Theorem 1.1 of [B2] p is an isomorphism and $\hat{F}(x, y)$ is free. This is a contradiction.

3. The case of free groups of countably infinite rank

Let $F = F(x_1, x_2, \dots)$ be a free group on the countably infinite set $X = \{x_1, x_2, \dots\}$. In order to obtain results on \hat{F} we shall first construct a metabelian group W .

Let A be the free abelian group on $X = \{x_1, x_2, \dots\}$ and let IA be the augmentation ideal of the integral group ring of A . Clearly IA/IA^2 is free abelian on $\{x_1 - 1, x_2 - 1, \dots\}$. The following lemma is a generalization of this fact (see [BG]).

LEMMA 3.1. IA^n/IA^{n+1} is the free abelian group on the set $\{\prod_{j=1}^n (x_{i(j)} - 1)\}$.

We now define W as the semi-direct product $W = IA \rtimes A$ where IA is regarded as right A -module in the usual way. We prove

LEMMA 3.2. $W_n = IA^n$, $n \geq 2$.

Proof. Let $u, v \in IA$, $x, y \in A$; then using Lemma 2.3 we obtain

$$[ux, vy] = [u, vy]^x [x, vy] = [u, y]^x [u, v]^{yx} [x, y][x, v]^y = [u, y]^x [x, v]^y,$$

since both IA and A are abelian. Thus we have $W_2 = [W, W] = [IA, A] = IA^2$. Using induction it is now easy to prove $W_n = IA^n$ for $n > 2$. We leave the details to the reader. We now consider $\hat{W} = \varprojlim W/W_n = \varprojlim (IA/IA^n \rtimes A) = (\varprojlim IA/IA^n) \rtimes A$. By Lemma 3.1 the group IA^n/IA^{n+1} is free abelian on the n -fold products $\prod_{i=1}^n (x_{i(j)} - 1)$. We may thus identify IA/IA^n , as abelian group, with the augmentation ideal of the quotient of the polynomial ring on $y_i = x_i - 1$, $i = 1, 2, \dots$ modulo the ideal generated by the n -fold products. As a consequence we see that $\varprojlim IA/IA^n$, as an abelian group, is isomorphic to the augmentation ideal J of the power series ring on $y_i = x_i - 1$, $i = 1, 2, \dots$. The operation of x_j, x_j^{-1} on $y_i = x_i - 1$ is given by

$$(x_i - 1) \circ x_j = (x_i - 1)(x_j - 1) + (x_i - 1) = y_i y_j + y_i, \quad (3.1)$$

$$(x_i - 1) \circ x_j^{-1} = y_i(1 - y_j + y_j^2 - y_j^3 + \dots). \quad (3.2)$$

We note for further reference that in $\hat{W} = J \upharpoonright A$

$$[x_i - 1, x_j] = -(x_i - 1) + (x_i - 1) \circ x_j = y_i y_j, \quad (3.3)$$

$$[x_i - 1, x_j^{-1}] = -y_i y_j + y_i y_j^2 - y_i y_j^3 + \cdots. \quad (3.4)$$

Here we have used the fact that conjugation of $x_i - 1$ by x_j, x_j^{-1} in \hat{W} is just operation of x_j, x_j^{-1} on y_i in J . Since

$$\hat{W}_2 = [J, A] \quad \hat{W}_3 = [J, A, A], \text{ etc.} \quad (3.5)$$

We obtain from (3.3), (3.4) the following key result.

LEMMA 3.3. $\hat{W}_k = J \cdot IA^{k-1}$, $k \geq 2$. In particular, an element $v \in J$ is in \hat{W}_2 if and only if it can be written as a finite linear combination

$$v = \sum_{i=1}^n w_i y_i, \quad w_i \in J. \quad (3.6)$$

With this result it is now possible to settle various questions about our group \hat{F} . We first reprove a result of Bousfield–Kan [BK, p. 114].

THEOREM 3.4. Let $F = F(X)$ be a free group where X is countably infinite. Then $h_2: F_{ab} \rightarrow \hat{F}_{ab}$ is not surjective.

COROLLARY 3.5. Let $F = F(X)$ be a free group where X is countably infinite. Then \hat{F} is not parafree on X .

Proof. We enumerate the elements of X as follows

$$X = \{x_{11}, x_{21}, x_{22}, x_{31}, x_{32}, x_{33}, x_{41}, \dots\} \quad (3.7)$$

and consider (see [BK]) the element $\Lambda = (\lambda_1 F_2, \lambda_2 F_3, \dots) \in \hat{F}$ where

$$\lambda_1 = e, \lambda_k = [x_{21}, x_{22}][x_{31}, x_{32}, x_{33}] \cdots [x_{k1}, \dots, x_{kk}], k \geq 2. \quad (3.8)$$

We shall show that $\Lambda \notin \hat{F}_2$ but $\tau_2(\Lambda) = e \in F_{ab}$; hence τ_2 is not injective. This implies, by Lemma 1.1, that h_2 is not surjective.

Consider the free abelian group A on X and the group $W = IA \upharpoonright A$. Define a

map $f: F \rightarrow W$ by

$$f(x_{kj}) = \begin{cases} x_{kj} - 1 \in IA \subseteq W & \text{for } j = 1, \\ x_{kj} \in A \subseteq W & \text{for } 2 \leq j \leq k. \end{cases} \quad (3.9)$$

We then obtain a map $\hat{f}: \hat{F} \rightarrow \hat{W}$ with

$$\hat{f}(\Lambda) = (x_{21} - 1)(x_{22} - 1) + (x_{31} - 1)(x_{32} - 1)(x_{33} - 1) + \cdots \quad (3.10)$$

(see (3.3)). It is then clear from Lemma 3.3 that $\hat{f}(\Lambda) \notin \hat{W}_2$, so that $\Lambda \notin \hat{F}_2$. This completes the proof of Theorem 3.4.

We note that it might conceivably be the case that \hat{F} is parafree on some set other than X . That this is not the case follows from Corollary 3.9 where the existence of non-trivial divisible elements in \hat{F}_{ab} is proved. We first state

THEOREM 3.6. *Let $F = F(X)$ be a free group where X is countably infinite. Then \hat{F} contains a subgroup which is free on an uncountable set Y of elements which are linearly independent mod \hat{F}_2 .*

For the proof of this result we shall need the following

LEMMA 3.7. *There exists an uncountable set Σ of sequences $\sigma = (\sigma_0, \sigma_1, \dots)$ of natural numbers σ_i with the following properties:*

- (i) $\sigma_0 \geq 2$, $\sigma_{i+1} > \sigma_i$, $i \geq 0$;
- (ii) if $\{\sigma^{(1)}, \dots, \sigma^{(n)}\}$ is a finite subset of Σ then there exists $i \geq 0$ such that for every $k \geq i$ the entries $\sigma_k^{(1)}, \dots, \sigma_k^{(n)}$ are different.

Proof. Let Ω be an uncountable set of sequences $\omega = (\omega_0, \omega_1, \dots)$ of numbers $0, 1, 2, \dots, 9$ with $\omega_0 \geq 2$. Define, for any such ω , a sequence

$$\sigma(\omega) = (\sigma_0(\omega), \sigma_1(\omega), \dots) \quad (3.11)$$

by setting

$$\sigma_i(\omega) = \omega_0 \cdot 10^i + \omega_1 \cdot 10^{i-1} + \cdots + \omega_{i-1} \cdot 10^1 + \omega_i \cdot 10^0. \quad (3.12)$$

It is plain that $\omega \neq \omega'$ implies $\sigma(\omega) \neq \sigma(\omega')$. Also, it is clear that $\sigma_0(\omega) = \omega_0 \geq 2$ and that $\sigma_{i+1}(\omega) > \sigma_i(\omega)$, $i \geq 0$. Moreover, if $\{\sigma^{(1)}, \dots, \sigma^{(n)}\}$ is a finite subset of Σ with $\sigma^{(l)} = \sigma(\omega^{(l)})$, then there exists $i \geq 0$ such that for $k \geq i$ the elements $\sigma_k^{(1)}, \dots, \sigma_k^{(n)}$ are different. We may thus set $\Sigma = \{\sigma(\omega) \mid \omega \in \Omega\}$.

Proof (of Theorem 3.6). Let $F = F(X)$ where

$$X = \{x_{01}, x_{02}, \dots, x_{11}, x_{12}, \dots, x_{21}, x_{22}, \dots\} \quad (3.13)$$

Define for any sequence $\sigma = (\sigma_0, \sigma_1, \dots) \in \Sigma$ an element

$$\Lambda(\sigma) = (\lambda_1 F_2, \lambda_2 F_3, \dots) \in \hat{F} \quad (3.14)$$

by setting

$$\begin{cases} \lambda_i = e & \text{for } i < \sigma_0, \\ \lambda_i = [x_{01}, \dots, x_{0\sigma_0}][x_{11}, \dots, x_{1\sigma_1}] \cdots [x_{l1}, \dots, x_{l\sigma_l}] & \text{for } \sigma_l \leq i < \sigma_{l+1}. \end{cases} \quad (3.15)$$

Note that $\Lambda(\sigma)$ is an element of \hat{F} since σ is strictly increasing. Next we shall show that the elements $\Lambda(\sigma)$, $\sigma \in \Sigma$ generate a free subgroup of \hat{F} . For this it is enough to show that any *finite* set of elements

$$\Lambda^{(l)} = \Lambda(\sigma^{(l)}), \quad 1 \leq l \leq n$$

freely generate a free subgroup. By Lemma 3.7 we may conclude that there exists an $i \geq 0$ such that the entries $\sigma_i^{(1)}, \dots, \sigma_i^{(n)}$ are all different. We now consider the projection $p: F(X) \rightarrow F(x_{i1}, x_{i2}, x_{i3}, \dots)$; then p induces a map $\hat{p}: \hat{F}(X) \rightarrow \hat{F}(x_{i1}, x_{i2}, x_{i3}, \dots)$ with

$$\hat{p}(\Lambda(\sigma^{(l)})) = [x_{i1}, \dots, x_{i\sigma_i^{(l)}}] \in F(x_{i1}, x_{i2}, x_{i3}, \dots) \subseteq \hat{F}(x_{i1}, x_{i2}, x_{i3}, \dots) \quad (3.16)$$

It follows at once that $\hat{p}(\Lambda(\sigma^{(l)}))$ freely generate a free subgroup of $F(x_{i1}, x_{i2}, \dots)$. Hence the elements $\Lambda(\sigma^{(l)})$, $1 \leq l \leq n$ freely generate a free subgroup of $\hat{F}(X)$.

It remains to show that the elements $\Lambda(\sigma)$, $\sigma \in \Sigma$ are linearly independent mod \hat{F}_2 . For this we consider the group $W = IA \upharpoonright A$ where A is the free abelian group on X and the map $f: F \rightarrow W$ defined by

$$f(x_{ik}) = \begin{cases} (x_{i1} - 1) \in IA \subseteq W & \text{for } k = 1, \quad i \geq 0, \\ x_{ik} \in A \subseteq W & \text{for } k \geq 2, \quad i \geq 0. \end{cases} \quad (3.17)$$

For the induced map $\hat{f}: \hat{F} \rightarrow \hat{W}$ we then obtain

$$\hat{f}(\Lambda(\sigma)) = (x_{01} - 1) \cdots (x_{0\sigma_0} - 1) + (x_{11} - 1) \cdots (x_{1\sigma_1} - 1) + \cdots \quad (3.18)$$

It follows from Lemma 3.3 that $\hat{f}(\Lambda(\sigma)) \notin \hat{W}_2$. Moreover no non-trivial linear

combination of elements $\hat{f}(\Lambda(\sigma))$ lies in \hat{W}_2 . For, if $\sigma^{(1)}, \dots, \sigma^{(n)}$ are different elements of Σ then, by Lemma 3.7, there exists $i \geq 0$ such that for all $k \geq i$ the entries $\sigma_k^{(1)}, \dots, \sigma_k^{(n)}$ are different. Hence a nontrivial linear combination of $\hat{f}(\Lambda(\sigma^{(1)})), \dots, \hat{f}(\Lambda(\sigma^{(n)}))$ cannot lie in \hat{W}_2 .

We note that in the course of the above proof we also have proved the following result

COROLLARY 3.8. *Let $F = F(X)$ be a free group where X is countably infinite. Then \hat{F}_{ab} is uncountably infinite.*

We next prove

THEOREM 3.9. *Let $F = F(X)$ be a free group where X is countably infinite. Then \hat{F}_{ab} contains uncountably many linearly independent divisible elements.*

Proof. Let $X = \{x_{01}, x_{02}, \dots, x_{11}, x_{12}, \dots, x_{21}, x_{22}, \dots\}$ and consider for $\sigma \in \Sigma$ (Lemma 3.7) the element

$$\Gamma(\sigma) = (\gamma_1 F_2, \gamma_2 F_3, \dots) \in \hat{F} \quad (3.19)$$

defined by

$$\begin{cases} \gamma_i = e & \text{for } i < \sigma_2, \\ \gamma_i = ([x_{21}, \dots, x_{2\sigma_2}][x_{31}, \dots, x_{3\sigma_3}] \cdots [x_{l1}, \dots, x_{l\sigma_l}]^{l \cdots})^3)^2 & \text{for } \sigma_l \leq i < \sigma_{l+1}. \end{cases} \quad (3.20)$$

(The fact that we start with σ_2 is merely a notational convenience.) We shall show that each $\Gamma(\sigma)$ gives rise to a divisible element in \hat{F}_{ab} . We first recall that F_{ab} is a direct summand of \hat{F}_{ab} . Thus in order to exhibit divisible elements in \hat{F}_{ab} we may consider the quotient of \hat{F}_{ab} by F_{ab} , in other words we may consider the quotient of \hat{F} by the normal subgroup N generated by F and \hat{F}_2 . In order to show that for $k \geq 2$ the element $\Gamma(\sigma)$ is a k -th power modulo N we consider the element

$$\Delta = \Delta(\sigma) = (\delta_1 F_2, \delta_2 F_3, \dots) \in \hat{F} \quad (3.21)$$

where

$$\begin{cases} \delta_i = e & \text{for } i < \sigma_k \\ \delta_i = ([x_{k1}, \dots, x_{k\sigma_k}][x_{l1}, \dots, x_{l\sigma_l}]^{l \cdots})^{k+1})^{k!} & \text{for } \sigma_i \leq i < \sigma_{l+1}, l \geq k. \end{cases} \quad (3.22)$$

Clearly, modulo N the elements $\Gamma(\sigma)$ and $\Delta(\sigma)$ are equivalent. But $\Delta(\sigma)$ is a k -th power. Hence $\Gamma(\sigma)$ is a k -th power modulo N . An argument similar to the one used in the proof of Theorem 3.6 shows that the elements $\Gamma(\sigma)$, $\sigma \in \Sigma$ are linearly independent in \hat{F}_{ab} .

COROLLARY 3.10. *Let $F = F(X)$ be a free group, where X is countably infinite. Then \hat{F}_k/\hat{F}_{k+1} , $k \geq 1$ contains non-trivial divisible elements.*

Proof. Let $X = \{x_{01}, x_{02}, \dots, x_{11}, x_{12}, \dots, x_{21}, x_{22}, \dots\}$ and let $\Gamma = \Gamma(\sigma) \in \hat{F}$ be one of the elements defined in (3.19), (3.20). We consider the element

$$\Gamma^* = [\Gamma, x_{01}, x_{02}, \dots, x_{0,k-1}] \in \hat{F}_k. \quad (3.23)$$

Since the k -fold commutator is a linear map from the k -fold tensor product of \hat{F}_{ab} into \hat{F}_k/\hat{F}_{k+1}

$$[\cdot \cdots \cdot]: \hat{F}_{ab} \otimes \cdots \otimes \hat{F}_{ab} \rightarrow \hat{F}_k/\hat{F}_{k+1} \quad (3.24)$$

the element Γ^* gives certainly rise to a divisible element in \hat{F}_k/\hat{F}_{k+1} . It remains to prove that Γ^* is non-trivial. We use the map $f: F \rightarrow W$ defined by

$$\begin{cases} f(x_{0l}) = x_{0l} \in A \subseteq W, & l \geq 1. \\ f(x_{il}) = x_{il} \in A \subseteq W, & i \geq 1, \quad l \geq 2. \\ f(x_{i1}) = (x_{i1} - 1) \in IA \subseteq W, & i \geq 1. \end{cases} \quad (3.25)$$

For the induced map $\hat{f}: \hat{F} \rightarrow \hat{W}$ we obtain

$$\begin{aligned} \hat{f}(\Gamma^*) = \\ [2(x_{21} - 1) \cdots (x_{2\sigma_2} - 1) + 3!(x_{31} - 1) \cdots (x_{3\sigma_3} - 1) + \cdots](x_{01} - 1) \cdots (x_{0,k-1} - 1) \end{aligned} \quad (3.26)$$

By Lemma 3.3, $\hat{f}(\Gamma^*) \notin \hat{W}_{k+1}$ and hence $\Gamma^* \notin \hat{F}_{k+1}$.

The following result should be compared with Theorem 4.2 of [B2].

THEOREM 3.11. *The 2-generator subgroups of \hat{F} are free.*

Proof. Let $a, b \in \hat{F} = \hat{F}(X)$. We have to consider two cases.

(i) Let $[a, b] \neq e$. In this case we shall deduce our result from Theorem 4.2 of [B2]. Clearly there exists $i \geq 1$ such that for $\tau_i: \hat{F} \rightarrow F/F_i$ we have $\tau_i[a, b] \neq e$. Consider then a finite subset $Y \subseteq X$ and the projection $p: F(X) \rightarrow F(Y)$, such that for $p\tau_i: \hat{F}(X) \rightarrow F(Y)/F_i(Y)$ we have $[p\tau_i a, p\tau_i b] \neq e$. We may thus consider $\hat{p}: \hat{F}(X) \rightarrow \hat{F}(Y)$. Since $\hat{F}(Y)$ is parafree and $[pa, pb] \neq e$ we may conclude from

Theorem 4.2 of [B2] that $\hat{p}a, \hat{p}b$ generate a free subgroup. Hence a, b generate a free subgroup in $\hat{F}(X)$.

(ii) Let $[a, b] = e$. We shall proceed as in the proof of Theorem 4.2 of [B2]. By Theorem 3.1 of [B2] the group $\hat{F} = \hat{F}(X)$ can be embedded in the power series ring $\mathbf{Z}[[X]]$ and hence in $\mathbf{Q}[[X]]$. Under that embedding let

$$\begin{cases} a = 1 + a_i + \cdots, & a_i \neq 0 \\ b = 1 + b_j + \cdots, & b_j \neq 0 \end{cases} \quad (3.27)$$

i.e. a_i, b_j are the first non-zero terms in the power series corresponding to a, b . The elements a_i, b_j are Lie elements in $\mathcal{L}[X]$, the Lie algebra over \mathbf{Q} generated by X in $\mathbf{Q}[X]$. Since $\mathcal{L}[X]$ is a free Lie algebra over \mathbf{Q} it follows that a_i, b_j generate a free sub- \mathbf{Q} -Lie algebra (see Sirsov [S], Witt [W]). Since $[a, b] = e$ we have in $\mathbf{Q}[[X]]$

$$ab = 1 + a_i + b_j + a_i b_j + \cdots = 1 + a_i + b_j + b_j a_i + \cdots = ba. \quad (3.28)$$

Hence in $\mathcal{L}[X]$ we have $[a_i, b_j] = 0$, so that the Lie algebra generated by a_i, b_j is abelian. Since it is free it must be isomorphic to \mathbf{Q} . It follows that there exist integers $m, n > 0$ such that

$$ma_i = nb_j. \quad (3.29)$$

In particular $i = \deg a_i = \deg b_j = j$. We now compute $c = a^{-m} \cdot b^n$ in $\mathbf{Z}[[X]]$.

$$\begin{aligned} c = a^{-m} b^n &= (1 - ma_i + \cdots)(1 + nb_j + \cdots) \\ &= 1 - ma_i + nb_j + \{\text{terms of degree} \geq i + 1\} \\ &= 1 + \{\text{terms of degree} \geq i + 1\}. \end{aligned} \quad (3.30)$$

But c , being an element in the subgroup generated by a, b commutes with a and b and hence satisfies $[a, c] = e$. A repetition of the above argument with c at the place of b shows that the power series expansion of c has the form

$$c = 1 + c_k + \cdots, \quad c_k \neq 0 \quad (3.31)$$

with $k = i$. This is a contradiction to (3.30) so that $c = 1$, i.e. $a^m = b^n$. Since \hat{F} is torsion-free it follows that the subgroup generated by a, b is infinite cyclic, hence free.

4. Two subgroups of \hat{F}

Let $F = F(X)$ be the free group on the set X . Here we shall exhibit two subgroups \bar{F}, \tilde{F} of $\hat{F} = \hat{F}(X)$ which are parafree on X . Note that \hat{F} is not itself parafree on X if X is at least countably infinite.

THEOREM 4.1. *There exists a subgroup $\bar{F} \subseteq \hat{F} = \hat{F}(X)$ which is parafree on X and has the property that it contains all subgroups of \hat{F} which are parafree on X .*

Proof. We first show that the system of subgroups of \hat{F} which are parafree on X is directed. Thus let U, V be two subgroups of \hat{F} which are parafree on X . Let W be the subgroup of \hat{F} generated by U, V . We claim that W is parafree on X . Since $F \subseteq W \subseteq \hat{F}$ we may apply Lemma 1.1, so that we have to show that $F_{ab} \rightarrow W_{ab}$ is surjective. Consider the obvious epimorphism $U *_F V \twoheadrightarrow W$. We then have the series of maps

$$F_{ab} \twoheadrightarrow U_{ab} \rightarrow (U *_F V)_{ab} \twoheadrightarrow W_{ab} \quad (4.1)$$

so that we only have to show that $U_{ab} \rightarrow (U *_F V)_{ab}$ is surjective. But this is trivial since U, V are parafree on X . We may thus define \bar{F} by $\bar{F} = \varinjlim U$, where U is a subgroup of \hat{F} which is parafree on X . It is then clear that $F_{ab} \rightarrow \bar{F}_{ab}$ is surjective, so that \bar{F} is parafree on X , by Lemma 1.1.

It is obvious that the above construction is independent of the fact that the group we start with is free. Thus if G is an arbitrary group we may find in \hat{G} a group \bar{G} with $hG \subseteq \bar{G} \subseteq \hat{G}$ such that h induces isomorphisms $h_i : G/G_i \xrightarrow{\sim} \bar{G}/\bar{G}_i$, $i \geq 2$, and with the following universal property. If $f : G \rightarrow H$ is a homomorphism such that $f_i : G/G_i \xrightarrow{\sim} H/H_i$, $i \geq 2$ and H is residually nilpotent, then there exists a unique $\bar{f} : H \rightarrow \bar{G}$ such that $\bar{f} \circ f = h : G \rightarrow \bar{G}$. By construction of \bar{G} we have

$$\text{COROLLARY 4.2. } \bar{\bar{G}} = \bar{G}.$$

As in Corollary 2.6 we obtain from Corollary 4.2

COROLLARY 4.3. *Let F be free, then \bar{F} is freely indecomposable.*

We shall now construct another subgroup \tilde{F} of $\hat{F} = \hat{F}(X)$ which is parafree on X . Thus we have $\tilde{F} \subseteq \bar{F}$, by Theorem 4.1, but we shall later show that $\tilde{F} \neq \bar{F}$ if X is (at least) countably infinite.

Consider the directed system of finite subsets Y of X and the associated

directed system of groups $\hat{F}(Y)$. Define

$$F = \varinjlim \hat{F}(Y), \quad Y \subseteq X, Y \text{ finite} \quad (4.2)$$

If we consider $\hat{F}(X)$ as a subgroup of the power series ring $\mathbb{Z}[[X]]$ (see Theorem 3.1 of [B2]), then \tilde{F} may be described as the subgroup of those power series of \hat{F} which involve only finitely many elements of X .

PROPOSITION 4.4. *The group \tilde{F} is parafree on X .*

Proof. Clearly $F \subseteq \tilde{F} \subseteq \hat{F}$. Since $\tilde{F}_{ab} = \varinjlim (\hat{F}(Y))_{ab} = \varinjlim (F(Y))_{ab}$ is free abelian on X , the map $F_{ab} \rightarrow \tilde{F}_{ab}$ is surjective. By Lemma 1.1 we conclude that \tilde{F} is parafree on X .

PROPOSITION 4.5. *Let $F = F(X)$ where X is countably infinite. Then \tilde{F} is a proper subgroup of \bar{F} .*

Proof. We shall exhibit a subgroup U of \hat{F} which is parafree on X , but not contained in \tilde{F} . Since $U \subseteq \bar{F}$ by Theorem 4.1 it then follows that $\tilde{F} \neq \bar{F}$.

Let $X = \{x_1, x_2, \dots, y_1, y_2, \dots\}$. Define elements

$$Z^{(i)} = (\zeta_1^{(i)} F_2, \zeta_2^{(i)} F_3, \dots) \in \hat{F}, \quad i = 0, 1, \dots \quad (4.3)$$

by setting

$$\begin{cases} \zeta_1^{(i)} = x_i, \\ \zeta_2^{(i)} = [x_{i+1}, y_{i+1}]x_i, \\ \zeta_3^{(i)} = [[x_{i+2}, y_{i+2}]x_{i+1}, y_{i+1}]x_i, \text{ etc.} \end{cases} \quad (4.4)$$

where $x_0 = e$. Modulo any F_k and hence in \hat{F} we have

$$Z^{(i)} = [Z^{(i+1)}, y_{i+1}]x_i, \quad i = 0, 1, \dots \quad (4.5)$$

Consider now the subgroup U of \hat{F} generated by $x_1, x_2, \dots, y_1, y_2, \dots, Z^{(0)}, Z^{(1)}, Z^{(2)}, \dots$. We claim that U is parafree on X . By Lemma 1.1 we only have to show that $F_{ab} \rightarrow U_{ab}$ is surjective. But it is clear that

$$\begin{cases} Z^{(0)} \equiv e \pmod{U_2}, \\ Z^{(i)} \equiv x_i \pmod{U_2}, \quad i \geq 1. \end{cases} \quad (4.6)$$

Finally it is plain that none of the elements $Z^{(i)}$ is contained in \tilde{F} .

BIBLIOGRAPHY

- [BG] F. BACHMANN, L. GRÜNENFELDER, *Ueber Lie-Ringe von Gruppen und ihre universelle Enveloppen*, Comment. Math. Helv. 47, (1972), 332–340.
- [B1] G. BAUMSLAG, *Groups with the same lower central sequence as a relatively free group. I. The groups*, Trans. Amer. Math. Soc. 129 (1967), 308–321.
- [B2] —, *Groups with the same lower central sequence as a relatively free group. II. Properties*, Trans. Amer. Math. Soc. 142 (1969), 507–538.
- [BK] A. K. BOUSFIELD, D. M. KAN, *Homotopy Limits, Completions and Localizations*. Lecture Notes in Mathematics, vol. 304; Springer 1972.
- [S] A. SIRSOV, *Unteralgebren von freien Liealgebren (russisch)*, Math. Sbornik N.S. 33 (75), (1953), 441–452.
- [W] E. WITT, *Die Unterringe der freien Lieschen Ringe*, Math. Z. 64 (1956), 195–216.

CCNY

Convent Ave. and 138th Street
New York, N.Y. 10031

ETH

Rämistrasse 101
8092 Zürich

Received July 3, 1976

