Semi-continuity of the face-function for a convex set.

Autor(en): **Eifler, Larry Q.**

Objekttyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **52 (1977)**

PDF erstellt am: **28.04.2024**

Persistenter Link: https://doi.org/10.5169/seals-40002

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.
Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.
Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*
ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

http://www.e-periodica.ch

Comment. Math. Helvetici **52** (1977) 325-328 Birkhauser Verlag, Basel

Semi-continuity of the face-function for a convex set

LARRY Q. EIFLER

Introduction

Throughout this paper, let K be a compact convex subset of a locally convex topological vector space E. Given $x \in K$, set $F(x) = \text{cl}\{y \in K : [y, x + \varepsilon(x-y)] \subseteq K$ for some $\varepsilon > 0\}$. We call F the face-function on K since $F(x)$ is the smallest closed face of K containing x if $F(x)$ is finite dimensional. If $f : K \to \mathbf{R}$ is continuous, we define the lower envelope f_e of f by $f_e = \sup\{g : g$ is a continuous affine function on K satisfying $g \leq f\}$. Following Klee and Martin [3], set $K_e = \{x \in K : f_e$ is continuous at x for each continuous function $f : K \to \mathbf{R}\}$ and set $K_l = \{x \in K : F$ is lower semi-continuous at $x\}$. Klee and Martin proved that $K_e \subseteq K_l$ in general and that $K = K_e = K_l$ if K is 2-dimensional. They left open whether $K_e = K_l$. We show that $K_e = K_l$ is K is finite dimensional and produce an infinite dimensional example where $K = K_l \neq K_e$.

Lower semi-continuity of F

Let $x \to F(x)$ be the face-function on K defined above. We say that F is lower semi-continuous at x if for each $y \in F(x)$ and for each neighborhood U of y, $\{z \in K : F(z)$ meets $U\}$ is a neighborhood of x. Note that $F(x)$ is a compact convex subset K for each $x \in K$. If F is lower semi-continuous on K, then the set of extreme points $ex(K)$ of K is closed. If $ex(K)$ is closed and if K is 2 or 3-dimensional, then Clausing and Magerl [1] have shown that $K_e = K$ and so $K_l = K$.

Let $P(K)$ denote the space of Radon probability measures on K and equip $P(K)$ with the weak* topology. Given $\mu \in P(K)$, there exists a unique point $r(\mu)$ in K such that $\int g \, d\mu = g(r(\mu))$ for each continuous affine function g on K. The map $r : P(K) \to K$ is the resultant or barycentric map. If $x \in K$, we let δ_x denote the point mass measure at x. The map r is an open map of $P(K)$ onto K if and only if $K = K_e$. See [2 or 5]. We say that r is open at $\mu \in P(K)$ if for each meighborhood U of μ in $P(K)$, $r(U) = \{r(\nu) : \nu \in U\}$ is a neighborhood of $r(\mu)$. We say that r is λ-open at $\mu \in P(K)$ where $0 < \lambda < 1$ if for each neighborhood U of μ

in $P(K)$, $\lambda r(U)+(1-\lambda)K$ is a neighborhood of $r(\mu)$. We first establish criteria for determining when r is open at μ. These results are of interest aside from their application to the study of the lower semi-continuity of the face-function.

LEMMA 1. *Let $\mu \in P(K)$. Then r is open at μ if r is λ-open at μ for some $0<\lambda<1$.*

Proof. Assume that r is λ-open at μ. Set $x=r(\mu)$. If $x_\alpha \to x$, then there exist $\mu_\alpha \to \mu$ and $y_\alpha \in K$ such that $\lambda r(\mu_\alpha)+(1-\lambda)y_\alpha = x_\alpha$. But $y_\alpha \to x$. Hence, there exist $\nu_\alpha \to \mu$ and $z_\alpha \in K$ such that $\lambda r(\nu_\alpha)+(1-\lambda)z_\alpha = y_\alpha$. Thus,

$$x_\alpha = \lambda(2-\lambda)\left\{r\left(\frac{\mu_\alpha}{2-\lambda}+\frac{(1-\lambda)\nu_\alpha}{2-\lambda}\right)\right\}+(1-\lambda)^2 z_\alpha.$$

One obtains that r is $\lambda(2-\lambda)$-open at μ. Hence, r is ρ-open at μ for each $0<\rho<1$. This implies that r is open at μ.

LEMMA 2. *Let $x \in K$. The following are equivalent.*

(1) *r is open at μ if $r(\mu)=x$ and*
(2) *r is open at μ if $r(\mu)=x$ and if μ is supported by 2 points.*

Proof. We only need to show $(2 \Rightarrow 1)$. Let U be a neighborhood of μ where $r(\mu)=x$ and μ is supported by n points. We show that $r(U)$ is a neighborhood of x by induction on n. The result holds for $n=2$. So assume the result holds for $n \le m$. Fix $\mu \in P(K)$ such that $r(\mu)=x$ and μ is supported by $\{x_1, \ldots, x_{m+1}\}$. Let $x=\sum_{i=1}^{m+1} \lambda_i x_i$. We assume each $\lambda_i > 0$. Set $y_k = (\lambda_k x_k + \lambda_{k+1}x_{k+1})/(\lambda_k+\lambda_{k+1})$ and set $\mu_k = \sum_{i=1}^{k-1} \lambda_i \delta_{x_i} + (\lambda_k+\lambda_{k+1})\delta_{y_k} + \sum_{i>k}^{m+1} \lambda_i \delta_{x_i}$. Suppose $x_\alpha \to x$. Then there exist $\mu_k^\alpha \to \mu_k$ such that $r(\mu_k^\alpha)=x_\alpha$. Set $\nu_\alpha = \sum_{k=1}^{m+1}(1/m+1)\mu_k^\alpha$. Then $r(\nu_\alpha)=x_\alpha$. Since $\nu_\alpha \to \sum_{k=1}^{m+1}(1/m+1)\mu_k$, we have $\lim_\alpha \sup \nu_\alpha(V) \ge (m/m+1)\lambda_k$ if V is an open set containing x_k. Thus, there exist $\mu_\alpha \to \mu$ and $z_\alpha \in K$ such that $(m/m+1)r(\mu_\alpha)+(1/m+1)z_\alpha = x_\alpha$. Hence, r is open at μ by Lemma 1. By approximating measures by measures with finite support, we see that r is open at μ if $r(\mu)=x$.

THEOREM. *Let $x \in K$. The following are equivalent.*

(1) *f_e is continuous at x for each $f \in C_{\mathbf{R}}(K)$*
(2) *r is open at μ if $r(\mu)=x$*
(3) *r is open at μ if $r(\mu)=x$ and if μ is supported by 2 points.*

Proof. The implication (1) $\Rightarrow$ (2) follows from Proposition 3.1 in Phelps [4, p.

21]. The implication (2) $\Rightarrow$ (1) follows from the separation form of the Hahn–Banach theorem and taking limits in the hyperspace of $P(K)$. See [2] for details. The implication (2 $\Leftrightarrow$ 3) is simply Lemma 2.

COROLLARY. *Assume K is finite dimensional. Then $K_e = K_l$.*

Proof. The inclusion $K_e \subseteq K_l$ was established in [3]. Let $x \in K_l$. Suppose $\mu \in P(K)$ such that μ is supported by $\{y, z\}$ and $r(\mu) = x$. We only need to show that r is $\frac{1}{2}$-open at μ by Lemma 1 and the above theorem. Set $x = \lambda y + (1-\lambda)z$ where $0 \leq \lambda \leq 1$. We may assume $y \neq z$ and $0 < \lambda < 1$. Let U be a neighborhood of μ in $P(K)$. Set $\Omega = \frac{1}{2}r(U) + \frac{1}{2}K$. Assume Ω is not a neighborhood of x. Then there exist $x_n \to x$ such that $x_n \notin \Omega$ and $\dim F(x_n) = q$ where q is least possible. By taking subsequences, we may assume that there exist $y_n, z_n \in F(x_n)$ such that $y_n \to y$ and $z_n \to z$ since $\limsup F(x_n) \supseteq F(x) \supseteq \{y, z\}$. Set $w_n = \lambda y_n + (1-\lambda)z_n$. We may assume $w_n \neq x_n$. Set $\varepsilon_n = \max\{\varepsilon : x_n + \varepsilon(w_n - x_n) \in K\}$. Then $\dim F(x_n + \varepsilon_n(w_n - x_n)) < \dim F(x_n) = q$. But, $w_n \in r(U)$ for n large. If $\varepsilon_n \geq 1$ and if $w_n \in r(U)$, then

$$\frac{\varepsilon_n}{1+\varepsilon_n} w_n + \frac{1}{1+\varepsilon_n}\{x_n + \varepsilon_n(x_n - w_n)\} = x_n \in \Omega.$$

Hence, $\varepsilon_n < 1$ for n large. Thus, $x_n + \varepsilon_n(w_n - x_n) \to x$ and $x_n + \varepsilon_n(w_n - x_n) \in \Omega$ which is impossible by the minimality of q.

Example 1. Let K be the convex hull of $\{(e^{i\theta}, \pm 1) : 0 \leq \theta \leq \pi\} \cup \{(1, \pm i)\}$ in $\mathbf{C}^2$. Then K is 4-dimensional and $ex(K)$ is closed. The face-function is not lower semi-continuous at $(1, 0)$ since $F(1, 0)$ is a square and $F(e^{i\theta}, 0)$ is an interval if $0 < \theta \leq \pi$.

Example 2. Let $X = \{0, 1, \frac{1}{2}, \frac{1}{3}, \ldots\}$ and equip X with the usual metric from $\mathbf{R}$. Let K denote the closed unit ball in the space of real Radon measures on X, i.e., the dual of $C_{\mathbf{R}}(X)$. Equip K with the weak* topology. Then K is a compact convex set. Given $\mu \in K$, set $\|\mu\| = \mu^+(X) + \mu^-(X)$. If $\|\mu\| < 1$, then $F(\mu) = K$. If $\|\mu\| = 1$, then $F(\mu)$ is the closed convex hull of $\{sgn\,(\mu(x)) \cdot \delta_x : x \in X\}$. One easily checks that the face-function is lower semi-continuous on K and so $K = K_l$. The zero measure 0 is not in K_e by criterion (2) in the theorem since $\frac{1}{2}[\delta_1 + (-1)\delta_1] = 0$ and $\frac{1}{2}[\delta_{1/n} + (-1)\delta_{1/n+1}] \to 0$.

REFERENCES

[1] A. CLAUSING and G. MAGERL, *Generalized Dirichlet problems and continuous selections of representing measures*, Math. Ann. *216* (1975), 71–78.

[2] L. Q. EIFLER, *Openness of convex averaging*, Glasnik Matematicki (to appear).
[3] V. KLEE and M. MARTIN, *Semicontinuity of the face-function of a convex set*, Comment. Math, Helvetici *46* (1971), 1–12.
[4] R. R. PHELPS, *Lectures on Choquet's theorem*, Van Nostrand, Princeton, 1966.
[5] J. VESTERSTROM, *On open maps, compact convex sets, and operator algebras*, J. London Math. Soc. (2), *6* (1973), 289–297.

Received October 12, 1976