

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 52 (1977)

Artikel: Flat manifolds with non-zero Euler characteristics.
Autor: Smillie, John
DOI: <https://doi.org/10.5169/seals-40008>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Flat manifolds with non-zero Euler characteristics

By JOHN SMILLIE

A flat structure on a smooth vector bundle ξ can be given either by a connection on ξ with zero curvature or by a reduction of the structure group of ξ from $Gl(\mathbb{R}^n)$ to $Gl(\mathbb{R}^n)$ with the discrete topology. Using either definition we see that if ξ and ξ' admit flat structures and f is a map then $\xi + \xi'$ and $f^*\xi$ admit flat structures. A flat structure on a manifold M is a flat structure on $T(M)$. We say a manifold is flat if it admits a flat structure.

Let M_g denote the surface of genus g for $g \geq 1$. Milnor (1) proves that an oriented bundle ξ over M_g admits a flat structure if and only if

$$|e(\xi)[M_g]| \leq -\frac{1}{2}\chi(M_g) \quad (1)$$

where $e(\xi)$ is the Euler class of ξ and $\chi(M)$ is the Euler characteristic of M . If M_g is a flat manifold this formula implies $\chi(M_g) = 0$. According to Hirsch-Thurston (2) it is an old conjecture that the Euler characteristic of any flat manifold is zero. The question is raised by Milnor (1) and Kamber-Tondeur (3, p. 47). In this paper we prove the following.

THEOREM. *There are flat manifolds M^{2n} with non-zero Euler characteristic for all $n > 1$.*

Starting with surfaces we construct manifolds M^{2n} by taking products and connected sums. To prove the existence of a flat structure on $T(M^{2n})$ we establish that $T(M^{2n})$ is isomorphic to a second bundle ζ . ζ admits a flat structure because it is the sum of pullbacks of 2-plane bundles which admit flat structures. A bundle isomorphism allows us to transfer the flat structure from ζ to $T(M)$.

ALMOST TRIVIAL BUNDLES. We will call a bundle ξ^n almost trivial if $\xi^n + \varepsilon^1 = \varepsilon^{n+1}$ where ε^n denotes the trivial bundle of dimension n .* Sums and pullbacks of almost trivial bundles are almost trivial. If $T(M)$ is almost trivial then M is said to be almost parallelizable.


* If the base space of ξ^n is n -dimensional then ξ^n is almost trivial if and only if ξ^n is stably trivial.

PROPOSITION 2. *An oriented 2-plane bundle ξ over a surface M_g is almost trivial if $e(\xi)[M_g]$ is even.*

Proof. $\xi + \varepsilon^1$ admits a two-frame over M_g if a certain obstruction class $\mathcal{O}_2(\xi + \varepsilon^1)$ vanishes. \mathcal{O}_2 can be identified with $w_2(\xi + \varepsilon^1) = w_2(\xi)$ which vanishes when $e(\xi)[M_g]$ is even. Assuming $\mathcal{O}_2 = 0$, $\xi + \varepsilon^1 = \eta + \varepsilon^2$ but $w_1(\eta) = w_1(\xi) = 0$ so $\eta = \varepsilon^1$.

PROPOSITION 3. *If n dimensional manifolds N and M are almost parallelizable then $N \# M$ is almost parallelizable.*

Proof. We can immerse N and M in \mathbb{R}^{n+1} . We can connect them by a tube. Smoothing the corners we see that we have constructed an immersion of $N \# M$ in \mathbb{R}^{n+1} hence $N \# M$ is almost parallelizable.



LEMMA. *If ξ is an almost trivial bundle over M then there is a map $g : M \rightarrow S^n$ so that $\xi = g^* T(S^n)$.*

Proof. Choose a bundle map $f : \xi + \varepsilon^1 \rightarrow \mathbb{R}^{n+1}$. We have a non-zero section $s : M \rightarrow \xi + \varepsilon^1$ associated to ε^1 . Using a Gram-Schmidt procedure we may assume that $fs(x)$ is of unit length and perpendicular to $f(\xi_x)$.

$$\begin{array}{ccc} \xi & \xrightarrow{f_{\ast} p \times f|_{\xi}} & S^n \times \mathbb{R}^{n+1} \\ \downarrow p & & \downarrow \pi \\ M & \xrightarrow{fs} & S^n \end{array}$$

Set $g = fs$. ξ is the pullback by g of the sub-bundle $T(S^n) \subset S^n \times \mathbb{R}^{n+1}$ consisting of pairs (x, y) where y is perpendicular to x .

PROPOSITION 4. *Let η_1 and η_2 be almost trivial oriented $2n$ -plane bundles over an oriented $2n$ -manifold. If $e(\eta_1) = e(\eta_2)$ then η_1 and η_2 are isomorphic.*

Proof. The lemma yields maps g_i such that $\eta_i = g_i^* T(S^{2n})$. $e(\eta_i) = 2g_i^* u$ where $u \in H^{2n}(S^{2n}; \mathbb{Z})$ is dual to the orientation class. The homotopy class of $g_i : M^{2n} \rightarrow S^{2n}$ is determined by the cohomology class $g_i^* u$ hence g_1 and g_2 are homotopic. The isomorphism type of $g_i^* T(S^{2n})$ depends only on the homotopy type of g_i hence η_1 and η_2 are isomorphic.

PROOF OF THEOREM. We construct a flat 4-manifold N with $\chi(N)=4$ and a flat 6-manifold Q with $\chi(Q)=8$. Products of these two manifolds with themselves give examples of flat manifolds with non-zero Euler characteristics of any even dimension greater than 2.

Let P be a parallelizable 4-manifold such as $S^1 \times S^3$. Let $N = (M_3 \times M_3) \# \underbrace{P \# \cdots \# P}_{6\text{-copies}}$ where M_3 is the surface of genus 3. Clearly $M_3 \times M_3$ is almost parallelizable. N is almost parallelizable by proposition 3. We have the formulas

$$\begin{aligned}\chi(M_g) &= 2 - 2g \\ \chi(M^{2n} \# N^{2n}) &= \chi(M^{2n}) + \chi(N^{2n}) - 2.\end{aligned}$$

We calculate that $\chi(N)=4$.

Let ξ be the oriented bundle over M_3 with $e(\xi)[M_3]=2$. Proposition 2 implies that ξ is almost trivial. $\xi \times \xi$ is a bundle over $M_3 \times M_3$. Let $f: N \rightarrow M_3$ be a degree one map that, for example, sends the six P summands to a point. $f^*\xi \times \xi$ is almost trivial and $e(f^*\xi \times \xi)[N]=4$. $T(N)$ is almost trivial and $e(T(N))[N]=\chi(N)=4$. It follows from proposition 4 that $T(N)$ and $f^*\xi \times \xi$ are isomorphic. Milnor's formula implies that ξ admits a flat structure hence $f^*\xi \times \xi$ admits a flat structure. Choosing an isomorphism we transfer the flat structure from $f^*\xi \times \xi$ to $T(N)$.

Let Q be the 6-manifold $\{(M_3 \times M_3) \# \underbrace{P \# \cdots \# P}_{9\text{-copies}}\} \times M_3$. We calculate that $\chi(Q)=8$. We have a flat bundle $\eta=\{h^*(\xi \times \xi)\} \times \xi$ over Q where h is the appropriate degree one map. $e(\eta)[Q]=8$. $T(Q)$ and η are almost trivial and have the same Euler class. By proposition 4 they are isomorphic hence $T(Q)$ admits a flat structure.

REFERENCES

- [1] J. MILNOR, *On the existence of a connection with curvature zero*, Comm. Math. Helv. 32 (1957), 215–223.
- [2] M. HIRSCH and W. THURSTON, *Foliated bundles, invariant measures and flat manifolds*, Annals of Math. 101 (1975), 369–390.
- [3] F. KAMBER and P. TONDEUR, *Flat manifolds*, Lecture notes in mathematics 67, Springer-Verlag, Heidelberg, 1968.

*Dept. of Mathematics
University of Chicago,
Chicago, Ill. 60637 U.S.A.*

Received November 1976

