

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 52 (1977)

Artikel: A remark on Abel's Theorem and the mapping of linear series.
Autor: Martens, Henrik H.
DOI: <https://doi.org/10.5169/seals-40019>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A remark on Abel's Theorem and the mapping of linear series

HENRIK H. MARTENS

Let X and Y be closed Riemann surfaces of positive genera, and let $\phi : X \rightarrow Y$ be a holomorphic map onto.

Remark. If $D = \sum m_i Q_i$ is a positive divisor of degree n and (projective) dimension r on X , then $\phi(D) = \sum m_i \phi(Q_i)$ is a positive divisor of degree n and (projective) dimension $\geq r$ on Y . If the complete linear series determined by D is without fixed points, then so is that determined by $\phi(D)$.

Proof. An arbitrary positive divisor of degree r on Y may be written as $\phi(D')$ where D' is a positive divisor of degree r on X . If D is of dimension r , there is a positive divisor D'' on X such that $D \sim D' + D''$ (linear equivalence). Connecting the points of D and $D' + D''$ by curves and applying Abel's theorem to the pullbacks to X of the holomorphic differentials on Y , we immediately see that $\phi(D) \sim \phi(D') + \phi(D'')$ on Y . Since $\phi(D')$ was an arbitrary positive divisor of degree r on Y , we conclude that $\phi(D)$ is of dimension $\geq r$.

The preimage under ϕ of the set of points occurring in $\phi(D)$ is a finite set of points on X . If D determines a complete linear series without fixed points, then we can find a positive divisor D' on X such that $D' \sim D$, and no point in the preimage occurs in D' . Then, since $\phi(D) \sim \phi(D')$ on Y , the complete linear series determined by $\phi(D)$ must be without fixed points.

COROLLARY 1. *If X can be displayed as an n -sheeted covering of the Riemann sphere, then so can Y . In particular, if X is hyperelliptic then Y is hyperelliptic or elliptic.*

Proof. X can be displayed as an n -sheeted covering of the Riemann Sphere if and only if it admits a positive divisor of degree n and dimension ≥ 1 that determines a complete linear series without fixed points.

COROLLARY 2. *If D and $\phi(D)$ both are divisors of dimension 1, then the branch points of the coverings of the Riemann sphere determined by D are mapped on the branch points of the coverings determined by $\phi(D')$ without reduction of branching order.*

Proof. Under the stated assumption, the branch points of any cover are precisely those that appear with multiplicity ≥ 2 in the divisors of the linear series.

APPLICATION. We consider the case when X is hyperelliptic. Then Y is hyperelliptic or elliptic, and in any case a display of Y as a 2-sheeted cover of the Riemann sphere must have $2h + 2$ branch points, where h is the genus of Y . If g is the genus of X , the $2g + 2$ branch points of X must map on the branch points of Y , whence $2g + 2 \leq m(2h + 2)$ or

$$g \leq m(h + 1) - 1$$

where m is the degree of ϕ . Thus, for instance, as noted by Accola and Farkas, a hyperelliptic surface of genus > 3 cannot be a 2-sheeted covering of an elliptic surface. Combining the above formula with the Riemann–Hurwitz relation, we get

$$m(h - 1) + 1 + \frac{1}{2}b_\phi = g \leq m(h + 1) - 1$$

or

$$b_\phi \leq 4(m - 1)$$

where b_ϕ is the total branch order of ϕ .

(If only Y is known to be hyperelliptic, certain restrictions are imposed on X . Thus, if Y is of genus ≥ 3 , no function of order 3 can exist on Y and hence not on X . The above formulas can be generalized to non-hyperelliptic cases).

Each divisor of degree 2 of the hyperelliptic series on X is mapped on a divisor of degree 2 in the image series on Y . Hence a sheet interchange on X followed by ϕ is equivalent to ϕ followed by a sheet interchange on Y . Thus, the function field on Y pulls back to a subfield on X invariant under the hyperelliptic involution.

(The above argument can be carried out for complete, irreducible algebraic curves, over a general groundfield, whose canonical images in the jacobian variety are invariant (modulo translation) under the involution $u \rightarrow (-u)$. This yields a geometric interpretation of a result of Tamme [*Ein Satz über Hyperelliptische Funktionenkörper*, J. Reine Angew. Math. 257 (1972) 217–220]. Of course, our main remark can also be proved in the more general context).

The result for hyperelliptic surfaces given in Corollary was mentioned by R. D. M. Accola, *Advances in the Theory of Riemann Surfaces*, Ann. of Math. Studies, No. 66, Princeton Univ. Press, Princeton N.J. 1971, pp. 7–18. A special case of our main remark was proven by M. Newman, Math. Ann. 196 (1972) 198–217.

Institut Mittag-Leffler, S-182 62 Djursholm, Schweden
ab juni 1977: *Institutt for Matematikk*
7034 NTH Trondheim, Norway

Received March 7, 1977

