

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 53 (1978)

Artikel: Level sets of infinite length.
Autor: Piranian, George / Weitsman, Allen
DOI: <https://doi.org/10.5169/seals-40761>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Level sets of infinite length

GEORGE PIRANIAN AND ALLEN WEITSMAN

Dedicated to Professor Albert Pfluger on his seventieth birthday

The geometry of level sets plays an important role in the analysis of various function-theoretic problems. Often, however, the level sets are so complicated that one must either choose sets associated with special levels (see [3], [4]) or resort to the approximation of level sets by shorter curves (see [1, pp. 550–553]).

In [3] and [4], there are some weak estimates on the length $\ell(r, R)$ of the sets $\{z : |f(z)| = R, |z| < r\}$ associated with a function f meromorphic in the plane. For such an f we do not know whether the quantity $\ell(r, R)$ can be bounded in terms of Nevanlinna's characteristic function T without reference to exceptional levels. But the following is implicit in the results in [3, pp. 121–123] and [4, p. 44]: *If f is meromorphic in $|z| \leq 2r$ and $f(0) = 1$, then each subinterval $[\alpha, \beta]$ of $(0, \infty)$ contains a set I of measure $(\beta - \alpha)/2$ such that the inequality*

$$\ell(r, R) \leq 2\pi r \sqrt{\frac{\beta T(2r, f)}{(\beta - \alpha) \log 2}}$$

holds for each R in I .

For functions univalent in the disk $D = \{|z| < 1\}$, it follows from the basic inequalities of length and area [2, p. 18] that almost all level sets have finite length; but we do not know of any univalent function in D having even one level set of infinite length. Neither do we know whether there exists a bounded holomorphic function f in D such that $\ell(1, R) = \infty$ for all R in some interval (α, β) , or for all R in some set of positive measure. In this paper, we show that bounded holomorphic functions can have level sets of infinite length.

THEOREM. *If M is a finite or countable set in the interval $(0, 1)$, then there exists a holomorphic function f in D such that $|f(z)| < 1$ for all z in D and $\ell(1, R) = \infty$ for each R in M .*

Because for $0 \leq r < 1$ and $0 < R < \infty$ the quantity $\ell(r, R)$ is a continuous

function of r and R (for each function f holomorphic in D), the set of values R for which $\ell(1, R) = \infty$ is of type G_δ . It follows immediately that if the set M in the theorem is dense in $(0, 1)$, then the set of values R for which $\ell(1, R) = \infty$ is residual in $(0, 1)$.

To prove the theorem, we begin with the case where the set M consists of a single value R , and we construct a nondecreasing continuous function μ such that the level set $\ell(1, R)$ of the function

$$f(z) = \exp \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{z + e^{it}}{z - e^{it}} d\mu(t) \quad (1)$$

has infinite length.

We write $z = \rho e^{i\theta}$, and we observe that

$$|f(\rho e^{i\theta})| = \exp \frac{-1}{2\pi} \int_{-\pi}^{\pi} P(\rho, \theta, t) d\mu(t),$$

where P denotes the Poisson kernel

$$P(\rho, \theta, t) = \frac{1 - \rho^2}{1 - 2\rho \cos(t - \theta) + \rho^2}.$$

If $\mu'(t)$ exists at the point t , then the radial limit of $|f(z)|$ at e^{it} is $\exp(-\mu'(t))$. We shall now construct a sequence $\{\mu_n\}$ such that on some set of positive measure the sequence $\{\mu'_n(t) + \log R\}$ converges to 0 with alternating signs.

On some subinterval $[t_0, t_1]$ of $(-\pi, \pi)$, let μ_0 be linear, and let $\mu'_0(t)$ be slightly less than $-\log R$. On $[-\pi, t_0]$ and on $[t_1, \pi]$, let μ_0 be constant. Then the function

$$|f_0(\rho e^{i\theta})| = \exp \frac{-1}{2\pi} \int_{-\pi}^{\pi} P(\rho, \theta, t) d\mu_0(t)$$

is slightly greater than R , on some arc γ_0 joining e^{it_0} to e^{it_1} .

To construct μ_1 , we divide the interval (t_0, t_1) into subintervals in such a way that for each $\varepsilon > 0$ both $[t_0, t_0 + \varepsilon]$ and $[t_1 - \varepsilon, t_1]$ contain infinitely many of the subintervals. On each of the subintervals, we define μ_1 so that

- (i) $\mu_1(t) = \mu_0(t)$ at the endpoints,
- (ii) μ'_1 is constant and slightly greater than $-\log R$, on a portion beginning at the left end of the subinterval,
- (iii) $\mu'_1(t) = 0$ on the remainder of the subinterval (see Figure 1).




Figure 1

If the subdivision of (t_0, t_1) is fine enough, then the function f_1 determined by μ_1 has modulus greater than R on the previously established arc γ_0 . But on a family $\{\gamma_{1k}\}$ of other arcs (corresponding to the intervals on which $\mu'_1 > -\log R$) separated from the origin by the arc γ_0 , the value of $|f_1(z)|$ is less than R . Moreover, if the positive values of μ'_0 and μ'_1 are sufficiently close to $-\log R$, the sum of the diameters of the arcs γ_{1k} is greater than $(t_1 - t_0)/2$.

On each segment on which $\mu'_1(t) = 0$, let $\mu_2(t) = \mu_1(t)$. Each of the remaining segments of (t_0, t_1) we divide into infinitely many subsegments. On each of these subsegments we define μ_2 so that $\mu'_2(t)$ is constant and slightly less than $-\log R$ in a large part of the segment, while $\mu'_2(t) = -2 \log R$ in the remainder (see Figure 2).




Figure 2

The sequence $\{\mu_n\}$ constructed by the obvious continuation of our process converges to a function μ , and the function (1) has the property that on each of

certain arcs γ_{nk} its modulus is greater or less than R according as the index n is even or odd. If the values $\mu'_n(t)$ converge to $-\log R$ rapidly enough in the portions where they are neither 0 nor $-2 \log R$, then for each n the sum of the diameters of the arcs γ_{nk} is greater than $(t_1 - t_0)/2$. If γ_{nh} separates the origin from $\gamma_{n+1,k}$ in D , then by continuity of f an arc of the level set $\{z : |f(z)| = R\}$ separates γ_{nh} from $\gamma_{n+1,k}$. It follows that the level set has infinite length.

We have proved the theorem for the special case where the set M is a singleton. In the general case, let $M = \{R_j\}$. We divide the interval $(-\pi, \pi)$ into infinitely many disjoint subintervals T_j , and we construct the sequence $\{\mu_n\}$ as in the special case, except that now the limit of $\mu'_n(t)$ is $-\log R_j$ in a substantial part of T_j . A detailed description would require the development of tedious notation; but the underlying principle is simple: if in the passage from μ_{n-1} to μ_n we make the new system of subdivisions fine enough, then on the arcs of the families $\{\gamma_{mk}\}$ ($m = 1, 2, \dots, n-1$) the appropriate differences $|f_n| - R_j$ and $|f_{n-1}| - R_j$ have the same sign.

Because the limit μ of the sequence $\{\mu_n\}$ is absolutely continuous, our function (1) is an outer function. We do not know whether a nonconstant inner function can have a level set of infinite length. Since the radial limit of an inner function has modulus 1 almost everywhere, the construction of an example may well require delicate computations.

This paper developed from conversations held in Lexington during the function-theoretic conference of May 19th to 22nd, 1976, in honor of Professor S. M. Shah on the occasion of his retirement. The authors acknowledge with gratitude the warm hospitality of the Mathematics Department of the University of Kentucky.

REFERENCES

1. CARLESON, L., *Interpolations by bounded analytic functions and the corona problem*, Ann. of Math. (2) 76 (1962), 547-559.
2. HAYMAN, W. K., *Multivalent functions*, Cambridge Tracts in Mathematics and Mathematical Physics, No. 48; Cambridge University Press, Cambridge, England, 1958.
3. WEITSMAN, A., *Meromorphic functions with maximal deficiency sum and a conjecture of F. Nevanlinna*, Acta Math. 123 (1969), 115-139.
4. —, *A theorem on Nevanlinna deficiencies*, Acta Math. 128 (1972), 41-52.

University of Michigan,
Ann Arbor, Michigan 48109
Purdue University, West Lafayette,
Indiana 47907

Received March 16, 1977