

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 54 (1979)  
  
**Artikel:** A note on groups with torsion-free abelianization and trivial multiplier.  
**Autor:** Strebel, Ralph  
**DOI:** <https://doi.org/10.5169/seals-41567>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# A note on groups with torsion-free abelianization and trivial multiplier

RALPH STREBEL

## 1. Introduction

1.1. A basic result on free groups  $F$  asserts that the factors  $\{F_j/F_{j+1}\}_{1 \leq j < \omega}$  of successive terms of the lower central series are free abelian (Magnus [4], Witt [11]). This can be proved using Lie algebra techniques and a proof (e.g. [1], pp. 35–39; or [7], LA 4.10–4.13) will then rely on three corner stones:

- \* The canonical Lie algebra homomorphism  $\sigma : L_X \rightarrow \text{Ass}_X$  from the free Lie  $\mathbf{Z}$ -algebra on the set  $X$  into the Lie algebra of the free associative  $\mathbf{Z}$ -algebra on  $X$  is *injective*.
- \* To every group  $G$  is associated a graded Lie algebra  $\text{gr } G$ . Its underlying additive group is the direct sum  $\bigoplus G_j/G_{j+1}$  of the factors of successive terms of the lower central series of  $G$ . Its Lie bracket is on homogeneous components induced by commutation in the group, sc.

$$(g \cdot G_{j+1}, h \cdot G_{k+1}) \mapsto (g^{-1}h^{-1}gh) \cdot G_{j+k+1} \quad (g \in G_j, h \in G_k)$$

and then extended linearly.

Reverting to free algebras and groups, let  $\gamma : L_X \rightarrow \text{gr } F_X$  denote the Lie algebra homomorphism defined by the assignments

$$X \ni x \mapsto x \cdot F_2 \in \text{gr}^1 F_X \subseteq \text{gr } F_X.$$

- \* There exists a Lie algebra homomorphism  $\alpha : \text{gr } F_X \rightarrow \text{Ass}_X$ , making the triangle

$$\begin{array}{ccc} L_X & \xrightarrow{\sigma} & \text{Ass}_X \\ \downarrow \gamma & \nearrow \alpha & \\ \text{gr } F_X & & \end{array}$$

commute.

1.2. In this note it is shown that the proof sketched above can be adapted to the more general situation in which  $F_X$  gets replaced by a group  $G$  whose abelianization  $G_{ab}$  is torsion-free and whose multiplier  $H_2(G, \mathbf{Z})$ , i.e. whose second homology group with integral coefficients, is a torsion group. Such a group will be referred to as being TFT. The place of the free associative  $\mathbf{Z}$ -algebra  $\text{Ass}_X$  will be taken by the tensor algebra  $TG_{ab}$ . The main problem is the existence of a Lie algebra homomorphism

$$\alpha : \text{gr } G \rightarrow TG_{ab}$$

extending the identification  $\text{gr } G \simeq TG_{ab}$  in degree 1.

**THEOREM 1.** *If  $G$  is TFT then the identification  $\text{gr}^1 G \simeq T^1 G_{ab}$ , taking  $g \cdot G_2 \in \text{gr } G$  to  $g \cdot G_2 \in TG_{ab}$ , extends (necessarily uniquely) to a Lie algebra isomorphism  $\alpha : \text{gr } G \simeq TG_{ab}$  from the graded Lie  $\mathbf{Z}$ -algebra  $\text{gr } G$  associated with  $G$  onto the Lie subalgebra of  $TG_{ab}$  generated by  $G_{ab} = T^1 G_{ab}$ . Moreover,  $\alpha$  induces an isomorphism  $U\alpha : U(\text{gr } G) \simeq TG_{ab}$  between associative  $\mathbf{Z}$ -algebras, thus providing a model for the universal algebra of  $\text{gr } G$ .*

1.3. Since the additive group underlying the tensor algebra  $TG_{ab}$  is torsion-free if  $G_{ab}$  is so, Theorem 1 immediately entails the approved.

**COROLLARY 1.** *The factors  $G_j/G_{j+1}$ ,  $j = 1, 2, \dots$ , of the lower central series of a TFT group  $G$  are torsion-free.*

An application of Corollary 1 can be found in [10] (cf. 4.1.).

The next corollary indicates that even in the special case of a TFT group the finer commutator structure gets lost in the passage from  $G$  to  $\text{gr } G$ . (Examples testifying the loss will be given in 4.6.ff.)

**COROLLARY 2.** *The Lie algebra  $\text{gr } G$  of a TFT group is determined by its first homogeneous component  $G_{ab}$ .*

1.4. Our second main result deals with subgroups of TFT groups. We state it as

**THEOREM 2.** *Let  $\varphi : G \rightarrow \bar{G}$  be a homomorphism for which  $\varphi^1 : G_{ab} \rightarrow \bar{G}_{ab}$  is injective. Suppose  $G$  is TFT and  $H_2(\bar{G}, \mathbf{Z})$  is a torsion group. Then  $\text{gr } \varphi : \text{gr } G \rightarrow \text{gr } \bar{G}$  is injective. Put differently,  $\varphi$  induces injective homomorphisms  $\varphi_* : G/G_j \rightarrow \bar{G}/\bar{G}_j$  for all  $j \geq 2$ .*

If  $G$  and  $\bar{G}$  in Theorem 2 are both free the claim reduces to a well-known result of Malcev on subgroups of free nilpotent groups ([5]; cf. [6], 42.51). Theorem 2 may also be compared with the following result:

**THEOREM** (Stallings [8], Stambach [9]). *Let  $\varphi: G \rightarrow \bar{G}$  be a homomorphism inducing an isomorphism  $\varphi^1: G_{ab} \rightarrow \bar{G}_{ab}$  and a surjection  $H_2(\varphi): H_2(G, \mathbf{Z}) \rightarrow H_2(\bar{G}, \mathbf{Z})$ . Then  $\text{gr } \varphi: \text{gr } G \rightarrow \text{gr } \bar{G}$  is an isomorphism of graded Lie algebras.*

## 2. The proof of Theorem 1

2.1. Let  $R$  be a non-trivial commutative ring with 1. If  $G$  is a group, let  $RG$  denote its group algebra (over  $R$ ) and  $\varepsilon: RG \rightarrow R$  the associated augmentation, i.e. the  $R$ -algebra homomorphism sending every  $g \in G$  to  $1 \in R$ . The kernel of  $\varepsilon$  is called the augmentation ideal  $I = I(RG)$  and, as an  $R$ -module, it is freely generated by the elements  $g - 1$  ( $g \in G \setminus \{e\}$ ). The powers  $\{I^j\}_{0 \leq j < \omega}$  form an integral filtration of  $RG$  whose associated graded  $R$ -algebra will be denoted by  $\text{gr } RG$ .

Define a descending chain of subsets of  $G$  by setting

$$D_R^j(G) = \{g \in G \mid g - 1 \in I^j\} \quad (1 \leq j < \omega).$$

Then  $D_R^1(G) = G$ , each  $D_R^j(G)$  is a (normal) subgroup and for every pair  $(j, k) \in \mathbf{N}^2$  the commutator  $[D_R^j(G), D_R^k(G)]$  is contained in  $D_R^{j+k}(G)$  (see, e.g., [2], §4.5, Prop. 2, p. 42). Hence  $\{D_R^j(G)\}_{1 \leq j < \omega}$  is a central series of  $G$  and we can form the associated graded Lie  $\mathbf{Z}$ -algebra  $\text{gr } \{D_R(G)\}$ . The function  $g \mapsto g - 1$  induces then an *injective* Lie algebra homomorphism

$$\beta: \text{gr } \{D_R(G)\} \rightarrow \text{gr } RG.$$

(It is clear that  $\beta$  is actually a natural transformation between functors from the category of groups to the category of graded Lie  $\mathbf{Z}$ -algebras.)

2.2. We specialize now to the case  $R = \mathbf{Z}$ . Then  $D_{\mathbf{Z}}^2(G) = G_2 = G'$  and  $\beta$  gives an isomorphism  $\beta^1: G/G_2 \xrightarrow{\sim} I/I^2$ ,  $gG_2 \mapsto (g-1) + I^2$ . If  $TG_{ab}$  is the tensor algebra on  $G_{ab} = G/G'$  the isomorphism  $\beta^1$  will extend uniquely to a homomorphism  $\mu: TG_{ab} \rightarrow \text{gr } \mathbf{Z}G$  of graded associative  $\mathbf{Z}$ -algebras, given in degree  $j$  by

$$g_1 G_2 \otimes g_2 G_2 \otimes \cdots \otimes g_j G_2 \mapsto (g_1 - 1)(g_2 - 1) \cdots (g_j - 1) + I^{j+1}.$$



Clearly  $\mu$  is always surjective. For TFT groups it is even bijective according to the following

LEMMA. If  $G_{ab}$  is torsion-free and  $H_2(G, \mathbf{Z})$  is a torsion group then  $\mu : TG_{ab} \xrightarrow{\sim} \text{gr } \mathbf{Z}G$  is an isomorphism of graded associative  $\mathbf{Z}$ -algebras.

2.3. *Proof.* For every  $j \geq 0$  the short exact sequence  $I^{j+1} \hookrightarrow I^j \twoheadrightarrow I^j/I^{j+1}$  of right  $G$ -modules induces a long exact sequence. In dimensions 2, 1 and 0 it looks like this:

$$\begin{array}{ccccccc} & & H_2(G, I^j/I^{j+1}) & \xleftarrow{\pi_*} & H_2(G, I^j) & & \\ \partial_2 \swarrow & & & & & & \\ & H_1(G, I^{j+1}) & \longrightarrow & H_1(G, I^j) & \xrightarrow{\pi_*} & H_1(G, I^j/I^{j+1}) & \searrow \partial_1 \\ & 0 \longleftarrow (I^j/I^{j+1}) \otimes_G \mathbf{Z} & \xleftarrow{\pi_*} & I^j \otimes_G \mathbf{Z} & \longleftarrow & I^{j+1} \otimes_G \mathbf{Z} & \end{array} \quad (1)$$

One readily verifies that the composite

$$\bar{\mu} : I/I^2 \otimes I^j/I^{j+1} = H_1(G, \mathbf{Z}) \otimes I^j/I^{j+1} \xrightarrow{\sim} H_1(G, I^j/I^{j+1})$$

$$\xrightarrow{\partial_1} I^{j+1} \otimes_G \mathbf{Z} \xrightarrow{\sim} I^{j+1}/I^{j+2}$$

is the obvious multiplication map. Taking into account that  $I^j \otimes_G \mathbf{Z} \rightarrow (I^j/I^{j+1}) \otimes_G \mathbf{Z}$  is an isomorphism and using the universal coefficient theorem, the sequence (1) can be rewritten as

$$\begin{array}{ccccccc} & & (H_2(G, \mathbf{Z}) \otimes I^j/I^{j+1} \oplus \text{Tor}_1^{\mathbf{Z}}(G_{ab}, I^j/I^{j+1})) & & & & \\ \partial_2 \swarrow & & & & & & \\ & H_1(G, I^{j+1}) & \longrightarrow & H_1(G, I^j) & \longrightarrow & I/I^2 \otimes I^j/I^{j+1} & \xrightarrow{\bar{\mu}} I^{j+1}/I^{j+2} \longrightarrow 0. \end{array} \quad (2)$$

This exact sequence allows, first of all, to prove that all homology groups  $H_1(G, I^j)$  ( $0 \leq j < \omega$ ) are torsion groups. To see this recall that  $H_2(G, \mathbf{Z})$  is a torsion group by hypothesis and  $\text{Tor}_1^{\mathbf{Z}}(?, ?)$  by nature, and that  $H_1(G, \mathbf{Z}G) = 0$ ; then use the exactness of (2). Secondly, (2) implies that all multiplication maps  $\bar{\mu} : I/I^2 \otimes I^j/I^{j+1} \rightarrow I^{j+1}/I^{j+2}$  are bijective. As all  $H_1(G, I^j)$  are torsion groups it will do to show inductively that  $I/I^2 \otimes I^j/I^{j+1}$  is torsion-free. This follows from the hypothesis that  $G_{ab} \cong I/I^2$  be torsion-free and the fact that the tensor product (over  $\mathbf{Z}$ ) of torsion-free groups is again torsion-free. The proof is now easily completed.

2.4. *The proof of Theorem 1.* Assume  $G_{ab}$  is torsion-free and  $H_2(G, \mathbf{Z})$  is a torsion group. By Lemma 2.2 the map  $\mu: TG_{ab} \rightarrow \text{gr } \mathbf{Z}G$  is bijective so that we can define a Lie algebra homomorphism  $\alpha$  as the composite

$$\text{gr } G \xrightarrow{\iota} \text{gr } \{D_{\mathbf{Z}}(G)\} \xrightarrow{\beta} \text{gr } \mathbf{Z}G \xleftarrow{\mu} TG_{ab}.$$

Here  $\iota$  denotes the Lie algebra homomorphism stemming from the inclusions  $G_j \subseteq D_{\mathbf{Z}}^j(G)$ . Note that  $\text{gr } G$  is generated by its first homogeneous component and that  $\alpha^1: \text{gr}^1 G \rightarrow T^1 G_{ab}$  is the identity on  $G_{ab}$ . These facts, together with the universal property of  $TG_{ab}$ , imply that  $\alpha: \text{gr } G \rightarrow TG_{ab}$  is the canonical map of  $\text{gr } G$  into its universal algebra and so prove the addendum to Theorem 1.

2.5. We are left with proving that  $\alpha$  is injective. If  $F_X$  is free on the set  $X$  then  $(F_X)_{ab}$  is free-abelian and  $H_2(F_X, \mathbf{Z}) = 0$ . Hence  $\alpha$  is defined and gives the classical Lie algebra homomorphism

$$\alpha: \text{gr } F_X \rightarrow T(F_X)_{ab} \cong \text{Ass}_X, \quad x \cdot F_2 \mapsto x \quad (x \in X).$$

The theory of basic sequences (see, e.g. [1]) or the Poincaré–Birkhoff–Witt theorem (see e.g. [7]) can then be used to prove that  $\alpha$  is injective.

Now let  $\varphi^1: F_{ab} \hookrightarrow G_{ab}$  be a finitely generated free-abelian subgroup of our torsion-free abelianization  $G_{ab}$ . Lift the inclusion to a group homomorphism  $\varphi: F \rightarrow G$ . The lift gives rise to the commutative square

$$\begin{array}{ccc} \text{gr } F & \xrightarrow{\alpha_F} & T F_{ab} \\ \downarrow \text{gr } \varphi & & \downarrow T \varphi^1 \\ \text{gr } G & \xrightarrow{\alpha_G} & T G_{ab} \end{array}$$

In it  $\alpha_F$  is injective, and because  $F_{ab}$  and  $G_{ab}$  are both torsion-free abelian groups and  $\varphi^1$  is injective,  $T \varphi^1$  is likewise injective. Consequently the restriction of  $\alpha_G$  to the image of  $\text{gr } \varphi$  is injective. But  $\text{gr } G$  is generated by its first homogeneous component  $G_{ab}$  and  $G_{ab}$ , being torsion-free, is a union of finitely generated free-abelian subgroups. This proves that  $\alpha$  is injective and establishes the claim of Theorem 1. The proofs of the corollaries present no problems.

2.7. *Remark.* The injectivity of  $\alpha$  could also have been inferred from a (rather difficult) theorem of M. Lazard [3] asserting that the canonical map of a Lie  $R$ -algebra into its universal algebra is injective if  $R$  is a principal ideal domain.

### 3. The proof of Theorem 2

3.1. We first return to the set-up of Subsection 2.1 and choose  $R$  to be the rational numbers  $\mathbf{Q}$ . The commutative square

$$\begin{array}{ccc} G/D_{\mathbf{Z}}^2(G) & \xrightarrow{\beta_{\mathbf{Z}}^1} & I/I^2 \\ \downarrow \text{can} & & \downarrow \text{can} \\ G/D_{\mathbf{Q}}^2(G) & \xrightarrow{\beta_{\mathbf{Q}}^1} & \text{gr}^1 \mathbf{Q}G \cong I/I^2 \otimes \mathbf{Q} \end{array}$$

shows that  $D_{\mathbf{Q}}^2(G)$  equals  $\ker \{G \rightarrow G_{ab} \otimes \mathbf{Q}\}$  whence

$$\beta_{\mathbf{Q}}^1 \otimes \mathbf{Q}: G_{ab} \otimes \mathbf{Q} \cong G/D_{\mathbf{Q}}^2(G) \otimes \mathbf{Q} \rightarrow \text{gr}^1 \mathbf{Q}G$$

is an isomorphism. It extends uniquely to a homomorphism

$$\mu_{\mathbf{Q}}: T(G_{ab} \otimes \mathbf{Q}) \rightarrow \text{gr} \mathbf{Q}G$$

of graded associative  $\mathbf{Q}$ -algebras. Clearly  $\mu_{\mathbf{Q}}$  is onto. An easy modification of the proof of Lemma 2.2 reveals that  $\mu_{\mathbf{Q}}$  is also injective provided merely that  $H_2(G, \mathbf{Z})$  is a torsion group. For a group  $G$  whose multiplier is a torsion group one can therefore define a homomorphism

$$\alpha_{\mathbf{Q}}: \text{gr} \{D_{\mathbf{Q}}(G)\} \xrightarrow{\beta_{\mathbf{Q}}^1} \text{gr} \mathbf{Q}G \xleftarrow{\mu_{\mathbf{Q}}} T(G_{ab} \otimes \mathbf{Q})$$

of graded Lie  $\mathbf{Z}$ -algebras.

3.2. Now let  $G$  be TFT, let  $\bar{G}$  be a group with  $H_2(\bar{G}, \mathbf{Z})$  a torsion group and let  $\varphi: G \rightarrow \bar{G}$  be a group homomorphism. Then the canonical maps  $\alpha(G)$ ,  $\alpha_{\mathbf{Q}}(G)$  and  $\alpha_{\mathbf{Q}}(\bar{G})$  are all three defined and they combine to produce the following commutative diagram

$$\begin{array}{ccc} \text{gr } G & \xrightarrow{\alpha(G)} & T G_{ab} \\ \downarrow \iota & & \downarrow T\kappa \\ \text{gr} \{D_{\mathbf{Q}}(G)\} & \xrightarrow{\alpha_{\mathbf{Q}}(G)} & T(G_{ab} \otimes \mathbf{Q}) \\ \downarrow \text{gr}_{\mathbf{Q}} \varphi & & \downarrow T(\varphi^1 \otimes \mathbf{Q}) \\ \text{gr} \{D_{\mathbf{Q}}(\bar{G})\} & \xrightarrow{\alpha_{\mathbf{Q}}(\bar{G})} & T(\bar{G}_{ab} \otimes \mathbf{Q}) \end{array}$$

In it  $\iota$  denotes the canonical Lie algebra homomorphism stemming from the inclusions  $G_j \subseteq D_{\mathbf{Q}}^j(G)$ , and  $\kappa: G_{ab} \rightarrow G_{ab} \otimes \mathbf{Q}$  is the obvious canonical  $\mathbf{Z}$ -module homomorphism.

By assumption  $G_{ab}$  is torsion-free. Therefore  $\kappa$  and  $T\kappa$  are injective. By Theorem 1 the same is true for  $\alpha(G)$ . If, as is required in the hypotheses of Theorem 2,  $\varphi^1: G_{ab} \rightarrow \bar{G}_{ab}$  is injective  $T(\varphi^1 \otimes \mathbf{Q})$  will also be injective. Hence the composite  $\iota \circ \text{gr}_{\mathbf{Q}} \varphi: \text{gr } G \rightarrow \text{gr } \{D_{\mathbf{Q}}(G)\}$  is seen to be injective and the claim of Theorem 2 follows upon noting that  $\iota \circ \text{gr}_{\mathbf{Q}} \varphi$  factors through  $\text{gr } \varphi: \text{gr } G \rightarrow \text{gr } \bar{G}$ .

#### 4. Examples and counter-examples

**4.1. E-groups.** Let  $G$  be a group having torsion-free abelianization and trivial multiplier. If  $G_{ab}$  is even free-abelian the Stallings–Stammbach theorem quoted in 1.4 applies and proves that each  $G_j/G_{j+1}$  is isomorphic with the corresponding factor  $F_j/F_{j+1}$  of a suitable free group  $F$  and so, in particular, torsion-free.

This argument breaks down if  $G_{ab}$  is not free abelian, as it usually happens when  $G$  is the derived group of a knot group or, more generally, when  $G$  is an **E-group** in the sense of [10]. A group  $G$  is there called an **E-group** if  $G_{ab}$  is torsion-free and if the  $G$ -trivial module  $\mathbf{Z}$  admits a  $\mathbf{Z}G$ -projective resolution  $\cdots \rightarrow P_2 \xrightarrow{\partial_2} P_1 \rightarrow P_0 \rightarrow \mathbf{Z} \rightarrow 0$  for which the induced differential  $1 \otimes \partial_2: \mathbf{Z} \otimes_G P_2 \rightarrow \mathbf{Z} \otimes_G P_1$  is injective. The condition on  $1 \otimes \partial_2$  implies that  $H_2(G, \mathbf{Z})$  is zero; the converse, however, is false (see 4.2).

**E-groups** have the following stability property: if  $G \in \mathbf{E}$  and  $N \triangleleft G$  is a normal subgroup with torsion-free, abelian factor group then  $N \in \mathbf{E}$ . In particular, the terms of the derived series of an **E-group** are **E-groups** and so are the terms of the lower central series.

**4.2. Groups  $G$  with  $G_{ab}$  torsion-free,  $H_2(G, \mathbf{Z}) = 0$  but  $G \notin \mathbf{E}$ .** It suffices to prove that  $G$  does not have the stability property enjoyed by **E-groups**. Let  $A$  be an abelian group possessing an automorphism  $\tau$  for which  $\tau - 1: A \rightarrow A$  is bijective and  $\tau \wedge \tau - 1 \wedge 1: A \wedge A \rightarrow A \wedge A$  is onto. Let  $C = \langle t \rangle$  be an infinite cyclic group and define  $G$  to be the split extension  $A \triangleleft C$  where  $t$  induces on  $A$  the given  $\tau$ . Then  $A = G_2$ ,  $G_{ab} \cong \mathbf{Z}$  and  $H_2(G, \mathbf{Z}) = 0$ , although  $A$  is in general neither torsion-free nor has it trivial multiplier (take e.g.  $A = (\mathbf{Z}/5\mathbf{Z}) \oplus (\mathbf{Z}/5\mathbf{Z})$  and let  $\tau$  operate by multiplication by 2).

**4.3.** We give next two examples demonstrating that  $\alpha: \text{gr } G \rightarrow TG_{ab}$  need not exist if the hypotheses of Theorem 1 are weakened. Consider first an *abelian group*  $A$ . Then  $\text{gr } A$  is a *commutative* graded Lie algebra concentrated in degree 1 and its universal algebra is the symmetric algebra  $SA$  of  $A$ . Hence  $\alpha: \text{gr } A \rightarrow TA$  can only exist if  $TA$  is commutative. The commutativity of  $\otimes^2 A$ , in turn, is equivalent with the vanishing of the exterior square  $\Lambda^2 A \cong H_2(A, \mathbf{Z})$ ; for the canonical map  $A \wedge A \rightarrow A \otimes A$  taking  $a \wedge b$  to  $a \otimes b - b \otimes a$  is injective. For a

torsion-free abelian group we thus get the following conclusion: The identification  $\text{gr}^1 A \simeq T^1 A$  extends to a Lie algebra homomorphism  $\alpha : \text{gr} A \rightarrow T A$  if and only if  $H_2(A, \mathbf{Z}) = 0$ .

4.4. Groups  $G$  with  $H_2(G, \mathbf{Z}) = 0$  but  $G_{ab}$  not torsion-free. The exact sequence

$$H_2(G, \mathbf{Z}) \longrightarrow I/I^2 \otimes I/I^2 \xrightarrow{\bar{\mu}} I^2/I^3 \longrightarrow 0$$

(cf. sequence (2) in 2.3.) shows that  $\mu^2 : \otimes G_{ab} \simeq I^2/I^3$  is bijective. Consequently the identification  $\text{gr}^1 G \simeq T^1 G_{ab}$  extends to

$$\alpha^2 : G_2/G_3 \longrightarrow I^2/I^3 \xleftarrow{\mu^2} \otimes^2 G_{ab}$$

taking  $[g, h] \cdot G_3$  to  $g \cdot G_2 \otimes h \cdot G_2 - h \cdot G_2 \otimes g \cdot G_2$ . (The existence of  $\alpha^2$  can also be deduced from the 5-term sequence associated with the extension  $G_2 \triangleleft G \twoheadrightarrow G_{ab}$ , namely

$$H_2(G, \mathbf{Z}) \rightarrow H_2(G_{ab}, \mathbf{Z}) \xrightarrow{\chi} G_2/G_3 \rightarrow G_{ab} \simeq G_{ab} \rightarrow 0, \quad (3)$$

and from the facts that  $H_2(G_{ab}, \mathbf{Z}) \cong G_{ab} \wedge G_{ab}$ , that under this isomorphism  $\chi$  becomes the obvious commutator map and that  $\Lambda^2 G_{ab}$  maps canonically into  $\otimes^2 G_{ab}$ .)

However, it is in general not possible to extend the identification  $\alpha^1 : \text{gr}^1 G \simeq T^1 G_{ab}$  to a Lie algebra homomorphism

$$\alpha_* : G/G_2 \oplus G_2/G_3 \oplus G_3/G_4 \rightarrow G_{ab} \oplus \otimes^2 G_{ab} \oplus \otimes^3 G_{ab}$$

of nilpotent Lie algebras of class two. To see this let  $G$  be a one-relator group of the form

$$G = \langle a, t; t^{-1}at = a^m \rangle = \langle a, t; [a, t] = a^{m-1} \rangle \quad (m \in \mathbf{Z} \setminus \{0, 1, 2\}).$$

Then  $G_{ab} = \text{gp}(aG_2) \times \text{gp}(tG_2) \cong (\mathbf{Z}/|m-1|\mathbf{Z}) \times \mathbf{Z}$  and  $H_2(G, \mathbf{Z}) = 0$ . The iterated commutator  $[a, [a, t]]$  represents the trivial element in  $G_3/G_4$ , whereas the corresponding Lie bracket in  $\otimes^3 G_{ab}$ , namely

$$[aG_2, [aG_2, tG_2]] = aG_2 \otimes aG_2 \otimes tG_2 - 2 \cdot aG_2 \otimes tG_2 \otimes aG_2 + tG_2 \otimes aG_2 \otimes aG_2$$

has order  $|m-1| > 1$ .

4.5. Groups  $G, \bar{G}$  with trivial multiplier,  $\varphi: G \rightarrow \bar{G}$  with  $\varphi^1$  injective but  $G_{ab}$  not torsion-free. Our goal is to show that  $\varphi^2: G_2/G_3 \rightarrow \bar{G}_2/\bar{G}_3$  is not always injective. Let  $G$  be the one-relator group  $\langle a, t; t^{-1}at = a^m \rangle$  considered before and let  $\bar{G}$  arise out of  $G$  by adjoining a  $k^{\text{th}}$  root of  $t$ , i.e.

$$\bar{G} = G \underset{t=u^k}{*} (u) = \langle a, u; u^{-k}au^k = a^m \rangle \quad (k \geq 2),$$

and let  $\varphi: G \rightarrow \bar{G}$  be the canonical injection. Then  $H_2(G, \mathbf{Z}) = H_2(\bar{G}, \mathbf{Z}) = 0$  and  $\varphi^1: G_{ab} \rightarrow \bar{G}_{ab}$  is injective. The map  $\varphi^2: G_2/G_3 \rightarrow \bar{G}_2/\bar{G}_3$  can be identified with the exterior square  $\Lambda^2 \varphi^1: \Lambda^2 G_{ab} \rightarrow \Lambda^2 \bar{G}_{ab}$  (consult (3) above). Both  $\Lambda^2 G_{ab}$  and  $\Lambda^2 \bar{G}_{ab}$  are cyclic of order  $|m-1|$  and  $\Lambda^2 \varphi^1$  takes the generator  $aG_2 \wedge tG_2$  to  $aG_2 \wedge u^k G_2 = k(aG_2 \wedge uG_2)$ . Hence  $\varphi^2$  is injective if and only if  $k$  and  $m$  are relatively prime.

This example shows that the conclusion of Theorem 2 becomes false if  $G_{ab}$  is not assumed to be torsion-free, everything else remaining unchanged. It is clear that a strong assumption on  $H_2(\bar{G}, \mathbf{Z})$  is necessary to exclude cases like the abelianization  $\varphi: F \rightarrow F_{ab}$  of a free group. But I have not been able to determine to what extent the hypothesis on  $H_2(G, \mathbf{Z})$  could be weakened without jeopardizing the claim. (The theorem of Stallings–Stammbach quoted in 1.4. bears also on the issue.)

4.6. A family of  $2^{\aleph_0}$  non-isomorphic groups with trivial multiplier having all the same torsion-free abelianization. Let  $\{ {}_k F \}_{k \in \mathbf{N}}$  be a sequence of free groups of rank two, say  ${}_k F$  is free on  $x_k$  and  $y_k$ , and let  $ab: {}_k F \twoheadrightarrow ({}_k F)_{ab}$  be the abelianizations. If

$$\varphi = \{ \varphi_k: ({}_k F)_{ab} \rightarrow ({}_{k+1} F)_{ab} \}_{k \in \mathbf{N}}$$

is a given sequence of homomorphisms it can be lifted to a sequence

$$\Phi = \{ \Phi_j: {}_k F \rightarrow {}_{k+1} F \}_{k \in \mathbf{N}}$$

so as to produce a commutative ladder

$$\begin{array}{ccccccc} {}_1 F & \xrightarrow{\Phi_1} & {}_2 F & \xrightarrow{\Phi_2} & {}_3 F & \xrightarrow{\Phi_3} & {}_4 F \longrightarrow \cdots \\ \downarrow ab & & \downarrow ab & & \downarrow ab & & \downarrow ab \\ ({}_1 F)_{ab} & \xrightarrow{\varphi_1} & ({}_2 F)_{ab} & \xrightarrow{\varphi_2} & ({}_3 F)_{ab} & \xrightarrow{\varphi_3} & ({}_4 F)_{ab} \longrightarrow \cdots \end{array}$$

If the  $\varphi_k$  are injective the lifts  $\Phi_k$  are likewise injective, e.g. because of Theorem 2 and the residual nilpotency of free groups. The direct limit  $G_\Phi = \text{colim } \Phi$  is

therefore a locally free group with trivial multiplier and torsion-free abelianization  $(G_\Phi)_{ab} = \text{colim } \varphi$ ; and  $\text{gr } G_\Phi$  is isomorphic to the Lie algebra of  $T(G_\Phi)_{ab} \cong T(\text{colim } \varphi)$  generated by its first homogeneous component  $\text{colim } \varphi$ . In particular,  $\text{gr } G_\Phi$  depends only on  $\varphi$  and not on the choice of the lift  $\Phi$ .

Next let  $P$  be an infinite set of odd rational primes and let  $\lambda: \mathbf{N} \rightarrow P$  be an enumeration of  $P$ . Define the sequence  $\varphi = \{\varphi_k\}$  by

$$\varphi_k: x_k \cdot ({}_k F)_2 \mapsto x_{k+1}^{\lambda_k} \cdot ({}_{k+1} F)_2 \quad \text{and} \quad y_k \cdot ({}_k F)_2 \mapsto y_{k+1}^{\lambda_k} \cdot ({}_{k+1} F)_2.$$

The direct limit  $\text{colim } \varphi$  can be identified with the direct sum  $A_x \oplus A_y$  of two copies of the subgroup of the rationals generated by the elements  $1/p$  ( $p \in P$ ). For each  $S \subseteq \mathbf{N}$  define a lift  $\Phi(S)$  of  $\varphi$  by the formulae

$$\Phi_k(S): x_k \mapsto \begin{cases} x_{k+1}^{\lambda_k} & \text{if } k \in S \\ x_{k+1}^{\lambda_k} [y_{k+1}, x_{k+1}] & \text{if } k \notin S \end{cases} \quad \text{and} \quad y_k \mapsto y_{k+1}^{\lambda_k}.$$

We shall prove that  $\text{colim } \Phi(S)$  and  $\text{colim } \Phi(S')$  are isomorphic if and only if the symmetric difference of  $S$  and  $S'$  is finite. Since  $\mathbf{N}$  can be written as a disjoint union of infinitely many infinite subsets this will imply that there are  $2^{\aleph_0}$  many non-isomorphic locally free groups whose associated graded Lie  $\mathbf{Z}$ -algebras are isomorphic.

4.7. If the symmetric difference of  $S$  and  $S'$  is finite then clearly  $\text{colim } \Phi(S)$  and  $\text{colim } \Phi(S')$  are isomorphic. The converse will be established by showing that, up to a finite error,  $S$  can be recovered from the nilpotent quotient of class two  $G_{\Phi(S)}/(G_{\Phi(S)})_3$ .

Let  $F$  be free on  $x$  and  $y$ . The elements of  $H = F/F_3$  can be parametrized by the lattice points  $\mathbf{Z}^3$  via

$$\mathbf{Z}^3 \ni (a, b, c) \leftrightarrow x^a y^b (y^{-1} x^{-1} y x)^c \cdot F_3 \in F/F_3 = H.$$

The resulting group multiplication on  $\mathbf{Z}^3$  is then given by

$$(a, b, c) \cdot (a', b', c') = (a + a', b + b', ba' + c + c').$$

Note that this group multiplication has an obvious extension to points of  $\mathbf{Q}^3$ .

For positive powers and roots of elements of  $H = H_{\mathbf{Z}} \subseteq H_{\mathbf{Q}}$  one gets

$$(a, b, c)^m = \left( ma, mb, mc + \binom{m}{2} \cdot a \cdot b \right)$$

$$(a, b, c)^{1/m} = (a/m, b/m, c/m - \frac{1}{2} \cdot (m-1) \cdot (a/m) \cdot (b/m))$$

It follows that an element of  $H_{\mathbf{Z}}$  is an  $m^{\text{th}}$  power ( $m$  an *odd* integer) if and only if all three entries are integral multiples of  $m$ .

The endomorphism  $\Phi^{\epsilon}$  of  $H$  corresponding to the lifts  $\Phi_k$  with  $k \in S$  has the parametric description

$$(a, b, c)\Phi^{\epsilon} = (\lambda_k \cdot a, \lambda_k \cdot b, (\lambda_k)^2 \cdot c).$$

It has the property that the image of an element of  $H$  which is an  $m^{\text{th}}$  power is at least a  $(\lambda_k \cdot m)^{\text{th}}$  power and that the image of an element which is not a  $q^{\text{th}}$  power ( $q \neq \lambda_k$  odd prime) is still not a  $q^{\text{th}}$  power.

The endomorphism  $\Phi^{\epsilon}$  of  $H$  corresponding to the lifts  $\Phi_k$  with  $k \notin S$  has the description

$$(a, b, c)\Phi^{\epsilon} = (\lambda_k \cdot a, \lambda_k \cdot b, (\lambda_k)^2 \cdot c + a).$$

If  $q \neq \lambda_k$  is an odd prime then the image under  $\Phi^{\epsilon}$  of an element which is not a  $q^{\text{th}}$  power is still not a  $q^{\text{th}}$  power. Moreover, if  $(a, b, c)\Phi^{\epsilon}$  is a  $\lambda_k^{\text{th}}$  power then  $\lambda_k \mid a$ .

4.8. Now let  $S \subseteq \mathbf{N}$  and construct the group  $G_{\Phi(S)} = \text{colim } \Phi(S)$ . Then the nilpotent group  $N(S) = G_{\Phi(S)} / (G_{\Phi(S)})_3$  is the direct limit of the obvious chain

$${}_1H \xrightarrow{\Phi_1^*} {}_2H \xrightarrow{\Phi_2^*} {}_3H \xrightarrow{\Phi_3^*} \dots$$

where each  ${}_kH$  is isomorphic with the free nilpotent group  $H$  discussed above. The isolators  $I(n) = \{n' \in N(S) \mid n = (n')^j \text{ some } j \in \mathbf{Z}\}$  of an element  $n \in N(S)$  are of two types: if  $n$  stems from an element  $(a_k, b_k, c_k) \in {}_kH$  with  $a_k \neq 0$ , – note the choice of  $k$  does not matter – then  $I(n) \cong \text{gp}\{1/p \mid p \in \lambda(S)\}$ , otherwise  $I(n) \cong \text{gp}\{1/p \mid p \in P\}$ . The claim then follows from the classification of isomorphism types of subgroups of the rationals.

*Acknowledgment.* I would like to thank F. R. Beyl for some helpful discussions in connection with the last counter-example.

## REFERENCES

- [1] BAUMSLAG, G., *Lecture Notes on Nilpotent Groups*; Regional Conference Series in Mathematics, Vol. 2; Amer. Math. Soc. 1971.
- [2] BOURBAKI, N., *Groupes et Algèbres de Lie, Chapitres 2 et 3*; Éléments de Mathématique, Fascicule XXXVII; Actualités Sc. et Ind. 1349, Hermann 1972.
- [3] LAZARD, M., *Sur les algèbres enveloppantes universelles de certaines algèbres de Lie*; Publ. Sci. Univ. Alger (A)I (1954), 281–294.



- [4] MAGNUS, W., *Über Beziehungen zwischen höheren Kommutatoren*; J. reine angew. Math. 177 (1937), 105–115.
- [5] MALCEV, A. I., *Two remarks on nilpotent groups*; Math. Sb. (N.S.) 37 (79) (1955), 567–572.
- [6] NEUMANN, H., *Varieties of Groups*; Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 37; Springer 1967.
- [7] SERRE, J.-P., *Lie algebras and Lie groups*; Benjamin 1965.
- [8] STALLINGS, J., *Homology and central series of groups*; J. Algebra 2 (1965), 170–181.
- [9] STAMMBACH, U., *Anwendungen der Homologietheorie der Gruppen auf Zentralreihen und auf Invarianten von Präsentierungen*; Math. Z. 94 (1966), 157–177.
- [10] STREBEL, R., *Homological methods applied to the derived series of groups*; Comment. Math. Helv. 49 (1974), 302–332.
- [11] WITT, E., *Treue Darstellung Liescher Ringe*; J. reine angew. Math. 177 (1937), 152–160.

*Mathematisches Institut  
Universität Heidelberg  
Im Neuenheimer Feld 288  
D 6900 Heidelberg  
West Germany*

Received February 20, 1978