

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 54 (1979)

Artikel: On division by inner factors.
Autor: Anderson, J.M.
DOI: <https://doi.org/10.5169/seals-41578>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On division by inner factors

by J. M. ANDERSON

Professor Albert Pfluger Zugeeignet

§1. Introduction

Let G denote a Banach space of function $f(z)$ analytic in the open disc $|z| < 1$, and suppose that G is contained within the Hardy space H^1 . If $h \in H^1$ we may write $h(z) = 0(z) \cdot I(z)$ where $0(z)$ is the outer factor of h , $0(z) \in H^1$ and $I(z)$ is the inner factor, which is a function in H^∞ for which $\lim_{r \rightarrow 1} f(re^{i\theta})$ exists and is of modulus one almost everywhere on $|z| = 1$. Such an inner factor may be written as

$$I(z) = B(z) \cdot S(z),$$

where $B(z)$ is the Blaschke product formed from the zeros of $h(z)$, and $S(z)$ is a singular inner factor of the form

$$S(z) = \exp \left\{ -\frac{1}{2\pi} \int_0^{2\pi} \frac{e^{ix} + z}{e^{ix} - z} d\mu(x) \right\}. \tag{1}$$

Here, $\mu(x)$ is a monotonic increasing function on $(0, 2\pi]$ which is singular with respect to Lebesgue measure on $(0, 2\pi]$. For the details of such a factorization and other properties of H^p spaces we refer to [3].

DEFINITION 1. A Banach space G , contained within H^1 is said to have the f -property if, given any $h \in G$ and any inner function I which divides h (i.e. $hI^{-1} \in H^1$) then $hI^{-1} \in G$.

This notation was introduced by Havin [6], reporting on earlier work by numerous authors [2], [9], [10], [11]; see also [15] and the lecture by Kahane [8]. In addition to the obvious spaces H^p ($p > 1$) many other spaces, such as various Lipschitz spaces, have the f -property.

If $\phi(z) \in H^\infty$ and $h(z) \in H^2$ we may write

$$\phi(e^{i\theta})h(e^{i\theta}) \sim \sum_{-\infty}^{\infty} a_n e^{in\theta}$$

and define the projection operator $P(\bar{\phi}h)$ by

$$P(\bar{\phi}h) \sim \sum_0^{\infty} a_n e^{in\theta}.$$

(i.e. P is the Toeplitz operator associated with $\bar{\phi} \in L^\infty$). Clearly

$$P(\bar{\phi}h)(z) = \frac{1}{2\pi i} \int_{|w|=1} \frac{\bar{\phi}(w)h(w)}{w-z} dw, \quad |z| < 1.$$

DEFINITION 2. A Banach space G contained in H^1 is said to have the K -property if given any $h \in G$ and $\phi \in H^\infty$ then $P(\bar{\phi}h) \in G$.

This property appears to have been considered first by Korenblum [9]. Its usefulness consists in the fact that if ϕ is an inner function dividing $h \in G$ then $P(\bar{\phi}h) = h\phi^{-1}$. Thus a space having the K -property also possesses the f -property.

The first example of a space not possessing the f -property and hence not possessing the K -property was given by Gurarii [5]. The space in question is l^1 (called W^\dagger by Gurarii), defined by

$$l^1 = \left\{ f : f(z) = \sum_0^{\infty} f_n z^n, \|f\| = \sum_0^{\infty} |f_n| < \infty \right\}.$$

The particular function considered is

$$h(z) = \frac{\pi}{2} \Gamma^2\left(\frac{3}{8}\right) + \sum_{k=0}^{\infty} \frac{(-1)^{k+1} \Gamma(k + \frac{3}{8})}{(k + \frac{3}{8}) \Gamma(k + \frac{5}{8})} z^{(8k+3)^2}.$$

This function clearly belongs to l^1 but if we write

$$f(z) = \exp \left\{ \gamma \frac{1+z}{1-z} \right\} \cdot h(z),$$

where $\gamma = 2^{-9} \pi^2$ then $f \in H^1$ but $f \notin l^1$, ([5] p. 29).

The main difficulty in Gurarii's proof is in detecting the presence of the "jump" inner factor $\exp\{-\gamma \frac{1+z}{1-z}\}$ in $h(z)$. Gurarii does not assert that the resulting function $f(z)$ is outer—there might be other inner factors. It seems very difficult in general to detect the presence of singular inner factors of the form (1) when the representing function $\mu(t)$ is continuous.

§2. Results

In this paper we present another example of a subspace X of H^1 which does not have the f -property. We construct a function $F \in X$ such that

$$F(z) = \Phi(z) \cdot S(z),$$

where $\Phi(z)$ is an outer function not in X and $S(z)$ is an inner function of the form (1) with $\mu(t)$ a continuous, monotonic, increasing function of t , which is singular with respect to Lebesgue measure on $(0, 2\pi]$. This direct construction avoids the difficulty of showing the presence of an inner factor.

We denote by B the Banach space of Bloch functions, i.e. functions $f(z)$ analytic in $|z| < 1$ for which the norm

$$\|f\| = |f(0)| + \sup_{|z| < 1} (1 - |z|^2) |f'(z)|$$

is finite. All polynomials belong to B and the closure of the polynomials in the Bloch norm is denoted by B_0 . Alternatively, B_0 consists of those $f \in B$ for which

$$(1 - |z|^2)f'(z) \rightarrow 0 \quad \text{as} \quad |z| \rightarrow 1-.$$

It is evident that $H^\infty \subset B$, but $H^\infty \not\subset B_0$ —for these and other properties of Bloch functions we refer to [1]. The space X we wish to consider is

$$X = H^\infty \cap B_0.$$

It is easy to verify that if f and g belong to X so does $f \cdot g$ so that, with a suitable choice of norm, X is a subalgebra of H^∞ .

THEOREM 1. *X does not have the f -property (and so does not have the K -property).*

There is an interesting characterization of X which appears to be due, in the first instance, to M. Behrens. Let M denote the maximal ideal space of H^∞ and let \hat{f} denote the transform of f in the standard Gelfand representation. Then $H^\infty \cap B_0$ consists precisely of all those $f \in H^\infty$ such that f is constant on all the non-trivial Gleason parts of M other than the image of the disc $|z| < 1$. The parts are, of course, either single points or analytic discs, and the proof is a simple consequence of work of Hoffman [7]. The space X is thus a closed subalgebra of H^∞ , a fact which can also be verified directly.

§3. Interference

To construct the function in X whose outer part (which belongs to H^∞ and so to B) does not belong to B_0 , we require the following theorem of Shapiro [14 Theorem 4] regarding interference.

THEOREM A. *Let $\phi(x)$ be any increasing function on $[0, 2\pi]$ having continuous second derivative on the open interval $(0, 2\pi)$. Then there exists a function $f(x)$ continuous on $[0, 2\pi]$ such that*

$$\mu(x) = \phi(x) + f(x) \tag{2}$$

is increasing and singular,

$$|f(x+t) + f(x-t) - 2f(x)| < Kt|\log t|^{-\frac{1}{2}}, \tag{3}$$

for some absolute constant K and all x, t satisfying $0 < t < \frac{1}{2}, 0 \leq x-t < x+t \leq 2\pi$.

Clearly $\mu(x)$ is continuous. As regards the function f , it is sufficient for our purposes to note that (3) implies that f belongs to the Zygmund class λ_* . Thus, if we define the function $F(z)$ by

$$F(z) = \exp \left\{ - \int_0^{2\pi} \frac{e^{ix} + z}{e^{ix} - z} df(x) \right\}, \tag{4}$$

then

$$F'(z) = O(1 - |z|^2)^{-1}, \quad |z| \rightarrow 1-,$$

and so $F \in B_0$. (See [4] Theorem 1 for details.)

It is convenient to base the proof of Theorem 1 on another theorem, which seems of independent interest.

THEOREM 2. *There exists a monotonic increasing, infinitely differentiable function $\phi(x)$ defined for $0 < x < 2\pi$ such that the function*

$$\Phi(z) = \exp \left\{ + \int_0^{2\pi} \frac{e^{ix} + z}{e^{ix} - z} d\phi(x) \right\} \tag{5}$$

belongs to H^∞ , but not to B_0 .

Proof of Theorem 1. Let $\phi(x)$ be as in Theorem 2 and $f(x)$ as in Theorem A. Then $\mu(x) = \phi(x) + f(x)$, or $f(x) = \mu(x) + (-\phi(x))$ with $\mu(x)$ singular and $-\phi(x)$ absolutely continuous with respect to Lebesgue measure. Thus

$$\begin{aligned} F(z) &= \exp \left\{ - \int_0^{2\pi} \frac{e^{ix} + z}{e^{ix} - z} df(x) \right\} \\ &= \exp \left\{ + \int_0^{2\pi} \frac{e^{ix} + z}{e^{ix} - z} d\phi(x) \right\} \cdot \exp \left\{ - \int_0^{2\pi} \frac{e^{ix} + z}{e^{ix} - z} d\mu(x) \right\} \\ &= \Phi(z) \cdot S(z). \end{aligned}$$

Since $\Phi(z)$ and $S(z)$ belong to H^∞ so does $F(z)$, and $\Phi(z)$ is the outer part of $F(z)$. Thus $F(z) \in H^\infty \cap B_0 = X$ by (3) but, by construction, $\Phi(z)$ does not belong to B_0 . Thus X does not have the f -property.

§4. Proof of Theorem 2

We construct the required function $\phi(x)$ satisfying $\phi'(x) \leq 1$ for all x . This will imply that $|\Phi(z)| \leq e^{2\pi}$ for $|z| \leq 1$ and so $\Phi(z) \in H^\infty$. The standard example of a function in H^∞ but not in B_0 is the inner function $\exp\{-1 + z/1 - z\}$ which is of the form (1) with $\mu(x)$ discontinuous. Our construction is based on the modified function

$$\Phi(\delta, z) = \exp \left\{ \int_{-\delta}^{\delta} \frac{e^{ix} + z}{e^{ix} - z} d(x + \delta) \right\}.$$

An integration yields

$$\Phi(\delta, z) = \exp \left\{ -2\delta - 2i \log \frac{e^{i\delta} - z}{e^{-i\delta} - z} \right\}.$$

We now consider z_0 real and equal to $1 - \delta$. Then

$$(1 - |z_0|^2)\Phi'(\delta, z_0) = \delta(2 - \delta) e^{-2\delta} \exp \left[-2i \log \frac{e^{i\delta} - z_0}{e^{-i\delta} - z_0} \right] 2i \left(\frac{1}{e^{i\delta} - z_0} - \frac{1}{e^{-i\delta} - z_0} \right)$$

An easy calculation now shows that

$$\begin{aligned} \lim_{\delta \rightarrow 0} (1 - |z_0|^2)\Phi'(\delta, z_0) &= -4 \exp \left\{ -2i \log \frac{1+i}{1-i} \right\} \\ &= -4 e^\pi. \end{aligned}$$

We now define sequences $\{\delta_n\}_{n=1}^\infty$ and $\{\theta_n\}_{n=1}^\infty$ so that the function

$$\Phi(z) = \prod_{n=1}^{\infty} \Phi(\delta_n, ze^{i\theta_n})$$

has the required properties. We choose $\theta_1 = 0$ and $\theta_n = \pi(1 - 1/n)$, $n > 1$. We then choose each δ_n so small so that, with $z_n = (1 - \delta_n) e^{-i\theta_n}$,

$$(1 - |z_n|^2)\Phi^1(\delta_n, z_n) \approx -4 e^\pi$$

but

$$|\Phi(\delta_n, z_m)| \approx 1 \quad (m \neq n).$$

It is clear that this can be achieved simply by choosing the sequence $\{\delta_n\}$ sufficiently small. Thus for our function $\Phi(z)$ we have

$$\lim_{n \rightarrow \infty} (1 - |z_n|^2) \Phi'(z_n) > 0,$$

whereas $|z_n| \rightarrow 1$ as $n \rightarrow \infty$. Thus $\Phi(z) \notin B_0$.

If $\Phi(z)$ as constructed above has the representation (5), the representing function $\phi(x)$ will not be differentiable at the points $x = \theta_n + \delta_n$, $n = 1, 2, 3, \dots$. Nevertheless by adjusting the function $\phi(x)$ in intervals $[\theta_n - \delta_n - \epsilon_n, \theta_n - \delta_n + \epsilon_n]$ and $[\theta_n + \delta_n - \epsilon_n, \theta_n + \delta_n + \epsilon_n]$ where ϵ_n is now chosen very small compared with δ_n we may make the function $\Phi(x)$ as smooth as we please. Thus Theorem 2 is proved.

§5. Analytic VMO

We denote by BMOA the Banach space of functions $f(z)$ analytic for $|z| < 1$ for which the norm

$$\|f\| = |f(0)| + \sup_{|\xi| < 1} \|f_\xi\|_2 < \infty.$$

Here

$$(\|f_\xi\|_2)^2 = \frac{1}{2\pi} \int_0^{2\pi} \left| f\left(\frac{z+\xi}{1+\xi z}\right) - f(\xi) \right|^2 |dz|, \quad z = e^{i\theta}.$$

The subspace of BMOA for which

$$\|f_\xi\|_2 \rightarrow 0 \quad \text{as} \quad |\xi| \rightarrow 1- \tag{6}$$

is denoted by VMOA. A function belongs to VMOA if and only if its boundary values on $|z|=1$ belong to the space of functions of vanishing mean oscillation as defined, for example in [13]. This characterization of VMOA was introduced in [12] where it was shown that $f \in \text{VMOA} \Rightarrow f \in B_0$.

From the above characterization two interesting facts follow immediately:

(i) No inner function in H^∞ can belong to VMOA. To see this we note that if f is inner then there is a sequence $\{\xi_n\}$ with $|\xi_n| \rightarrow 1$ as $n \rightarrow \infty$ such that $f(\xi_n) \rightarrow 0$ as $n \rightarrow \infty$. Since

$$\frac{1}{2\pi} \int_0^{2\pi} \left| f\left(\frac{z+\xi}{1+\xi z}\right) \right|^2 d\theta = 1$$

for all ξ with $|\xi| < 1$ it follows that (6) does not hold.

(ii) If $Y = H^\infty \cap \text{VMOA}$ then $f, g \in Y$ implies $f \cdot g \in Y$. Thus Y , with suitable choice of norm can be made into a Banach algebra. The space Y is the ‘‘analytic’’ part of the space QC of quasicontinuous functions introduced by Sarason [13]. Since it was observed by Sarason that $QC = (H^\infty + C) \cap (H^\infty + C)$, we see that $Y = H^\infty \cap (H^\infty + C)$. In particular we see that Y is again a closed subalgebra of H^∞ , a fact which can also be verified directly.

If A is defined by

$$A = \{f(z) : f(z) \text{ analytic for } |z| < 1, \text{ continuous for } |z| \leq 1\}.$$

then we have that $A \subset Y \subset X \subset H^\infty$. It is easy to see that all of the above inclusions are strict. An example of a function in X but not in Y is a function of the form (4) where $f(x)$ is a monotonic singular function satisfying (3). That such functions exist is known from [14, Theorem 2]. An example of a function in Y but not in A is a function $f(z)$ mapping $|z| < 1$ conformally onto a bounded domain which has a bad prime end. Clearly $f \notin A$ but since f is univalent and in B_0 it is in VMOA ([12] Satz 1).

It would be nice to find a characterization of $Y = H^\infty$ VMOA similar to that of X mentioned in §2. Even though no inner function can belong to Y there are functions in Y which have an inner factor—the function constructed by Gurarii is in A and has a jump inner factor.

In conclusion it seems worthwhile to point out the following theorem

THEOREM 3. *Let $F(z)$ be given by (4) where $f(x)$ satisfies (3). Then there is a constant K_1 so that*

$$|F'(z)| < \frac{K_1}{1-|z|^2} \left[\log \left(\frac{1}{1-|z|^2} \right) \right]^{-\frac{1}{2}}$$

for $|z| < 1$.

For our purposes we needed only that $F'(z) = O(1-|z|^2)^{-1}$, $|z| \rightarrow 1-$. The proof of this theorem is a straightforward adaptation of the proofs of ([4] Theorem 1) and ([16] Theorem 7), and-so is omitted.

I am indebted to Professors J. Clunie and R. Rochberg for their help and advice in the preparation of this paper.

REFERENCES

1. ANDERSON, J. M. CLUNIE J. and Ch. POMMERENKE, *On Bloch functions and normal functions* J. Reine Angew. Math. 270 (1974) 12–37.
2. CARLESON, L., *A representation formula for the Dirichlet Integral* Math. Zeit. 73 (1960) 190–196.
3. DUREN, P. L., *Theory of H^p Spaces*, New York, 1970.
4. DUREN, P. L., SHAPIRO, H. S. and SHIELDS, A. L., *Singular measures and domains not of Smirnov type*, Duke Math. J. 33, (1966) 247–254.
5. GURARII, V. P., *On the factorization of absolutely convergent Taylor series and Fourier Integrals* (in Russian). Zap. Nauch. Sem. LOMI. 30 (1972), 15–32.
6. HAVIN, V. P., *On the factorization of analytic functions, smooth up to the boundary* (in Russian), Zap. Nauch. Sem. LOMI. 22 (1971) 202–205.
7. HOFFMAN, K., *Bounded analytic functions and Gleason parts*, Ann. of Math, 86, (1967) 74–111.
8. KAHANE, J. P., *Two problems on H^p classes* Symposium on Complex Analysis, Canterbury 1973, London Math. Soc. Lecture Note Series, 12, Cambridge, 1975, 93–95.

9. KORENBLYUM, B. I., *On an extremal property of outer functions* (in Russian) *Mat. Zamet.* 10 (1971), 53–56.
10. KORENBLYUM B. I. and KOROLEVICH, V. S., *On analytic functions, regular in the disc and smooth on its boundary* (in Russian) *Mat. Zamet.* 7 (1970) 165–172.
11. KORENBLYUM B. I. and FAIVYSHEVSKII, V. M., *On a class of projection operators connected with the division of analytic functions*, (in Russian), *Ukrain. Mat. Zhurnal*, 24 (1972) 692–695.
12. POMMERENKE, Ch., *Schlichte Funktionen und analytische Funktionen beschränkter mittlerer Oszillation* *Comment. Math. Helv.* 52 (1977) 591–602.
13. SARASON, D., *Functions of vanishing mean oscillation* *Trans. Amer. Math. Soc.* 207 (1975) 391–405.
14. SHAPIRO, H. S., *Monotonic singular functions of high smoothness*, *Michigan Math. J.* 15 (1968) 265–275.
15. SHIROKOV, N. A., “*Ideal and factorisation in algebras of functions smooth up to the boundary* (in Russian)” *Trudy Math. Inst. imeni V. A. Steklova* 130 (1978) 196–220.
16. ZYGMUND, A., *Smooth functions* *Duke Math. J.* 12 (1945) 47–76.

*Mathematics Department,
University College London
London WC1*

Received November 15, 1977.