

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 54 (1979)

Artikel: A Landesman-Lazer alternative theorem for a class of optimization problems.
Autor: Frehse, Jens
DOI: <https://doi.org/10.5169/seals-41580>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A Landesman–Lazer alternative theorem for a class of optimization problems

JENS FREHSE

In [1] we proved an alternative theorem for the existence of minima of functionals F_l defined by $F_l(u) = F(u) - (l, u)$ on, say, a reflexive Banach space B . Here $l \in B^*$ and F satisfies the following conditions

- (1) $F: B \rightarrow \mathbf{R} \cup \{\infty\}$ is lower semi-continuous in the weak topology of B .
- (2) F is of polynomial type, i.e. if for some pair $v, w \in B$
 - (i) $\sup t^{-1}F(w + tv) < \infty \quad (t > t_0 > 0)$ and
 - (ii) $\inf F(w + tv) > -\infty \quad (t \in \mathbf{R})$

then $F(w + tv)$ is constant in t .

- (3) F satisfies a surrogate convexity condition, i.e. for all $u, w \in B, \alpha \in [0, 1]$, $F(w) < \infty$

$$F(1 - \alpha)w + \alpha u \leq K_w + K_w \alpha F(u)$$

with some constant K_w .

- (4) F is semi-coercive, i.e. there exists a continuous projection $Q: B \rightarrow V$ onto a finite dimensional subspace V such that for all $K > K_0$

$$\sup \{ \|u\| / (1 + \|Qu\|) \mid F(u) \leq K \} < \infty$$

- (5) F is bounded from below on B .

Introducing the set

$$D = \{v \in B \mid F(w + tv) \text{ is constant in } t \in \mathbf{R} \text{ for all } w \in B, F(w) < \infty\}$$

we obtained the following

THEOREM 0. Under the conditions (1)–(5) the functional F_l has a minimum if and only if $l \perp D$. Furthermore, $\dim D \leq \dim V$.

A simple corollary (cf. [2], §1) yields that if in addition F has a Gateaux derivative $T: B \rightarrow B^*$ then the range of T is linear and has finite co-dimension.

In this paper we consider perturbations $F_l + G$ of the above functionals F_l where G has a so called *weak sub-asymptote* (cf. definition below). It then turns out that the set of $l \in B^*$ for which $F_l + G$ has a minimum becomes “thicker”, i.e. is open and contains the closed set D . Under additional conditions we can characterize these elements l in the form of an alternative theorem. Our results are in the spirit of the “classical” Landesman–Lazer-alternative theorems, cf. Landesman–Lazer [8]. For a rather complete list of references to this subject, cf. [9] and also [4]. These theorems state that the range of perturbed semi-coercive linear differential operators like $-\Delta u - \lambda u + \arctg u$ subject to boundary conditions is open and can be characterized by the asymptotes of the perturbation.

If $F_l + G$ is Gateaux-differentiable our result yields a Landesman–Lazer-type theorem of the usual form but covers cases with strongly non-linear principal part of polynomial type. Our proof of this result is very simple. The theorem has a non-variational analogue which was presented in [3]. A Landesman–Lazer theorem for a class of equations with strongly non-linear principle part was presented by Hess in [5], [6], [7]. His approach and his results are rather different from the setting in the present paper.

DEFINITION. A mapping $a_0: D \rightarrow \mathbf{R}$ is called a weak sub-asymptote of the mapping $G: B \rightarrow \mathbf{R}$ if for every sequence $(u_i \in B, i = 1, \dots)$ with $\|u_i\| \rightarrow \infty$, $\|u_i\|^{-1}u_i \rightarrow v \in D$ weakly ($i \rightarrow \infty$), $v \neq 0$, we have

$$a_0(v) \leq \liminf \|u_i\|^{-1} G(u_i) < \infty \quad (i \rightarrow \infty) \quad (6)$$

We shall also assume for the perturbation G

$$\sup \|u\|^{-1} G(u) < \infty \quad (u \in B, u \neq 0) \quad (7)$$

and

$$F + G \text{ is lower semi-continuous in the weak topology of } B. \quad (8)$$

THEOREM 1. Let B be a reflexive Banach space and $F: B \rightarrow \mathbf{R} \cup \infty$, $G: B \rightarrow \mathbf{R}$ mappings such that G has a weak sub-asymptote $a_0: D \rightarrow \mathbf{R}$ and F, G satisfy (1)–(5), (7) and (8).

Then the functional $\Phi_l: B \rightarrow \mathbf{R}$ defined by

$$\Phi_l(u) = F(u) + G(u) - (l, u)$$

has a minimum on B for all $l \in B^*$ for which

$$(l, v) < a_0(v), \quad v \in D, \quad v \neq 0. \quad (9)$$

If in addition

$$\liminf t^{-1}(G(w + tv) - G(w)) < a_0(v) \quad (t \rightarrow +0) \quad (10)$$

for all $w \in B$ and $v \in D, v \neq 0$, then (9) is also necessary for the existence of a minimum of Φ_l .

Proof. We may assume that $F \neq \text{const} = \infty$. Let (u_i) be a minimizing sequence for Φ_l . Suppose that (u_i) were unbounded. Then we may assume that $\|u_i\| \rightarrow \infty$ and that $\|u_i\|^{-1}u_i := v_i \rightarrow v$ weakly in $B (i \rightarrow \infty)$. By (4) there is a constant $C > 0$ such that $\|u_i\| \leq C + C\|Qu_i\|$ and hence $1 \leq C \liminf \|Qv_i\| (i \rightarrow \infty)$. Since $\dim QB < \infty$ we have $Qv_i \rightarrow Qv$ strongly and, therefore, $1 \leq C\|Qv\|$, and

$$v \neq 0.$$

We intend to show that $v \in D$. By the convexity condition (3)

$$F(1 - \alpha)w + \alpha u_i \leq K_w + \alpha K_w F(u_i) = K_w + \alpha K_w [\Phi_l(u_i) - G(u_i) + (l, u_i)]$$

for all w such that $F(w) < \infty$.

Since $\|u_i\| \rightarrow \infty$ we may set $\alpha = t\|u_i\|^{-1}$ for $t > 0, i > i(t)$. Passing to the limit $i \rightarrow \infty$ we obtain in view of the lower semi-continuity of F .

$$F(w + tv) \leq K_w - tK_w \liminf \|u_i\|^{-1}G(u_i) + tK_w(l, v) \quad (11)$$

and by (7)

$$t^{-1}F(w + tv) \leq K_{wt}, \quad t > 1.$$

From (5) and (2) we then conclude that $F(w + tv)$ is constant in $t \in \mathbf{R}$ for all w with $F(w) < \infty$ and hence

$$v \in D.$$

Since G has a weak sub-asymptote $a_0: D \rightarrow \mathbf{R}$, we obtain from (11)

$$(F(w) =)F(w + tv) \leq K_w - tK_w a_0(v) + tK_w(l, v)$$

and passing to the limit $t \rightarrow \infty$ we arrive at the inequality

$$a_0(v) \leq (l, v).$$

This contradicts (9) and hence the assumption of (u_i) being unbounded leads to a contradiction. The first statement follows in view of (8) and the reflexivity of B . The necessity of (9) can be seen from the following simple argument: If u is a minimum of Φ_t on B and if $v \in D$, then

$$F(u) + G(u) - (l, u) \leq F(u + tv) + G(u + tv) - (l, u + tv)$$

$$G(u) \leq G(u + tv) - t(l, v)$$

since $F(u + tv)$ is constant in t . Hence, by (10),

$$(l, v) \leq \liminf t^{-1}(G(u + tv) - G(u)) < a_0(v) \quad (t \rightarrow +0)$$

as claimed. The theorem is proved.

EXAMPLES. In the following let Ω be a bounded connected open set in \mathbf{R}^n and $H^{1,p}$ the usual Sobolev space over Ω . The corresponding Sobolev space of r -vector functions is denoted by $[H^{1,p}]^r$.

(i) Let $B = H^{1,p}(\Omega)$, $l \in B^*$, and

$$F(u) = \frac{1}{p} \int |\nabla u|^p dx, \quad G(u) = \int [u \operatorname{arctg} u - \frac{1}{2} \ln(1 + |u|^2)] dx$$

Here and in the following \int denotes integration over Ω . Then $D = \{z \in H^{1,p} \mid z = \text{const.}\}$ and $a_0(v) = \pi/2 \int |v| dx$. Since F and G satisfy the hypotheses of Theorem 1, cf [1], §3, the functional F_t defined by

$$F_t(u) = F(u) + G(u) - (l, u)$$

has a minimum when

$$|(l, 1)| < \frac{\pi}{2} |\Omega|. \tag{12}$$

The minimum u of F_l is a weak solution of the differential equation.

$$-\sum_{i=1}^n \partial_i (\partial_i u |\nabla u|^{p-2}) + \operatorname{arctg} u = l. \quad (13)$$

It is a simple exercise to prove that the above functional G satisfies condition (10); Hence (12) is also necessary for the existence of a minimum of F_l . The characterization of the range of the operator on the left hand side of (13) can be obtained using the methods of Peter Hess, cf. references.

A non-differentiable variant of this example is obtained when $F(u)$ is replaced by

$$F(u) = \int \left[\frac{1}{p} |\nabla u|^p + |\nabla u| \right] dx$$

and condition (12) remains as the necessary and sufficient condition for the existence of minima of the functional F_l .

(ii) Let $B = [H^{1,p}(\Omega)]^2$, $l \in B^*$ and

$$F(u) = \int [|\nabla u_1|^p + u_1^p + \lambda \sin u_1 + u_1 \partial_1 u_2 + |\nabla u_2|^p] dx, \quad u = (u_1, u_2),$$

$$G(u) = \int [u_2 \operatorname{arctg} u_2 - \frac{1}{2} \ln(1 + |u_2|^2)] dx.$$

Again, F and G satisfy the hypotheses of the theorem. The surrogate convexity of F follows by splitting the integrand into a sum of convex and bounded functions. The set D has the form

$$D = \{(0, c) \in [H^{1,p}(\Omega)]^2 \mid c \in \mathbf{R}\}.$$

The functional F_l , $l = (l_1, l_2) \in B^*$ has a minimum if and only if

$$|\langle l_2, 1 \rangle| < \frac{\pi}{2} |\Omega|.$$

REFERENCES

- [1] FREHSE, J., *An existence theorem for a class of non-coercive optimization and variational problems.* Math. Z. 159 (1978), 51–63.
- [2] FREHSE, J., *Existence and alternative theorems for semi-coercive problems and applications to*

non-linear elliptic equations. From: Quaderni dei gruppi di ricerca matematica del consiglio nazionale delle ricerche. Convegno su: Sistemi ellittici non lineari ed applicazioni. Ferrara, 20–24 Sett. 1977. Istituto matematico dell'Università di Ferrara. Editrice Universitaria, Ferrara 1978.

- [3] FREHSE, J., *Landesman–Lazer-alternative theorems for a class of non-linear functional equations*. Math. Ann. 238 (1978), 59–65.
- [4] FUČIK, S., *Ranges of nonlinear operators*. Vol. I–V. Dept. of Mathematics Charles University Praha. to be published.
- [5] HESS, P., *On a class of strongly nonlinear elliptic variational inequalities*. Math. Ann. 211 (1974), 289–297.
- [6] HESS, P., *On semi-coercive nonlinear problems*. Indiana Univ. Math. J. 23 (1974), 645–654.
- [7] HESS, P., *Nonlinear perturbations of linear elliptic and parabolic problems at resonance: Existence of multiple solutions*. To appear in Ann. Sc. Norm. Sup. Pisa. iv, vol. 5, (1978), 527–538.
- [8] LANDESMAN, E. M., and LAZER, A. C., *Nonlinear perturbations of linear elliptic boundary value problems at resonance*. J. Math. Mech. 19 (1970), 609–623.
- [9] MAWHIN, J., *Landesman–Lazer's type problems for nonlinear equations*. Conference del Seminario di Matematica dell'Università di Bari. No. 147 (1977).

*Institut für Angewandte Mathematik
der Universität, Beringstraße 4–6
5300 Bonn, West-Germany*

Received June 26, 1978