

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 54 (1979)

Artikel: The Boolean algebra of spectra.
Autor: Bousfield, A.K.
DOI: <https://doi.org/10.5169/seals-41584>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Boolean algebra of spectra

A. K. BOUSFIELD

Introduction

Let \mathbf{Ho}^s denote the stable homotopy category of CW-spectra (cf. [Adams 2]), and for $E \in \mathbf{Ho}^s$ let E_* be the associated homology theory. For $E, G \in \mathbf{Ho}^s$ we say E_* and G_* have the same acyclic spectra if the following equivalent conditions hold:

- (i) For $X \in \mathbf{Ho}^s$, $E_*X = 0 \Leftrightarrow G_*X = 0$.
- (ii) For $f: X \rightarrow Y \in \mathbf{Ho}^s$, $f_*: E_*X \approx E_*Y \Leftrightarrow f_*: G_*X \approx G_*Y$.

This gives a very coarse equivalence relation for spectra, and we let $\mathbf{A}(\mathbf{Ho}^s)$ consist of all the equivalence classes $\langle E \rangle$ for $E \in \mathbf{Ho}^s$, where $\langle E \rangle$ is given by all $G \in \mathbf{Ho}^s$ such that E_* and G_* have the same acyclic spectra. We partially order $\mathbf{A}(\mathbf{Ho}^s)$ by writing $\langle E \rangle \leq \langle G \rangle$ when each G_* -acyclic spectrum is E_* -acyclic. Our purpose in this note is to study the algebraic structure of $\mathbf{A}(\mathbf{Ho}^s)$ when it is equipped with the relation \leq and the operations \vee and \wedge induced from the usual wedge and smash product for spectra.

We say that $\langle E \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ has a *complement* $\langle E \rangle^c \in \mathbf{A}(\mathbf{Ho}^s)$ if $\langle E \rangle \wedge \langle E \rangle^c = \langle 0 \rangle$ and $\langle E \rangle \vee \langle E \rangle^c = \langle S \rangle$ where S is the sphere spectrum, and we note that $\langle E \rangle^c$ is unique when it exists. We let $\mathbf{BA}(\mathbf{Ho}^s) \subset \mathbf{A}(\mathbf{Ho}^s)$ consist of those $\langle E \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ with complements, and we observe that $\mathbf{BA}(\mathbf{Ho}^s)$ is a Boolean algebra. We prove that $\langle E \rangle \in \mathbf{BA}(\mathbf{Ho}^s)$ whenever E is a (possibly infinite) wedge of finite CW-spectra. It would be most interesting to determine the sublattice of $\mathbf{BA}(\mathbf{Ho}^s)$ given by such $\langle E \rangle$. We show that $\langle S^0 \cup_\alpha e^n \rangle = \langle S^0 \rangle$ for each $\alpha \in [S^{n-1}, S^0]$ with $n \neq 1$, and that $\langle DE \rangle = \langle E \rangle$ when E is a finite CW-spectrum and DE is its Spanier-Whitehead dual. This incidentally implies that $G_*E = 0 \Leftrightarrow G^*E = 0$, for $G, E \in \mathbf{Ho}^s$ with E finite. Some other members of $\mathbf{BA}(\mathbf{Ho}^s)$ are $\langle K \rangle$ and $\langle SZ_{(J)} \rangle$ where K is the spectrum of complex K -theory and $SZ_{(J)}$ is the Moore spectrum associated with a subring $Z_{(J)} \subset Q$. Indeed, $\langle K \rangle$ and $\langle SZ_{(J)} \rangle$ are of the form $\langle E \rangle^c$ where E is an appropriate wedge of finite CW-spectra, though the proof for K will be postponed to [Bousfield 3].

We also introduce a distributive lattice $\mathbf{DL}(\mathbf{Ho}^s)$ given by all $\langle E \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ with $\langle E \rangle \wedge \langle E \rangle = \langle E \rangle$, and we show that $\mathbf{BA}(\mathbf{Ho}^s) \subset \mathbf{DL}(\mathbf{Ho}^s) \subset \mathbf{A}(\mathbf{Ho}^s)$ where both

containments are proper. It turns out that $\langle E \rangle \in \mathbf{DL}(\mathbf{Ho}^s)$ whenever E is a (possibly infinite) wedge of ring spectra and finite CW-spectra. In fact, most familiar spectra represent elements of $\mathbf{DL}(\mathbf{Ho}^s)$.

The class $\mathbf{A}(\mathbf{Ho}^s)$ has applications to the homological localization theory of spectra, cf. [Bousfield 3], [Ravenel]. In particular, the E_* -localization is equivalent to the G_* -localization iff $\langle E \rangle = \langle G \rangle$, and a determination of $\mathbf{A}(\mathbf{Ho}^s)$ would provide an inventory of the possible homological localization functors.

Our results on the structure of $\mathbf{A}(\mathbf{Ho}^s)$, $\mathbf{BA}(\mathbf{Ho}^s)$, and $\mathbf{DL}(\mathbf{Ho}^s)$ are established in §2. Some of our proofs involve $[E,]_*$ -colocalizations of spectra, and we develop the required theory in §1.

We essentially use the notation and terminology of [Adams 2]. However, we let \mathbf{Ho}^s be the category of CW-spectra and homotopy classes of maps of degree 0, cf. [Adams 2, p. 144]. Thus \mathbf{Ho}^s is an additive category equipped with an equivalence $\Sigma : \mathbf{Ho}^s \rightarrow \mathbf{Ho}^s$ induced by the “shift” suspension Σ of CW-spectra. We write $[X, Y]$ for the group of morphisms $X \rightarrow Y \in \mathbf{Ho}^s$, and write $[X, Y]_n$ for $[\Sigma^n X, Y]$ where $n \in \mathbb{Z}$. By a cofibre sequence we mean a sequence in \mathbf{Ho}^s equivalent to $X \xrightarrow{f} Y \xrightarrow{i} Y \cup_f CX$ for some cellular map f of degree 0 between CW-spectra, cf. [Adams 2, p. 155]. Recall that \mathbf{Ho}^s has arbitrary coproducts induced by the wedge \vee for CW-spectra, and for $X, Y \in \mathbf{Ho}^s$ there is a natural smash product $X \wedge Y \in \mathbf{Ho}^s$ which is associative, commutative, and unitary (with the sphere spectrum S as unit) up to coherent natural equivalences, cf. [Adams 2, p. 158]. We call $E \in \mathbf{Ho}^s$ a ring spectrum if it is equipped with an associative (but not necessarily commutative) multiplication $E \wedge E \rightarrow E$ and a two sided unit $S \rightarrow E$ in \mathbf{Ho}^s . As usual, we let $X * Y = \pi_* X \wedge Y = [S, X \wedge Y]_*$ for $X, Y \in \mathbf{Ho}^s$.

§1. $[E,]_*$ -colocalizations of spectra

In preparation for §2 and for [Bousfield 3], we now develop the $[E,]_*$ -colocalization theory of spectra. Some of the concepts here have previously been developed by J. P. May (unpublished) and in [Bousfield 2].

For $E \in \mathbf{Ho}^s$, a map $f : A \rightarrow B \in \mathbf{Ho}^s$ is called an $[E,]_*$ -equivalence if $f_* : [E, A]_* \approx [E, B]_*$, and a spectrum $C \in \mathbf{Ho}^s$ is called $[E,]_*$ -colocal if $g_* : [C, X]_* \approx [C, Y]_*$ whenever $g : X \rightarrow Y$ is an $[E,]_*$ -equivalence. It is easy to check:

- (1.1) E is $[E,]_*$ -colocal.
- (1.2) If $\{X_\alpha\}$ is a set of $[E,]_*$ -colocal spectra, then $\vee_\alpha X_\alpha$ is $[E,]_*$ -colocal.
- (1.3) If $W \rightarrow X \rightarrow Y$ is a cofibre sequence in \mathbf{Ho}^s and any two of W, X, Y are $[E,]_*$ -colocal, then so is the third.
- (1.4) If X is $[E,]_*$ -colocal, then so is $X \wedge Y$ for all $Y \in \mathbf{Ho}^s$.

A map $\varphi : X \rightarrow A \in \mathbf{Ho}^s$ is called an $[E,]_*$ -colocalization of A if X is $[E,]_*$ -colocal and φ is an $[E,]_*$ -equivalence. Note that the $[E,]_*$ -colocalizations of A are initial among the $[E,]_*$ -equivalences with target A , and are terminal among the maps from $[E,]_*$ -colocal spectra to A . $[E,]_*$ -colocalizations are clearly unique up to equivalence and

PROPOSITION 1.5. *Each spectrum $A \in \mathbf{Ho}^s$ has an $[E,]_*$ -colocalization.*

Proof. We inductively construct a transfinite sequence of inclusions of CW-spectra

$$A = B_0 \subset B_1 \subset \dots \subset B_s \subset B_{s+1} \subset \dots$$

where $B_\lambda = \bigcup_{s < \lambda} B_s$ for each limit ordinal λ and where $B_s \subset B_{s+1}$ is given by the push-out square

$$\begin{array}{ccc} \bigvee_{\alpha \in I} M_\alpha & \xrightarrow{f: M_\alpha \rightarrow B_s} & B_s \\ \downarrow & & \downarrow \\ \bigvee_{\alpha \in I} \text{Cone}(M_\alpha) & \xrightarrow{f: M_\alpha \rightarrow B_s} & B_{s+1} \end{array}$$

in which $\{M_\alpha\}_{\alpha \in I}$ consists of all cofinal subspectra of the spectra $\sum^n E$ for $n \in \mathbb{Z}$, and f ranges over all cellular functions $M_\alpha \rightarrow B_s$ of degree 0, cf. [Adams, p. 140, 154]. Now let σ be the number of stable cells in E and let γ be the first infinite ordinal of cardinality greater than σ . Then for each $\alpha \in I$, each cellular function $M_\alpha \rightarrow B_\gamma$ of degree 0 extends over $\text{Cone}(M_\alpha)$ because the image of M_α is contained in B_s for some $s < \gamma$. Thus $[E, B_\gamma]_* = 0$. Since A is a closed subspectrum of B_γ (cf. [Adams 2, p. 154]), there is an associated cofibre sequence

$$\sum^{-1}(B_\gamma/A) \rightarrow A \rightarrow B_\gamma$$

in \mathbf{Ho}^s . The morphism $\sum^{-1}(B_\gamma/A) \rightarrow A$ is clearly an $[E,]_*$ -equivalence, so it suffices to show $\sum^{-1}(B_\gamma/A)$ is $[E,]_*$ -colocal. For this it suffices to show inductively that B_s/A is $[E,]_*$ -colocal for all s . If B_s/A is $[E,]_*$ -colocal, then so is B_{s+1}/A because there is a cofibre sequence

$$B_s/A \rightarrow B_{s+1}/A \rightarrow B_{s+1}/B_s \in \mathbf{Ho}^s$$

where B_{s+1}/B_s is equivalent to a wedge of iterated (de)-suspensions of E . If B_s/A

is $[E,]_*$ -colocal for all $s < \lambda$ where λ is a limit ordinal, then B_λ/A is $[E,]_*$ -colocal because there is a cofibre sequence

$$\bigvee_{s < \lambda} B_s/A \xrightarrow{1-g} \bigvee_{s < \lambda} B_s/A \rightarrow B_\lambda/A \in \mathbf{Ho}^s$$

where g is induced by the maps $B_s/A \rightarrow B_{s+1}/A$. This completes the induction and the proof 1.5.

For each $A \in \mathbf{Ho}^s$ let $\varphi: {}^E A \rightarrow A \in \mathbf{Ho}^s$ denote the $[E,]_*$ -colocalization given by $\sum^{-1}(B_\gamma/A) \rightarrow A$ above, and note that it is functorial and idempotent in the obvious sense. To clarify the nature of $[E,]_*$ -colocal spectra, we let *Class-E* denote the smallest class of spectra in \mathbf{Ho}^s such that: (i) $E \in \text{Class-E}$; (ii) if $\{X_\alpha\}$ is a set of spectra in *Class-E*, then $\bigvee_\alpha X_\alpha \in \text{Class-E}$; and (iii) if $W \rightarrow X \rightarrow Y$ is a cofibre sequence in \mathbf{Ho}^s and any two of W, X, Y are in *Class-E*, then so is the third.

PROPOSITION 1.6. *Class-E equals the class of $[E,]_*$ -colocal spectra in \mathbf{Ho}^s .*

Proof. *Class-E* is contained in the class of $[E,]_*$ -colocal spectra by (1.1)–(1.3). Conversely, if X is $[E,]_*$ -colocal, then $X \in \text{Class-E}$ because ${}^E X \simeq X$ and ${}^E X \in \text{Class-E}$ by the proof of 1.5.

We call a spectrum $W \in \mathbf{Ho}^s$ $[E,]_*$ -trivial if $[E, W]_* = 0$, and we note that $[V, W]_* = 0$ whenever V is $[E,]_*$ -colocal and W is $[E,]_*$ -trivial. Each spectrum A can be canonically built from $[E,]_*$ -colocal and $[E,]_*$ -trivial spectra as follows. Extend $\varphi: {}^E A \rightarrow A$ to the cofibre sequence

$$(1.7) \quad {}^E A \xrightarrow{\varphi} A \xrightarrow{\nu} A^E \in \mathbf{Ho}^s$$

given by $\sum^{-1}(B_\gamma/A) \rightarrow A \rightarrow B_\gamma$ above, and observe that A^E is $[E,]_*$ -trivial. Indeed, ν is clearly the $[E,]_*$ -trivialization of A , i.e. ν is the initial example of a map from A to an $[E,]_*$ -trivial spectrum. It is useful to observe:

(1.8) If $V \rightarrow X \rightarrow W$ is a cofibre sequence in \mathbf{Ho}^s with V $[E,]_*$ -colocal and with W $[E,]_*$ -trivial, then $V \rightarrow X \rightarrow W$ is equivalent to the cofibre sequence

$${}^E X \xrightarrow{\varphi} X \xrightarrow{\nu} X^E$$

It is straightforward to check that the $[E,]_*$ -colocalization and $[E,]_*$ -trivialization functors on \mathbf{Ho}^s commute with suspension and preserve cofibre sequences. In [Bousfield 3] we will show that for each $E \in \mathbf{Ho}^s$ there exists a spectrum $aE \in \mathbf{Ho}^s$ such that the E_* -localization and E_* -acyclization functors are respectively equivalent to the $[aE,]_*$ -trivialization and $[aE,]_*$ -colocalization functors on \mathbf{Ho}^s . Thus, many examples of trivialization and colocalization functors will be implicitly studied in [Bousfield 3].

§2. On the structure of $\mathbf{A}(\mathbf{Ho}^s)$

We now examine the structure of the class $\mathbf{A}(\mathbf{Ho}^s)$ of “acyclicity types” of spectra, and we establish the results mentioned in the introduction concerning the distributive lattice $\mathbf{DL}(\mathbf{Ho}^s) \subset \mathbf{A}(\mathbf{Ho}^s)$ and the Boolean algebra $\mathbf{BA}(\mathbf{Ho}^s) \subset \mathbf{DL}(\mathbf{Ho}^s)$.

$\mathbf{A}(\mathbf{Ho}^s)$ has the following relations and operations:

(2.1) For $\langle X \rangle, \langle Y \rangle \in \mathbf{A}(\mathbf{Ho}^s)$, define $\langle X \rangle \leq \langle Y \rangle$ if each Y_* -acyclic spectrum is X_* -acyclic. This is a partial order relation. Clearly $\langle 0 \rangle$ is the smallest element of $\mathbf{A}(\mathbf{Ho}^s)$ and $\langle S \rangle$ is the largest. Note that if X is $[Y,]_*$ -colocal (or equivalently, if $X \in \text{Class-}Y$), then $\langle X \rangle \leq \langle Y \rangle$.

(2.2) For a set $\{\langle X_\alpha \rangle\}$ of elements in $\mathbf{A}(\mathbf{Ho}^s)$, define $\vee_\alpha \langle X_\alpha \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ by $\vee_\alpha \langle X_\alpha \rangle = \langle \vee_\alpha X_\alpha \rangle$. Note that $\vee_\alpha \langle X_\alpha \rangle$ is the least upper bound of $\{\langle X_\alpha \rangle\}$ in $\mathbf{A}(\mathbf{Ho}^s)$, and \vee is associative, commutative, and idempotent. Of course, $\langle 0 \rangle \vee \langle X \rangle$ and $\langle S \rangle \vee \langle X \rangle = \langle S \rangle$.

(2.3) For $\langle X \rangle, \langle Y \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ define $\langle X \rangle \wedge \langle Y \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ by $\langle X \rangle \wedge \langle Y \rangle = \langle X \wedge Y \rangle$. This is well-defined: if $\langle X \rangle = \langle X_1 \rangle$ and $\langle Y \rangle = \langle Y_1 \rangle$, then clearly $\langle X \wedge Y \rangle = \langle X_1 \wedge Y \rangle = \langle X_1 \wedge Y_1 \rangle$. Note that $\langle X \rangle \wedge \langle Y \rangle$ is a lower bound of $\langle X \rangle, \langle Y \rangle \in \mathbf{A}(\mathbf{Ho}^s)$, and that if $\langle X \rangle \leq \langle X_1 \rangle$ and $\langle Y \rangle \leq \langle Y_1 \rangle$ then $\langle X \rangle \wedge \langle Y \rangle \leq \langle X_1 \rangle \wedge \langle Y_1 \rangle$. Clearly \wedge is associative and commutative, with $\langle S \rangle \wedge \langle X \rangle = \langle X \rangle$ and $\langle 0 \rangle \wedge \langle X \rangle = \langle 0 \rangle$. Also the distributive law $\langle X \rangle \wedge (\vee_\alpha \langle Y_\alpha \rangle) = \vee_\alpha (\langle X \rangle \wedge \langle Y_\alpha \rangle)$ and absorption law $\langle X \rangle \vee (\langle X \rangle \wedge \langle Y \rangle) = \langle X \rangle$ hold.

(2.4) For each $\langle X \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ there is an element $a\langle X \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ such that $a\langle X \rangle$ is the greatest member of $\mathbf{A}(\mathbf{Ho}^s)$ with $\langle X \rangle \wedge a\langle X \rangle = \langle 0 \rangle$. Moreover, $aa\langle X \rangle = \langle X \rangle$ for each $\langle X \rangle \in \mathbf{A}(\mathbf{Ho}^s)$, and $\langle X \rangle \leq \langle Y \rangle$ if and only if $a\langle Y \rangle \leq a\langle X \rangle$. This will be shown in [Bousfield 3], and we remark that $a\langle X \rangle = \langle aX \rangle$ where aX is the spectrum mentioned at the end of §1. It turns out that $\mathbf{DL}(\mathbf{Ho}^s)$ is not closed under $a()$, although $a()$ gives the complement in $\mathbf{BA}(\mathbf{Ho}^s)$. We won't use $a()$ in this paper.

So far, $\mathbf{A}(\mathbf{Ho}^s)$ resembles a Boolean algebra with complement $a()$, but the following lemma shows that \wedge is not idempotent in $\mathbf{A}(\mathbf{Ho}^s)$.

LEMMA 2.5. *Let $X \in \mathbf{Ho}^s$ be a finite CW-spectrum with H_*X finite, and let $cX \in \mathbf{Ho}^s$ be the Brown-Comenetz dual of X . If $X \neq 0$, then $\langle cX \rangle \wedge \langle cX \rangle = \langle 0 \rangle \neq \langle cX \rangle$.*

Proof. Using [Brown-Comenetz, 1.14] it is easy to show $H_*(cX; Z) = 0$, and thus $\langle H \rangle \wedge \langle cX \rangle = \langle 0 \rangle$ where H is the spectrum for integral homology. Since $\pi_i cX$ is the Pontrjagin dual of $\pi_{-i} X$, it vanishes for sufficiently large i . Hence

$(cX)(n, \infty) \in \text{Class-}H$ for each n where $(cX)(n, \infty)$ is the $(n - 1)$ -connected section of cX . The cofibre sequence

$$\bigvee_{n \leq 0} (cX)(n, \infty) \rightarrow \bigvee_{n \leq 0} (cX)(n, \infty) \rightarrow cX \in \mathbf{Ho}^s$$

now shows $cX \in \text{Class-}H$, and thus $\langle cX \rangle \leq \langle H \rangle$. The lemma now follows since $\langle cX \rangle \wedge \langle cX \rangle \leq \langle H \rangle \wedge \langle cX \rangle = \langle 0 \rangle$ and since $(cX)_*(S) \neq 0 = 0_*(S)$.

To avoid the pathological spectra revealed by 2.5, we introduce

2.6 The distributive lattice of spectra $\mathbf{DL}(\mathbf{Ho}^s)$

Let $\mathbf{DL}(\mathbf{Ho}^s)$ consist of all $\langle X \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ with $\langle X \rangle \wedge \langle X \rangle = \langle X \rangle$. For instance, if E is a ring spectrum, then $\langle E \rangle \in \mathbf{DL}(\mathbf{Ho}^s)$ because E is a retract of $E \wedge E$ in \mathbf{Ho}^s . Also, if E a Moore spectrum or a finite CW spectrum, then $\langle E \rangle \in \mathbf{DL}(\mathbf{Ho}^s)$ by 2.9 and 2.13 below. Many other examples can be derived from the preceding, since $\mathbf{DL}(\mathbf{Ho}^s)$ is closed under the operation \vee (with any number of summands) and under \wedge ; the proof for \vee uses the equality $\langle X \rangle \vee (\langle X \rangle \wedge \langle Y \rangle) = \langle X \rangle$. With the operations \vee and \wedge , $\mathbf{DL}(\mathbf{Ho}^s)$ is clearly a distributive lattice with $0, 1$ as defined in the next paragraph.

We refer the reader to [Dwinger] or [Grätzer] for an exposition of distributive lattice theory, but for convenience we recall that a class L with binary operations \vee, \wedge is a *distributive lattice with $0, 1$* if:

- (i) $x \wedge x = x$ and $x \vee x = x$ for $x \in L$.
- (ii) $x \wedge y = y \wedge x$ and $x \vee y = y \vee x$ for $x, y \in L$.
- (iii) $x \wedge (y \wedge z) = (x \wedge y) \wedge z$ and $x \vee (y \vee z) = (x \vee y) \vee z$ for $x, y, z \in L$.
- (iv) $x \wedge (x \vee y) = x$ and $x \vee (x \wedge y) = x$ for $x, y \in L$.
- (v) $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ and $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ for $x, y, z \in L$.
- (vi) There exist elements $0, 1 \in L$ such that $0 \vee x = x$ and $1 \wedge x = x$ for all $x \in L$.

(Clearly, 0 and 1 are unique.) Now let L be a distributive lattice with $0, 1$. For $x, y \in L$ one writes $x \leq y$ if the equivalent conditions $x \wedge y = x$ and $x \vee y = y$ are satisfied. Then \leq is a partial order relation on L , and $x \vee y$ (resp. $x \wedge y$) is the l.u.b. (resp. g.l.b.) of $x, y \in L$, cf. [Dwinger, p. 44] or [Grätzer, p. 6]. We also recall that L is called a *Boolean algebra* if for each $x \in L$ there exists $y \in L$ with $x \wedge y = 0$ and $x \vee y = 1$.

For $\langle X \rangle, \langle Y \rangle \in \mathbf{DL}(\mathbf{Ho}^s)$ we conclude that $\langle X \rangle \wedge \langle Y \rangle$ is the g.l.b. of $\langle X \rangle$ and $\langle Y \rangle$, where $\mathbf{DL}(\mathbf{Ho}^s)$ has the partial ordering inherited from $\mathbf{A}(\mathbf{Ho}^s)$. Of course, we previously observed that $\langle X \rangle \vee \langle Y \rangle$ is the l.u.b. of $\langle X \rangle$ and $\langle Y \rangle$. Thus the algebraic structure of $\mathbf{DL}(\mathbf{Ho}^s)$ is contained in its partial ordering.

We call $\langle Y \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ the *complement* of $\langle X \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ if $\langle X \rangle \wedge \langle Y \rangle = \langle 0 \rangle$ and $\langle X \rangle \vee \langle Y \rangle = \langle S \rangle$. Note that if $\langle Y_1 \rangle$ is also the complement of $\langle X \rangle$, then $\langle Y \rangle = \langle Y_1 \rangle$

because

$$\langle Y \rangle = \langle Y \rangle \wedge (\langle X \rangle \vee \langle Y_1 \rangle) = \langle Y \rangle \wedge \langle Y_1 \rangle = (\langle X \rangle \vee \langle Y \rangle) \wedge \langle Y_1 \rangle = \langle Y_1 \rangle.$$

If $\langle X \rangle$ has a complement $\langle Y \rangle$, then $\langle X \rangle \in \mathbf{DL}(\mathbf{Ho}^s)$ because $\langle X \rangle = \langle X \rangle \wedge (\langle X \rangle \vee \langle Y \rangle) = \langle X \rangle \wedge \langle X \rangle$, but the members of $\mathbf{DL}(\mathbf{Ho}^s)$ need not have complements.

LEMMA 2.7. $\langle H \rangle \in \mathbf{DL}(\mathbf{Ho}^s)$, but $\langle H \rangle$ does not have a complement.

Proof. $\langle H \rangle \in \mathbf{DL}(\mathbf{Ho}^s)$ since H is a ring spectrum. Suppose $\langle H \rangle$ has a complement $\langle L \rangle$. Let $\langle cX \rangle$ be as in 2.5, and recall that $\langle cX \rangle \neq \langle 0 \rangle = \langle H \rangle \wedge \langle cX \rangle$ and $\langle cX \rangle \leq \langle H \rangle$. Thus

$$\langle cX \rangle = (\langle H \rangle \vee \langle L \rangle) \wedge \langle cX \rangle = \langle L \rangle \wedge \langle cX \rangle \leq \langle L \rangle \wedge \langle H \rangle = \langle 0 \rangle$$

and this contradicts $\langle cX \rangle \neq \langle 0 \rangle$. Therefore $\langle H \rangle$ cannot have a complement.

We now introduce

2.8 The Boolean algebra of spectra $\mathbf{BA}(\mathbf{Ho}^s)$

Let $\mathbf{BA}(\mathbf{Ho}^s)$ consist of all $\langle X \rangle \in \mathbf{A}(\mathbf{Ho}^s)$ such that $\langle X \rangle$ has a complement (written $\langle X \rangle^c$), and note that $\mathbf{BA}(\mathbf{Ho}^s) \subset \mathbf{DL}(\mathbf{Ho}^s)$. If E is a Moore spectrum or a (possibly infinite) wedge of finite CW spectra, then $\langle E \rangle \in \mathbf{BA}(\mathbf{Ho}^s)$ by 2.9 and 2.13 below. Many other members of $\mathbf{BA}(\mathbf{Ho}^s)$ can be derived from the preceding, since $\mathbf{BA}(\mathbf{Ho}^s)$ is clearly closed under $(\)^c$ and the binary operations \vee, \wedge ; indeed, for $\langle X \rangle, \langle Y \rangle \in \mathbf{BA}(\mathbf{Ho}^s)$

$$\langle X \rangle^{cc} = \langle X \rangle$$

$$(\langle X \rangle \vee \langle Y \rangle)^c = \langle X \rangle^c \wedge \langle Y \rangle^c$$

$$(\langle X \rangle \wedge \langle Y \rangle)^c = \langle X \rangle^c \vee \langle Y \rangle^c.$$

With these operations, $\mathbf{BA}(\mathbf{Ho}^s)$ is clearly a Boolean algebra.

As promised, we now prove

PROPOSITION 2.9. *If $E \in \mathbf{Ho}^s$ is a (possibly infinite) wedge of finite CW-spectra, then $\langle E \rangle \in \mathbf{BA}(\mathbf{Ho}^s)$. Moreover, $\langle E \rangle = \langle {}^E S \rangle$ and $\langle E \rangle^c = \langle S^E \rangle$.*

Proof. Assume $E = \vee_{\alpha} B_{\alpha}$ where each B_{α} is a finite CW spectrum. A spectrum $Y \in \mathbf{Ho}^s$ is $[E,]$ -trivial iff $(DB_{\alpha}) \wedge Y \simeq 0 \in \mathbf{Ho}^s$ for all α , where DB_{α} is the

Spanier-Whitehead dual of B_α . Thus if Y is $[E,]_*$ -trivial, then so is $X \wedge Y$ for all $X \in \mathbf{Ho}^s$. In particular, ${}^E S \wedge S^E$ is $[E,]_*$ -trivial as well as $[E,]_*$ -colocal (by 1.4), and thus ${}^E S \wedge S^E \simeq 0 \in \mathbf{Ho}^s$. Using the cofibering ${}^E S \rightarrow S \rightarrow S^E$ of (1.7), we conclude that $\langle {}^E S \rangle \vee \langle S^E \rangle = \langle S \rangle$, so $\langle {}^E S \rangle \in \mathbf{BA}(\mathbf{Ho}^s)$ with $\langle {}^E S \rangle^c = \langle S^E \rangle$. It remains to show $\langle E \rangle = \langle {}^E S \rangle$. Applying (1.8) to the cofibre sequence

$$X \wedge {}^E S \rightarrow X \wedge S \rightarrow X \wedge S^E,$$

we find that $X \wedge {}^E S \simeq {}^E X$ and $X \wedge S^E \simeq X^E$ for all $X \in \mathbf{Ho}^s$. Since E is $[E,]_*$ -colocal, this implies $E \wedge {}^E S \simeq E$, and thus $\langle E \rangle \leq \langle {}^E S \rangle$. Since ${}^E S$ is $[E,]_*$ -colocal, we know $\langle {}^E S \rangle \leq \langle E \rangle$, and therefore $\langle E \rangle = \langle {}^E S \rangle$.

We remark that the above spectra ${}^E S$ and S^E satisfy the strong idempotency conditions ${}^E S \wedge {}^E S \simeq {}^E S$ and $S^E \wedge S^E \simeq S^E$. Indeed S^E is a commutative ring spectrum whose multiplication map $S^E \wedge S^E \rightarrow S^E$ is an equivalence. We next observe

PROPOSITION 2.10. *If $E \in \mathbf{Ho}^s$ is a finite CW spectrum, then $\langle E \rangle = \langle DE \rangle$. Consequently, for any $G \in \mathbf{Ho}^s$, $G_*(E) = 0 \Leftrightarrow G^*(E) = 0$.*

Proof. Since $[E, S^E]_* = 0$, we have $(DE) \wedge S^E \simeq 0$ and thus $(DE) \wedge {}^E S \simeq DE$. Since $\langle {}^E S \rangle = \langle E \rangle$, this implies $\langle DE \rangle \wedge \langle E \rangle = \langle DE \rangle$. Dually one shows $\langle E \rangle \wedge \langle DE \rangle = \langle E \rangle$, and therefore $\langle DE \rangle = \langle E \rangle$. The last statement is deduced using $G^*(E) = G_*(DE)$.

We next prove “triangle (in)equalities” for cofibre sequences. Call a map $f: A \rightarrow X \in \mathbf{Ho}^s$ *smash nilpotent* if the m -fold smash product

$$f \wedge \cdots \wedge f: A \wedge \cdots \wedge A \rightarrow X \wedge \cdots \wedge X \in \mathbf{Ho}^s$$

is the 0 map for some $m \geq 1$. Note that the smash nilpotent maps form a subgroup of $[A, X]$, and a composite fg is smash nilpotent if either f or g is. Moreover, if $i \neq 0$ then each map $S^i \rightarrow S^0 \in \mathbf{Ho}^s$ is a smash nilpotent by [Nishida].

PROPOSITION 2.11. *If $A \xrightarrow{f} X \rightarrow B$ is a cofibre sequence in \mathbf{Ho}^s , then:*

- (i) $\langle A \rangle \leq \langle X \rangle \vee \langle B \rangle$, $\langle X \rangle \leq \langle A \rangle \vee \langle B \rangle$, and $\langle B \rangle \leq \langle A \rangle \vee \langle X \rangle$.
- (ii) If $A, X \in \mathbf{DL}(\mathbf{Ho}^s)$ and f is smash nilpotent, then $\langle B \rangle = \langle A \rangle \vee \langle X \rangle$.

Proof. Part (i) is obvious. For (ii) we assume $f \wedge \cdots \wedge f = 0$ in \mathbf{Ho}^s and form a cofibre sequence

$$A \wedge \cdots \wedge A \xrightarrow{f \wedge \cdots \wedge f} X \wedge \cdots \wedge X \rightarrow C \in \mathbf{Ho}^s$$

Then

$$\langle C \rangle = \langle A \wedge \cdots \wedge A \rangle \vee \langle X \wedge \cdots \wedge X \rangle = \langle A \rangle \vee \langle X \rangle$$

and also $\langle C \rangle \leq \langle B \rangle$ because C is $[B,]_*$ -colocal (i.e. $C \in \text{Class-}B$). Thus $\langle A \rangle \vee \langle X \rangle \leq \langle B \rangle$ and the opposite inequality is given by (i).

COROLLARY 2.12. *If $n \neq 1$ and $\alpha : S^{n-1} \rightarrow S^0$ in \mathbf{Ho}^s , then $\langle S^0 \cup_\alpha e^n \rangle = \langle S \rangle$.*

Because of Nishida's theorem, this follows from 2.11; or instead of using 2.11, we could have used the easy result that if $\sum^m A \xrightarrow{f} A \rightarrow B$ is a cofibre sequence in \mathbf{Ho}^s with f nilpotent in $[A, A]_*$, then $\langle B \rangle = \langle A \rangle$. By combining 2.12 with R. Wood's result $K \simeq (S^0 \cup_n e^2) \wedge KO$, we recover the result $\langle K \rangle = \langle KO \rangle$, cf. [Meier], [Ravenel].

Of course, 2.12 fails when $n = 1$, and we now consider $\langle SG \rangle$ where G is an abelian group and $SG \in \mathbf{Ho}$ is a Moore spectrum of type $(G, 0)$ (i.e. $\pi_i SG = 0$ for $i < 0$, $H_0 SG = G$, and $H_i SG = 0$ for $i > 0$). There is a short exact sequence

$$0 \rightarrow G \otimes \pi_n X \rightarrow \pi_n (SG \wedge X) \rightarrow \text{Tor}(G, \pi_{n-1} X) \rightarrow 0.$$

Thus

$$\langle S \rangle = \langle SQ \rangle \vee \bigvee_{p \text{ prime}} \langle SZ/p \rangle$$

$$\langle SQ \rangle \wedge \langle SZ/p \rangle = \langle 0 \rangle = \langle SZ/p \rangle \wedge \langle SZ/q \rangle \text{ for primes } p \neq q.$$

It follows that $\mathbf{BA}(\mathbf{Ho}^s)$ has a sub-Boolean algebra $\mathbf{MBA}(\mathbf{Ho}^s)$ whose members are the wedges of subsets of

$$I = \{ \langle SQ \rangle, \langle SZ/2 \rangle, \langle SZ/3 \rangle, \langle SZ/5 \rangle, \dots \}.$$

Moreover, there is an obvious Boolean algebra isomorphism between $\mathbf{MBA}(\mathbf{Ho}^s)$ and the power set $P(I)$. Note that for any set J of primes

$$\langle SZ_{(J)} \rangle = \langle SQ \rangle \vee \bigvee_{p \in J} \langle SZ/p \rangle$$

where $Z_{(J)}$ is the localization of Z at J . More generally,

PROPOSITION 2.13. *For each abelian group G , $\langle SG \rangle \in \mathbf{MBA}(\mathbf{Ho}^s)$.*

Proof. Let C be the class of all abelian groups A such that $G \otimes A = 0 = \text{Tor}(G, A)$, and note that $(SG)_* X = 0$ if $\pi_i X \in C$ for all i . The result now follows easily from [Bousfield 1,2.3] since C is a "special" class.

We conclude by noting that $\mathbf{BA}(\mathbf{Ho}^s)$ contains many elements outside $\mathbf{MBA}(\mathbf{Ho}^s)$. For p prime let

$$A(p): \Sigma^m SZ/p \rightarrow SZ/p \in \mathbf{Ho}^s$$

be the K_* -equivalence of [Adams 1, §12] where $m = 2p - 2$ for p odd and $m = 8$ for $p = 2$. It is easy to check that the cofibre of $A(p)$ represents an element of $\mathbf{BA}(\mathbf{Ho}^s)$ outside $\mathbf{MBA}(\mathbf{Ho}^s)$. In [Bousfield 3] we will show that $\langle K \rangle = \langle E \rangle^c$ where E is the wedge of the cofibres of all the $A(p)$.

REFERENCES

- ADAMS, J. F. 1: *On the groups $J(X)$ -IV*. *Topology* 5 (1966), 21–71.
 ADAMS, J. F. 2: *Stable homotopy and generalized homology*. The University of Chicago Press, 1974.
 BALBES, R. and DWINGER, P.: *Distributive Lattices*. The University of Missouri Press, 1974.
 BOUSFIELD, A. K., 1: *Types of acyclicity*. *J. Pure Appl. Algebra* 4 (1974), 293–298.
 BOUSFIELD, A. K., 2: *Constructions of factorization systems in categories*. *J. Pure Appl. Algebra* 9 (1977), 207–220.
 BOUSFIELD, A. K., 3: *The localization of spectra with respect to homology*. (To appear).
 BROWN E. H. and COMENETZ, M. *Pontrjain duality for generalized homology and cohomology theories*. *Amer. J. of Math.* 98 (1976), 1–27.
 GRÄTZER, G., *Lattice Theory*. W. H. Freeman and Company, San Francisco, 1971.
 MEIER, W., *Localization with respect to K -theory*. To appear.
 NISHIDA, G., *The nilpotency of elements of the stable homotopy groups of spheres*, *J. Math. Soc. Japan* 25 (1973), 707–732.
 RAVENEL, D., *Localization with respect to certain periodic homology theories*. Lecture given at Northwestern University, March 1977.

*Dept. of Mathematics,
 University of Illinois at Chicago Circle,
 Box 4348, Chicago, Ill. 60680 U.S.A.*

Received May 17, 1978