

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 55 (1980)  
  
**Artikel:** On cohomological periodicity for infinite groups.  
**Autor:** Talelli, Olympia  
**DOI:** <https://doi.org/10.5169/seals-42369>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 20.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## On cohomological periodicity for infinite groups

OLYMPIA TALELLI

### Introduction

The phenomenon of cohomological periodicity for finite groups has long been understood. Here we introduce the notion of “a group  $G$  having period  $q$  after  $k$ -steps” so as to allow infinite groups to have periodic cohomology. The definition is given in terms of a projective resolution of  $G$  and, as expected, it is equivalent to having the functors  $H^n(G, -)$  and  $H^{n+q}(G, -)$  naturally isomorphic for all  $n \geq k + 1$ . We then show that this definition coincides with the classical one for finite groups and moreover, we obtain that if an infinite group  $G$  has period  $q$  after  $k$ -steps then  $k \geq 1$ .

In §2 we investigate what it means for a countable locally finite group to have period  $q$  after  $k$ -steps. We obtain what one would expect, i.e. that a countable locally finite group  $G$  has period  $q$  after  $k$ -steps iff every finite subgroup of  $G$  has period  $q$ . Moreover, we have here that  $k = 1$ .

Then we show that there is an element  $g \in H^q(G, \mathbb{Z})$  such that cup product with  $g$

$$\cup g : H^i(G, -) \rightarrow H^{i+q}(G, -)$$

induces the natural isomorphism for all  $i \geq 2$ .

Finally, in §3 we characterize the infinite locally finite  $p$ -groups which have period  $q$  after  $k$ -steps. First, we point out two obvious candidates, i.e. the infinite locally cyclic  $p$ -group and the infinite locally quaternion group, and then we show that these are the only ones. This result depends heavily on the well known similar statement for periodic finite  $p$ -groups [Cartan + Eilenberg].

I wish to thank K. W. Gruenberg for his help during the preparation of this paper.

### §1. Periodicity after some “steps”

Let  $G$  be a group and  $ZG$  its integral group ring. We work in the category of left  $ZG$ -modules. If  $A$  is a  $ZG$ -module, by a resolution of  $A$  we shall always mean a projective resolution of  $A$ .

DEFINITION. A group  $G$  is said to have period  $q$  after  $k$ -steps if there is an exact sequence

$$0 \longrightarrow R_{k+q} \xrightarrow{\beta} P_{k+q-1} \longrightarrow \cdots \longrightarrow P_k \xrightarrow{\partial_k} P_{k-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow Z \longrightarrow 0$$

$\begin{array}{c} \nearrow \alpha \\ R_k \end{array}$

(1)

where  $Z$  is regarded as a trivial  $ZG$ -module,  $R_{k+q} = R_k$  and  $P_i$   $0 \leq i \leq k+q-1$  are projective  $ZG$ -modules. We take  $R_0 = Z$ . If  $k=0$  then  $G$  is said to have period  $q$ . Having (1) we can form a resolution of  $G$

$$P'_* : \cdots \longrightarrow P'_i \xrightarrow{\partial'_i} P'_{i-1} \longrightarrow \cdots \longrightarrow P'_0 \longrightarrow Z \longrightarrow 0$$

$\begin{array}{c} \searrow \\ R'_i \end{array}$

by defining

$$P'_i = P_i \quad 0 \leq i \leq k-1 \quad \partial'_i = \partial_i \quad 1 \leq i \leq k$$

and

$$\begin{aligned} P'_i &= P_{k+\lambda_i} & i &= k+nq+\lambda_i & \partial'_i &= \partial_{k+\lambda_i} & i &= k+nq+\lambda_i \\ & & n &\geq 0, 0 \leq \lambda_i < q & & & n &\geq 0, 0 < \lambda_i < q \\ & & & & \partial'_{k+\mu q} &= \beta\alpha & \mu &\geq 1 \end{aligned}$$

Such a resolution is naturally considered periodic after  $k$ -steps; we may refer to (1) itself as a resolution of  $G$  which is periodic after  $k$ -steps. If  $k=0$  we call such a resolution a periodic resolution of  $G$ .

Evidently the functors  $H^n(G, -)$  and  $H^{n+q}(G, -)$  are naturally isomorphic for all  $n \geq k+1$ .

Now let  $M, N$  be  $ZG$ -modules,  $g \in \text{Ext}_{ZG}^q(M, N)$  and consider the following diagram

$$\begin{array}{ccccccccccc} P_* : & \cdots & \longrightarrow & P_{i+q} & \longrightarrow & P_{i+q-1} & \longrightarrow & \cdots & \longrightarrow & P_{q+1} & \longrightarrow & P_q & \longrightarrow & P_{q-1} & \longrightarrow & \cdots & \longrightarrow & P_0 & \longrightarrow & M & \longrightarrow & 0 \\ & & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \searrow & & \nearrow & & & & & & \\ E_* : & \cdots & \longrightarrow & E_i & \longrightarrow & E_{i-1} & \longrightarrow & \cdots & \longrightarrow & E_1 & \longrightarrow & E_0 & \longrightarrow & N & \longrightarrow & 0 \end{array}$$

$\begin{array}{c} \nearrow \\ R_q \\ \downarrow \theta \end{array}$

where  $P_* \rightarrow M$ ,  $E_* \rightarrow N$  are resolutions of  $M, N$  respectively and  $\theta : R_q \rightarrow M$  a

$q$ -cocycle representing  $g$ . Then  $\theta$  lifts to a chain map  $P_* \rightarrow E_*$  of degree  $-q$  and this chain map induces cup product with  $g$

$$\bigcup g : \text{Ext}_{ZG}^i(N, -) \rightarrow \text{Ext}_{ZG}^{i+q}(M, -) \quad i \geq 0.$$

**THEOREM 1.1.** *Let  $G$  be a group. Then the following statements are equivalent:*

- (i)  $G$  has period  $q$  after  $k$ -steps
- (ii) There is a resolution of  $G$

$$\begin{array}{ccccccc} \cdots & \longrightarrow & P_i & \longrightarrow & P_{i-1} & \longrightarrow & \cdots \longrightarrow P_0 \longrightarrow Z \longrightarrow 0 \\ & & \searrow & & \nearrow & & \\ & & R_i & & & & \end{array}$$

and an element  $h \in \text{Ext}_{ZG}^q(R_k, R_k)$  such that for every  $ZG$ -module  $A$  cup product with  $h$

$$\bigcup h : \text{Ext}_{ZG}^i(R_k, A) \longrightarrow \text{Ext}_{ZG}^{i+q}(R_k, A)$$

is an isomorphism for  $i \geq 1$  and an epimorphism for  $i = 0$ . Note that  $\text{Ext}_{ZG}^i(R_k, A) = H^{i+k}(G, A)$  for  $i \geq 1$ ,  $k \geq 0$ .

*Proof.* (i)  $\Rightarrow$  (ii). Consider a resolution  $P'_*$  of  $G$  which is periodic, or period  $q$ , after  $k$ -steps

$$\begin{array}{ccccccc} P'_* : \cdots & \longrightarrow & P'_i & \longrightarrow & P'_{i-1} & \longrightarrow & \cdots \longrightarrow P'_0 \longrightarrow Z \longrightarrow 0 \\ & & \searrow & & \nearrow & & \\ & & R'_i & & & & \end{array}$$

and the resolution  $P'_* | R'_k$  of  $R'_k$

$$P'_* |_{R_k} : \cdots \longrightarrow P'_i \longrightarrow P'_{i-1} \longrightarrow \cdots \longrightarrow P'_k \xrightarrow{\alpha} R'_k \longrightarrow 0.$$

Now there is an element  $h \in \text{Ext}_{ZG}^q(R'_k, R'_k)$  defined by  $P'_{k+q} \xrightarrow{\alpha} R'_k$  or in Yoneda's interpretation of  $\text{Ext}_{ZG}^i(-, -)$  by the multiple extension

$$0 \longrightarrow R'_k \longrightarrow P'_{k+q-1} \longrightarrow \cdots \longrightarrow P'_k \xrightarrow{\alpha} R'_k \longrightarrow 0.$$

Clearly, for any  $ZG$ -module  $A$  cup product with  $h$

$$\bigcup h : \text{Ext}_{ZG}^i(R_k, A) \longrightarrow \text{Ext}_{ZG}^{i+q}(R_k, A)$$



is the identity for  $i \geq 1$  and an epimorphism for  $i = 0$ . (ii)  $\Rightarrow$  (i) can be easily deduced from the following Theorem. It is due to C. T. C. Wall [[7], Theorem. 1.2]; however, we give here a different proof.

**THEOREM 1.2.** *Let  $A, B$  be  $ZG$ -modules. If  $g \in \text{Ext}_{ZG}^n(A, B)$  is such that for any  $ZG$ -module  $E$  cup product with  $g$*

$$\cup g : \text{Ext}_{ZG}^i(B, E) \longrightarrow \text{Ext}_{ZG}^{i+n}(A, E)$$

*is injective for  $i = 1$  and surjective for  $i = 0$ , then  $g$  is represented by a multiple extension*

$$0 \longrightarrow B \longrightarrow K \longrightarrow P_{n-2} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow A \longrightarrow 0$$

*with  $K, P_i$   $0 \leq i \leq n-2$  projective  $ZG$ -modules.*

*Proof.* Let  $P_* \rightarrow A, E_* \rightarrow B$  be resolutions of  $A$  and  $B$  respectively, and let  $\theta : R_n \rightarrow B$  be a cocycle representing  $g$

$$\begin{array}{ccccccccccc} P_* : & \cdots & \longrightarrow & P_{n+1} & \longrightarrow & P_n & \longrightarrow & P_{n-1} & \longrightarrow & P_{n-2} & \longrightarrow & \cdots & \longrightarrow & P_0 & \longrightarrow & A & \longrightarrow & 0 \\ & & & \downarrow & & \downarrow & \nearrow & \downarrow & & & & & & & & & & & \\ & & & E_1 & \longrightarrow & E_0 & \longrightarrow & B & \longrightarrow & 0. & & & & & & & & \\ & & & & & \downarrow \theta & & & & & & & & & & & & & \end{array}$$

Now  $\theta$  lifts to a chain map  $P_* \rightarrow E_*$  of degree  $-n$  which induces cup product with  $g$ .

Let

$$\begin{array}{ccc} R_n & \xrightarrow{\gamma} & P_{n-1} \\ \theta \downarrow & & \downarrow \alpha \\ B & \xrightarrow{\beta} & K \end{array}$$

be a push-out diagram. Then  $\beta$  is injective and  $\text{coker } \beta \cong \text{coker } \gamma \cong R_{n-1}$ .

Now  $g$  is represented by  $0 \rightarrow B \rightarrow K \rightarrow P_{n-2} \rightarrow \cdots \rightarrow P_0 \rightarrow A \rightarrow 0$ . We shall show that  $K$  is a projective  $ZG$ -module.

Consider the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & R_n & \longrightarrow & P_{n-1} & \longrightarrow & R_{n-1} \longrightarrow 0 \\ & & \downarrow \theta & & \downarrow \alpha & & \downarrow id \\ 0 & \longrightarrow & B & \longrightarrow & K & \longrightarrow & R_{n-1} \longrightarrow 0. \end{array}$$

This gives rise to the following commutative diagram

$$\begin{array}{ccccccc}
 \cdots \rightarrow \operatorname{Hom}_{ZG}(B, E) & \rightarrow & \operatorname{Ext}_{ZG}(R_{n-1}, E) & \rightarrow & \operatorname{Ext}_{ZG}(K, E) & \rightarrow & \operatorname{Ext}_{ZG}(B, E) \rightarrow \cdots \\
 \theta^* \downarrow & \searrow & \downarrow \operatorname{id} & & \downarrow \alpha^* & & \downarrow \theta^* \\
 \cdots \rightarrow \operatorname{Hom}_{ZG}(R_n, E) & \xrightarrow{\delta} & \operatorname{Ext}_{ZG}(R_{n-1}, E) & \rightarrow & \operatorname{Ext}_{ZG}(P_{n-1}, E) & \rightarrow & \operatorname{Ext}_{ZG}(R_n, E) \rightarrow \cdots
 \end{array}$$

By hypotheses  $\theta^*: \operatorname{Ext}_{ZG}(B, E) \rightarrow \operatorname{Ext}_{ZG}(R_n, E) \simeq \operatorname{Ext}_{ZG}^{1+n}(A, E)$  is injective and

$$\delta\theta^*: \operatorname{Hom}_{ZG}(B, E) \rightarrow \operatorname{Ext}_{ZG}(R_{n-1}, E) \simeq \operatorname{Ext}_{ZG}^n(A, E)$$

is surjective. Moreover,  $\operatorname{Ext}_{ZG}(P_{n-1}, E) = 0$  since  $P_{n-1}$  is projective. Thus by the commutativity of the diagram we obtain that  $\operatorname{Ext}_{ZG}(K, E) = 0$ , and this holds for any  $ZG$ -module  $E$ . Hence  $K$  is a projective  $ZG$ -module.

*Remark.* If  $P_{n-1}, B$  are finitely generated  $ZG$ -modules then  $K$  is a finitely generated module since from the push-out diagram we have an epimorphism  $P_{n-1} \oplus B \xrightarrow{\alpha+\beta} K \longrightarrow 0$ .

**PROPOSITION 1.3.** *If  $G$  has period  $q$  after  $k$ -steps then so does every subgroup  $H$  of  $G$ .*

*Proof.* This follows from the fact that a projective  $ZG$ -module is a projective  $ZH$ -module.

**PROPOSITION 1.4.** *If  $G$  is an infinite group and has period  $q$  after  $k$ -steps then  $k \geq 1$ .*

*Proof.* Let  $H$  be a group which has period  $q$  and consider a periodic resolution of  $H$

$$0 \longrightarrow Z \xrightarrow{\beta} P_{q-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow Z \longrightarrow 0.$$

Now  $\operatorname{im} \beta \hookrightarrow H^0(H, P_{q-1})$ . But  $P_{q-1}$  is a direct summand of an induced  $ZH$ -module, and it is well known that if  $A$  is a non-trivial induced  $ZH$ -module then  $H^0(H, A) \neq 0$  iff  $H$  is finite. The result follows.

**LEMMA 1.5.** *If  $G$  is a finite group and has period  $q$  after  $k$ -steps, then  $G$  has period  $q$ .*

*Proof.* By Theorem 1.1 (i)  $\Rightarrow$  (ii) there is a resolution of  $G$

$$\begin{array}{ccccccc}
 \cdots \longrightarrow & P_i & \longrightarrow & P_{i-1} & \longrightarrow & \cdots & \longrightarrow P_0 \longrightarrow Z \longrightarrow 0 \\
 & \searrow & & \nearrow & & & \\
 & & & R_i & & & 
 \end{array} \tag{2}$$

and an element  $h \in \text{Ext}_{ZG}^q(R_k, R_k)$  such that for every  $ZG$ -module  $A$  cup product with  $h$

$$\bigcup h : \text{Ext}_{ZG}^i(R_k, A) \rightarrow \text{Ext}_{ZG}^{i+q}(R_k, A)$$

is an isomorphism for  $i \geq 1$ .

Now since  $G$  is a finite group, a projective  $ZG$ -module  $P$  is a direct summand of a coinduced  $ZG$ -module; hence  $H^i(G, P) = 0$  for all  $i \geq 1$ . Thus from (2) we obtain an isomorphism

$$\delta : \text{Ext}_{ZG}^q(R_k, R_k) \xrightarrow{\cong} H^q(G, Z).$$

Let  $h' \in H^q(G, Z)$  be the image of  $h \in \text{Ext}_{ZG}^q(R_k, R_k)$  under  $\delta$ . Then clearly for any  $ZG$ -module  $A$  cup product with  $h'$

$$\bigcup h' : H^i(G, A) \rightarrow H^{i+q}(G, A)$$

is an isomorphism for  $i \geq k + 1$ .

Let  $0 \rightarrow C \rightarrow P \rightarrow A \rightarrow 0$  be a short exact sequence with  $P$  a projective  $ZG$ -module. Then we obtain the following commutative diagram

$$\begin{array}{ccc} H^{n-1}(G, A) & \xrightarrow{\bigcup h'} & H^{n+q-1}(G, A) \\ \delta_n \downarrow & & \downarrow \delta_{n+q} \\ H^n(G, C) & \xrightarrow{\bigcup h'} & H^{n+q}(G, C) \end{array}$$

where  $\delta_n, \delta_{n+q}$  are connecting homomorphisms. But since  $H^i(G, P) = 0$  for all  $i \geq 1$ ,  $\delta_i$  is an isomorphism for  $i > 1$  and an epimorphism for  $i = 1$ . Thus it follows that cup product with  $h'$

$$\bigcup h' : H^i(G, A) \longrightarrow H^{i+q}(G, A)$$

is an isomorphism for  $i \geq 1$  and an epimorphism for  $i = 0$ . The result now follows from Theorem 1.1 (ii)  $\Rightarrow$  (i).

**PROPOSITION 1.6.** *If  $G$  has period  $q$  after  $k$ -steps then every finite subgroup  $H$  of  $G$  has period  $q$ .*

*Proof.* This follows from Proposition 1.3 and Lemma 1.5.

**COROLLARY 1.7.** *If  $G$  has period 2 after  $k$ -steps then every finite subgroup  $H$  of  $G$  is cyclic.*

*Proof.* It is known [[6], Lemma 5.2] that a finite group  $H$  has period 2 iff it is cyclic; hence the result follows from Proposition 1.6.

Let  $G$  be a group such that the functors  $H^i(G, -)$  and  $H^{i+q}(G, -)$  are naturally isomorphic for some  $j \geq 1$ . Consider a resolution of  $G$

$$\begin{array}{ccccccc} \cdots & \longrightarrow & P_i & \longrightarrow & P_{i-1} & \longrightarrow & \cdots \longrightarrow P_0 \longrightarrow Z \longrightarrow 0. \\ & & \searrow & & \nearrow & & \\ & & R_i & & & & \end{array} \quad (3)$$

Then clearly  $\text{Ext}_{ZG}(R_{j-1}, -) \cong^{\text{nat.}} \text{Ext}_{ZG}(R_{j+q-1}, -)$ . Thus by [[4], Thm. 2.6] there exist projective  $ZG$ -modules  $Q_1, Q_2$  such that

$$R_{j-1} \oplus Q_1 \xrightarrow{\alpha} R_{j+q-1} \oplus Q_2. \quad (*)$$

Now (3) gives rise to the following exact sequence

$$\begin{array}{ccccccccccc} 0 & \longrightarrow & R_{j+q-1} \oplus Q_2 & \xrightarrow{\alpha} & P_{j+q-2} \oplus Q_2 & \longrightarrow & P_{j+q-3} & \longrightarrow & \cdots & \longrightarrow & P_{j-2} & \longrightarrow \\ & & & & & & & & & & & \\ & & & & & & & & & & P_{j-1} \oplus Q_1 & \longrightarrow P_{j-2} \oplus Q_1 \longrightarrow P_{j-3} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow Z \longrightarrow 0 \end{array} \quad (4)$$

$$\begin{array}{ccc} & \searrow & \nearrow \\ & R_{j-1} \oplus Q_1 & \\ & \searrow & \nearrow \\ & R_{j-2} & \end{array}$$

Hence it follows from (4) and (\*) that:

**PROPOSITION 1.8.** *If the functors  $H^\lambda(G, -)$  and  $H^{\lambda+q}(G, -)$  are naturally isomorphic for some  $\lambda \geq 2$ , then  $G$  has period  $q$  after  $\lambda-1$ -steps.*

Now let  $j = 1$  and  $q > 0$ . Clearly it follows that  $H^n(G, -) \cong^{\text{nat.}} H^{n+q}(G, -)$  for all  $n \geq 1$ . Thus by Proposition 1.8  $G$  has period  $q$  after 1-step. But  $G$  is finite since there is a monomorphism  $\varrho\alpha : Z \oplus Q_1 \rightarrow P_{q-1} \oplus Q_2$  (same argument as for the proof of Proposition 1.4). Thus by Lemma 1.5  $G$  has period  $q$ . Hence we have proved:

**PROPOSITION 1.9.** *If the functors  $H^1(G, -)$  and  $H^{1+q}(G, -)$  are naturally isomorphic for some  $q > 0$ , then  $G$  is finite and has period  $q$ .*

## 2. Periodic countable locally finite groups

In this section we state and prove our main Theorem. The proof is based on direct limit arguments.

**PROPOSITION 2.1** [[1]]. *Let  $(P_i, \lambda_{ij})_I$  be a countable direct system of projective  $ZG$ -modules. Then there exists an exact sequence  $0 \rightarrow Q \rightarrow Q \rightarrow P \rightarrow 0$  where  $P = \varinjlim (P_i, \lambda_{ij})_I$  and  $Q$  is a projective  $ZG$ -module.*

**LEMMA 2.2.** *Let  $\cdots \rightarrow A_n \xrightarrow{\alpha_n} A_{n-1} \rightarrow \cdots \rightarrow A_1 \xrightarrow{\alpha_1} A_0 \xrightarrow{\pi} \text{coker } \alpha_1 \rightarrow 0$  be an exact sequence of  $ZG$ -modules such that for every  $n \geq 0$  there is a resolution  $P_{n*} \rightarrow A_n$  of the form  $0 \rightarrow P_{n,1} \xrightarrow{\partial_1^n} P_{n,0} \xrightarrow{\partial_0^n} A_n \rightarrow 0$ . If  $\{\alpha_*^j\}: P_{j*} \rightarrow P_{j-1*}$  are chain maps lifting  $\alpha_j: A_j \rightarrow A_{j-1}$   $j \geq 1$  then there is a resolution  $P_* \rightarrow \text{coker } \alpha_1$*

$$P_*: \cdots \longrightarrow P_k \xrightarrow{\partial_k} P_{k-1} \longrightarrow \cdots \longrightarrow P_1 \xrightarrow{\partial_1} P_0 \xrightarrow{\varepsilon} \text{coker } \alpha_1 \longrightarrow 0$$

where

$$P_k = P_{k-1,1} \oplus P_{k,0} \quad k \geq 0$$

$$\partial_k = \begin{cases} P_{k,0} & \xrightarrow{\alpha_0^k - (\partial_1^{k-2})^{-1} \alpha_0^{k-1} \alpha_0^k} P_{k-1,0} \oplus P_{k-2,1} \\ P_{k-1,1} & \xrightarrow{\partial_1^{k-1} - \alpha_1^{k-1}} P_{k-1,0} \oplus P_{k-2,1} \end{cases} \quad k \geq 1$$

and  $\varepsilon = \pi \partial_0^0$ .

Note that  $P_{ij}$  is understood to be zero if  $i < 0$  or  $j < 0$ .

*Proof.* We have the following commutative diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & A_n & \xrightarrow{\alpha_n} & A_{n-1} & \longrightarrow & \cdots \longrightarrow A_1 \xrightarrow{\alpha_1} A_0 \xrightarrow{\pi} \text{coker } \alpha_1 \longrightarrow 0 \\ & & \uparrow \partial_0^n & & \uparrow \partial_0^{n-1} & & \uparrow \partial_0^1 & & \uparrow \partial_0^0 \\ \cdots & \longrightarrow & P_{n,0} & \xrightarrow{\alpha_0^n} & P_{n-1,0} & \longrightarrow & \cdots \longrightarrow P_{1,0} \xrightarrow{\alpha_0^1} P_{0,0} \\ & & \uparrow \partial_1^n & & \uparrow \partial_1^{n-1} & & \uparrow \partial_1^1 & & \uparrow \partial_1^0 \\ \cdots & \longrightarrow & P_{n,1} & \xrightarrow{\alpha_1^n} & P_{n-1,1} & \longrightarrow & \cdots \longrightarrow P_{1,1} \xrightarrow{\alpha_1^1} P_{0,1} \end{array}$$

Now it is easily seen that this gives rise to the resolution  $P_* \rightarrow \text{coker } \alpha_1$ .

We shall need the following notion.

Let  $(G_i, \gamma_{ij})_I$  be a direct system of groups. For  $i \in I$  let  $(C_*^i, \partial_*^i)$  be a  $ZG_i$ -chain complex. If  $i \leq j$  then let  $c_{ij}: (C_*^i, \partial_*^i) \rightarrow (C_*^j, \partial_*^j)$  be a  $ZG_i$ -chain map, where  $(C_*^j, \partial_*^j)$  is considered as a  $ZG_i$ -chain complex via  $\gamma_{ij}: G_i \rightarrow G_j$ , such that

- (1)  $c_{ii}$  is the identity chain map of  $(C_*^i, \partial_*^i)$ , for all  $i \in I$  and
- (2) if  $i \leq j \leq k$  then  $c_{jk}c_{ij} = c_{ik}$  as  $ZG_i$ -chain maps.

We call  $((C_*^i, \partial_*^i), c_{ij})_I$  a direct system of chain complexes over the direct system of groups  $(G_i, \gamma_{ij})_I$ . In particular let  $(A_i, \alpha_{ij})_I$  be a direct system of modules over the direct system of groups  $(G_i, \gamma_{ij})_I$ . Clearly  $(A_i, \alpha_{ij})_I$  is a direct system of abelian groups and if  $A = \varinjlim (A_i, \alpha_{ij})_I$  then  $G = \varinjlim (G_i, \gamma_{ij})_I$  acts on  $A$  in a natural way [[2], ch. 2, §6, nos 6, 7].

Moreover, if  $B_i = ZG \otimes_{ZG_i} A_i$  and  $\beta_{ij} : B_i \rightarrow B_j$

$$x \otimes_{ZG_i} a_i \rightarrow x \otimes_{ZG_j} \alpha_{ij} a_i$$

then it follows that  $(B_i, \beta_{ij})_I$  is a direct system of  $ZG$ -modules and  $\varinjlim (B_i, \beta_{ij})_I \simeq A$  as  $ZG$ -modules.

Recall that if  $X$  is a class of groups, then a group  $G$  is said to be locally an  $X$ -group if every finite set of elements of  $G$  is contained in some  $X$ -subgroup of  $G$ .

**THEOREM 2.3.** *Let  $G$  be a countable locally finite group all of whose finite subgroups have period  $q$ . Then  $G$  has period  $q$  after 1-step.*

*Proof.* It is easily seen that there is a direct system of finite subgroups of  $G$  over  $I = \{1, 2, 3, \dots\}$ ,  $(G_i, e_{ij})_I$ , with  $e_{ij} : G_i \rightarrow G_j$  inclusions and  $G = \varinjlim (G_i, e_{ij})_I$ . By hypothesis for each  $i \in I$  we have a periodic resolution  $P_*^{i'}$  of  $G_i$

$$P_*^{i'} : 0 \longrightarrow Z \longrightarrow P_{q-1}^{i'} \longrightarrow \dots \longrightarrow P_1^{i'} \longrightarrow ZG_i \xrightarrow{\varepsilon_i} Z \longrightarrow 0.$$

Now consider the following diagram

$$\begin{array}{ccccccccccc} P_*^{i'} & 0 & \longrightarrow & Z & \longrightarrow & P_{q-1}^{i'} & \longrightarrow & \dots & \longrightarrow & P_1^{i'} & \longrightarrow & ZG_i & \xrightarrow{\varepsilon_i} & Z & \longrightarrow & 0 \\ \varrho'_{i+1} \downarrow & & & \downarrow h & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow id_Z & & \\ P_*^{i+1'} & 0 & \longrightarrow & Z & \longrightarrow & P_{q-1}^{i+1'} & \longrightarrow & \dots & \longrightarrow & P_1^{i+1'} & \longrightarrow & ZG_{i+1} & \xrightarrow{\varepsilon_{i+1}} & Z & \longrightarrow & 0. \end{array}$$

Since  $P_*^{i+1'}$  is evidently a resolution of  $G_{i+1}$  via  $e_{i+1} : G_i \rightarrow G_{i+1}$  we can lift  $id_Z$  to a  $ZG_i$ -chain map  $\varrho'_{i+1} : P_*^{i'} \rightarrow P_*^{i+1'}$ .

Then  $\varrho'_{i+1}$  induces a map  $h : Z \rightarrow Z$  (multiplication by  $h$ ) which need not be the identity on  $Z$ .

Our aim is to find a periodic resolution  $P_*^{i+1'}$  of  $G_{i+1}$  and a  $ZG_i$ -chain map  $\varrho_{i+1} : P_*^{i'} \rightarrow P_*^{i+1'}$  which lifts  $id_Z$  on the right and induces  $id_Z$  on the left. Now the map  $\varrho'_{i+1}$  induces an isomorphism between the cohomology groups of  $G_i$  defined using  $P_*^{i'}$  and those defined using  $P_*^{i+1'}$ . On  $H^q(G_i, Z)$  this map is obviously

multiplication by  $h$ . Since  $H^q(G_i, Z) \simeq Z/|G_i|Z$  by the periodicity, it follows that  $(h, |G_i|) = 1$ ; hence there are integers  $\lambda_1, \lambda_2$  such that  $1 = \lambda_1 h + \lambda_2 |G_i|$ .

Let  $\lambda = \lambda_1 + x |G_i|$  where  $x = p_1 \cdots p_m$  with  $p_j$   $1 \leq j \leq m$  primes such that  $p_j \nmid |G_{i+1}|$  and  $p_j \nmid |G_i|$ ,  $p_j \nmid \lambda_1$  for all  $1 \leq j \leq m$ . If no such primes exist, take  $x = 1$ . Then

$$(\lambda, |G_{i+1}|) = 1 \quad \text{and} \quad 1 = \lambda h + (\lambda_2 - xh) |G_i| (*).$$

Consider  $P_*^{i+1'}$  as a projective resolution of  $G_{i+1}$ . Then  $id_Z$  is a  $q$ -cocycle which defines an element  $g_{i+1} \in H^q(G_{i+1}, Z)$ . Clearly  $\lambda g_{i+1} \in H^q(G_{i+1}, Z)$  is represented by the  $q$ -cocycle  $\lambda : Z \rightarrow Z$  (multiplication by  $\lambda$ ). We shall show that  $\lambda g_{i+1} \in H^q(G_{i+1}, Z) = \text{Ext}_{ZG_{i+1}}^q(Z, Z)$  is represented by a multiple extension

$$P_*^{i+1} : 0 \longrightarrow Z \longrightarrow P_{q-1}^{i+1} \longrightarrow P_{q-2}^{i+1} \longrightarrow \cdots \longrightarrow P_1^{i+1} \longrightarrow ZG_{i+1} \longrightarrow Z \longrightarrow 0$$

with  $P_k^{i+1}$   $1 \leq k \leq q-1$  projective  $ZG_{i+1}$ -modules. By Theorem 1.2 it is enough to show that for any  $ZG_{i+1}$ -module  $A$  cup product with  $\lambda g_{i+1}$

$$\bigcup \lambda g_{i+1} : H^k(G_{i+1}, A) \longrightarrow H^{k+q}(G_{i+1}, A)$$

is injective for  $k=1$  and surjective for  $k=0$ . This follows easily since cup product with  $\lambda g_{i+1}$  is multiplication by  $\lambda$  for  $k=0, 1$  and we have that  $(\lambda, |G_{i+1}|) = 1$ . Note that  $|G_{i+1}| H^k(G_{i+1}, A) = 0$  for all  $k \geq 1$  and by the periodicity  $H^q(G_{i+1}, A) \simeq A^{G_{i+1}} / (\sum_{g \in G_{i+1}} g)A$  where  $A^{G_{i+1}} \simeq \text{Hom}_{ZG_{i+1}}(Z, A)$ .

Moreover, from the proof of Theorem 1.2, we obtain the following commutative diagram

$$\begin{array}{ccccccccccc} P_*^{i'} & 0 & \longrightarrow & Z & \xrightarrow{\mu_i} & P_{q-1}^{i'} & \longrightarrow & \cdots & \longrightarrow & P_1^{i'} & \longrightarrow & ZG_i & \longrightarrow & Z & \longrightarrow & 0 \\ e'_{i+1} \downarrow & & & \downarrow h & & \downarrow \alpha & & & & \downarrow & & \downarrow & & \downarrow id_Z & & \\ P_*^{i+1'} & 0 & \longrightarrow & Z & \xrightarrow{\gamma} & P_{q-1}^{i+1'} & \longrightarrow & P_{q-2}^{i+1'} & \longrightarrow & \cdots & \longrightarrow & P_1^{i+1'} & \longrightarrow & ZG_{i+1} & \longrightarrow & Z & \longrightarrow & 0 \\ L' \downarrow & & & \downarrow \lambda & & \downarrow \beta & & \parallel & & \parallel & & \parallel & & \parallel & & \downarrow id_Z & & \\ P_*^{i+1} & 0 & \longrightarrow & Z & \xrightarrow{\mu_{i+1}} & P_{q-1}^{i+1} & \longrightarrow & P_{q-2}^{i+1} & \longrightarrow & \cdots & \longrightarrow & P_1^{i+1} & \longrightarrow & ZG_{i+1} & \longrightarrow & Z & \longrightarrow & 0 \end{array}$$

Clearly  $L' e'_{i+1} : P_*^{i'} \rightarrow P_*^{i+1}$  is a  $ZG_i$ -chain map.

Now consider  $P_*^{i'}$  as a resolution of  $G_i$ . Then  $id_Z$  is a  $q$ -cocycle which defines an element  $g_i \in H^q(G_i, Z)$ .

Clearly the  $q$ -cocycle  $\lambda h : Z \rightarrow Z$  represents  $\lambda h g_i$ . But from (\*)  $\lambda h g_i = g_i$  since  $|G_i| g_i = 0$ . Hence the cocycles  $h$  and  $id_Z$  represent the same element. So there

exists  $\gamma: P'_{q-1} \rightarrow Z$  such that  $id_Z - \lambda h = \gamma \mu_i$ . If we now take  $\beta\alpha + \mu_{i+1}\gamma$  instead of  $\beta\alpha$  we still have a chain map  $P_*^{i'} \rightarrow P_*^{i+1}$  which now induces  $id_Z$  on the left. We call this chain map  $\varrho_{i+1}$ , i.e. we have the following commutative diagram

$$\begin{array}{ccccccccccc} P_*^{i'} & 0 \longrightarrow & Z & \longrightarrow & P_{q-1}^{i'} & \longrightarrow & \cdots & \longrightarrow & P_1^{i'} & \longrightarrow & ZG_i & \xrightarrow{\varepsilon_i} & Z & \longrightarrow & 0 \\ \varrho_{i+1} \downarrow & & \downarrow id_Z & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow id_Z & & \\ P_*^{i+1} & 0 \longrightarrow & Z & \longrightarrow & P_{q-1}^{i+1} & \longrightarrow & \cdots & \longrightarrow & P_1^{i+1} & \longrightarrow & ZG_{i+1} & \xrightarrow{\varepsilon_{i+1}} & Z & \longrightarrow & 0. \end{array}$$

Take  $P_*^{1'} = P_*^1$ . Then given  $P_*^1$  and  $P_*^{2'}$  we construct as above a periodic resolution  $P_*^2$  of  $G_2$  and a  $ZG_1$ -chain map  $\varrho_{12}: P_*^1 \rightarrow P_*^2$  which lifts  $id_Z$  on the right and induces  $id_Z$  on the left. In this way we construct inductively a direct system  $(P_*^i, \varrho_{ij})_I$  of periodic resolutions over the direct system of groups  $(G_i, e_{ij})_I$ . Clearly  $\varrho_{ij(i \leq j)}: P_*^i \rightarrow P_*^j$  are given as  $\varrho_{ij} = \varrho_{j-1j} \cdots \varrho_{i+1j}$  and they lift  $id_Z$  on the right and induce  $id_Z$  on the left.

Now since direct limit preserves exactness, taking the direct limit of  $(P_*^i, \varrho_{ij})_I$  we obtain the following exact sequence of  $ZG$ -modules

$$0 \longrightarrow Z \xrightarrow{\theta} \varinjlim P_{q-1}^i \xrightarrow{\alpha_{q-1}} \varinjlim P_{q-2}^i \longrightarrow \cdots \longrightarrow \varinjlim P_1^i \xrightarrow{\alpha_1} ZG \xrightarrow{\varepsilon} Z \longrightarrow 0 \quad (*)'$$

By splicing together copies of  $(*)'$  we obtain an exact sequence of  $ZG$ -modules

$$\begin{aligned} \cdots \xrightarrow{\alpha_1} ZG \xrightarrow{\alpha_q} \varinjlim P_{q-1}^i \xrightarrow{\alpha_{q-1}} \cdots \longrightarrow \varinjlim P_1^i \xrightarrow{\alpha_1} ZG \longrightarrow \\ \xrightarrow{\alpha_q} \varinjlim P_{q-1}^i \xrightarrow{\alpha_{q-1}} \cdots \longrightarrow \varinjlim P_1^i \xrightarrow{\alpha_1} ZG \xrightarrow{\varepsilon} Z \longrightarrow 0 \quad (1) \end{aligned}$$

where  $\alpha_q = \theta\varepsilon$ .

By Proposition 2.1 we have resolutions  $Q_{j*} \rightarrow \varinjlim P_j^i$  of the form

$$0 \longrightarrow Q_{j1} \longrightarrow Q_{j0} \longrightarrow \varinjlim P_j^i \longrightarrow 0 \quad \text{for all } 1 \leq j \leq q-1.$$

Now the hypotheses of Lemma 2.2 hold for (1). Moreover, it is clear that we can choose here the chain maps  $\{a_*^n\}$  of Lemma 2.2 so as

$$\{\alpha_*^{j+kq}\} = \{\alpha_*^j\}, \quad \{\alpha_*^{\lambda q}\} = \{\alpha_*^q\} \quad 1 \leq j \leq q-1, \quad k \geq 0, \quad \lambda \geq 1.$$



$$\begin{array}{c} \cdots \longrightarrow Q_{2,0} \oplus Q_{1,1} \xrightarrow{\partial_{q+2}} Q_{1,0} \xrightarrow{\partial_{q+1}} ZG \oplus Q_{q-1,1} \longrightarrow \cdots \\ \qquad \qquad \qquad \searrow \qquad \nearrow \\ \qquad \qquad \qquad R_{q+1} \end{array}$$

$$\begin{array}{c} \longrightarrow Q_{2,0} \oplus Q_{1,1} \xrightarrow{\partial_2} Q_{1,0} \xrightarrow{\partial_1} ZG \xrightarrow{\epsilon} Z \longrightarrow 0 \\ \qquad \qquad \qquad \searrow \qquad \nearrow \\ \qquad \qquad \qquad R_1 \end{array}$$

*Remark.* Note that in view of Proposition 1.4, Theorem 2.3 is the best possible we can obtain if  $G$  is an infinite countable locally finite group.

(i)  $G$  has period  $q$  after  $k$ -steps.  
(ii) Every finite subgroup of  $G$  has period  $q$ .  
Moreover,  $k = 0$  if  $G$  is finite and  $k = 1$  if  $G$  is infinite.

Now if  $G$  is a finite group and  $P$  a projective  $ZG$ -module then  $H^i(G, P) = 0$  for all  $i \geq 1$ . We have an analogous result for a countable locally finite group:

If  $G$  is a countable locally finite group and  $A$  a  $ZG$ -module such that for some  $n \geq 2$

Now let  $G$  be a countable locally finite group which has period  $q$  after  $k$ -steps. Then by Corollary 2.4  $G$  has period  $q$  after 1-step. Thus by Theorem 1.1 (i)  $\Rightarrow$  (ii) there is a resolution of  $G$

$$\cdots \longrightarrow P_i \longrightarrow P_{i-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow Z \longrightarrow 0$$

$\searrow \quad \nearrow$   
 $R_i$

and an element  $g' \in \text{Ext}_{ZG}^q(R_1, R_1)$  such that for every  $ZG$ -module  $A$  cup product with  $g'$

$$\bigcup g' : \text{Ext}_{ZG}^i(R_1, A) \longrightarrow \text{Ext}_{ZG}^{i+q}(R_1, A)$$

is an isomorphism for all  $i \geq 1$ .

By Proposition 2.5 if  $P$  is a projective  $ZG$ -module then  $H^j(G, P) = 0$  for all  $j \geq 2$ . Having this result one obtains from (2) that  $\text{Ext}_{ZG}^q(R_1, R_1) \xrightarrow{\delta} H^q(G, Z)$ ; note that  $q \geq 2$  since by Proposition 1.6  $q$  is a period of every finite subgroup of  $G$ , and that is known to be even [[3], ch. XII, p. 261].

Now if  $g \in H^q(G, Z)$  corresponds to  $g' \in \text{Ext}_{ZG}^q(R_1, R_1)$  under  $\delta$  then one clearly has that for any  $ZG$ -module  $A$  cup product with  $g$

$$\bigcup g : H^i(G, A) \longrightarrow H^{i+q}(G, A)$$

is an isomorphism for all  $i \geq 2$ . Thus we have proved that:

**PROPOSITION 2.6.** *Let  $G$  be a countable locally finite group which has period  $q$  after  $k$ -steps. Then there is an element  $g \in H^q(G, Z)$  such that for any  $ZG$ -module  $A$  cup product with  $g$*

$$\bigcup g : H^i(G, A) \longrightarrow H^{i+q}(G, A)$$

*is an isomorphism for all  $i \geq 2$ .*

### §3. Periodic locally finite $p$ -groups

The following well known theorem characterizes the finite  $p$ -groups which have period  $q > 0$ .

**THEOREM 3.1** [[3], ch. XII, Thm. 11.6]. *For a finite  $p$ -group  $G$  the following statements are equivalent:*

- (i)  $G$  has period  $q > 0$
  - (ii)  $G$  is either cyclic or is a (generalized) quaternion group.
- Moreover, a cyclic group has period 2 and a (generalized) quaternion group has period 4.

We shall characterize the infinite locally finite  $p$ -groups which have period  $q > 0$  after  $k$ -steps.

I. Consider an infinite locally cyclic  $p$ -group. It is easily seen that such a group is uniquely determined up to isomorphism by the prime  $p$  and is given by

$$C_p^\infty = \langle c_1, c_2, \dots, c_i, \dots; c_1^p = 1, c_2^p = c_1, \dots, c_{i+1}^p = c_i, \dots \rangle.$$

II. Recall that a (generalized) quaternion group  $Q_{2^i}$  is given by

$$Q_{2^i} = \langle x, y; x^{2^{i-2}} = y^2, xyx = y \rangle \quad i \geq 3.$$

One easily shows that a (generalized) quaternion group  $Q_{2^i}$  contains a normal cyclic subgroup  $C$  of index 2, which is characteristic if  $i > 3$ , and every element of  $Q_{2^i} \setminus C$  is of order four and inverts every element of  $C$ . Having this result and following the proof of Proposition 1.I.2 [8], one shows:

**PROPOSITION 3.2.** *Let  $G$  be a locally quaternion group. Then  $G$  has a locally cyclic normal subgroup  $C$  of index 2, and every element of  $G \setminus C$  is of order four and inverts every element of  $C$ .*

**COROLLARY 3.3.** *Up to isomorphism there exists only one infinite locally quaternion group, namely*

$$Q_2^\infty = \langle c_1, \dots, c_n, \dots, y; c_1^2 = 1, \dots, c_{n+1}^2 = c_n, \dots, y^2 = c_1, \\ c_i^y c_i = 1 \quad \text{all } i \geq 1 \rangle.$$

*Proof.* This follows easily from Proposition 3.2 and the fact that up to isomorphism there exists only one infinite locally cyclic 2-group, namely,  $C_2^\infty$ .

**PROPOSITION 3.4.** (i) *An infinite locally cyclic  $p$ -group has period 2 after 1-step.*  
(ii) *An infinite locally quaternion group has period 4 after 1-step.*

*Proof.* (i) Clearly it is enough to consider  $C_{p^\infty}$ . Now  $C_{p^\infty}$  is a countable group. Moreover, every finite subgroup of  $C_{p^\infty}$  is cyclic, hence by Theorem 3.1 it has period 2. The result now follows from Theorem 2.3.

(ii) By Corollary 3.3 it is enough to consider  $Q_{2^\infty}$ . Clearly  $Q_{2^\infty}$  is countable. Moreover, if  $K$  is a finite subgroup of  $Q_{2^\infty}$  then  $K$  is contained in some (generalized) quaternion group; hence by Theorem 3.1 and Proposition 1.3  $K$  has period 4. Now the result follows from Theorem 2.3.

**THEOREM 3.5.** *For an infinite locally finite  $p$ -group  $G$  the following statements are equivalent:*

(i)  $G$  has period  $q > 0$  after  $k$ -steps.

(ii)  $G$  is either  $C_{p^\infty}$  or  $Q_{2^\infty}$ .

Moreover,  $k = 1$ . If  $G = C_{p^\infty}$  then  $q = 2$  and if  $G = Q_{2^\infty}$  then  $q = 4$ .

*Proof.* (i)  $\Rightarrow$  (ii). Let  $G$  have period  $q$  after  $k$ -steps. By Proposition 1.6 every finite subgroup of  $G$  has period  $q$  and therefore by Theorem 3.1 every finite subgroup of  $G$  is either cyclic or is a (generalized) quaternion group. Thus  $G$  is either an infinite locally cyclic  $p$ -group or an infinite locally quaternion group, i.e.  $G$  is either  $C_{p^\infty}$  or by Corollary 3.3,  $G$  is  $Q_{2^\infty}$ . (ii)  $\Rightarrow$  (i) follows from Proposition 3.4.

## REFERENCES

- [1] BERSTEIN, I. *On the dimension of modules and algebras IX. Direct limits*. Nagoya Math. J. 13 (1958), 83–84.
- [2] BOURBAKI, N. *Algebra*, Chapter 2; Hermann (ASI 1236) (1962).
- [3] CARTAN H. and EILENBERG, S. *Homological Algebra*. Princeton University Press.
- [4] HILTON P. J. and REES, D. *Natural maps of extension functors and a theorem of R. G. Swan*. Proc. Camb. Philos. Soc. 57 (1961) 489–502.
- [5] ROBINSON, D. J. S. *On the cohomology of soluble groups of finite rank*. J. Pure Applied Algebra 6 (1975), 155–164.
- [6] SWAN, R. G. *Minimal resolutions for finite groups*, Topology 4, (1965), 193–208.
- [7] WALL, C. T. C. *Periodic projective resolutions*, to appear.
- [8] WEHRFRITZ B. and KEGEL, O. *Locally finite groups*, North-Holland 1973.

*Forschungsinstitut für Mathematik  
ETH Zentrum  
CH-8092 Zürich*

Received March 10, 1979.