

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 55 (1980)  
  
**Artikel:** On the Euler class of representations of finite groups over real fields.  
**Autor:** Eckmann, Beno / Mislin, Guido  
**DOI:** <https://doi.org/10.5169/seals-42379>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 04.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# On the Euler class of representations of finite groups over real fields

BENO ECKMANN and GUIDO MISLIN

## Introduction

For representations of finite groups over the rationals  $\mathbf{Q}$  there is a uniform bound, depending on the degree  $m$  of the representation only, for the order of the Euler class. This has been proved in [E–M], and the best possible such bound was shown there to be  $E_m = \text{denominator of } B_m/m$  if  $m$  is even, where  $B_m$  is the  $m$ -th Bernoulli number (and, of course,  $E_m = 2$  if  $m$  is odd). The Euler class of a representation  $\rho: G \rightarrow \text{GL}_m(\mathbf{R})$  is an element of  $H^m(G; \mathbf{Z}(\rho))$ ,  $\mathbf{Z}(\rho)$  being the group of integers turned into a  $G$ -module by multiplication with  $\text{sgn det } \rho$  and hence a trivial  $G$ -module if and only if  $\rho$  is “orientable.”

In the present paper we discuss analogous bounds for representations realizable over an arbitrary real field  $K \subset \mathbf{R}$  instead of the rationals  $\mathbf{Q}$ . The universal bound is expressed in terms of a certain operator  $\mathcal{E}_K(m)$  on finite Abelian groups, depending on  $K$  and  $m$  only.  $\mathcal{E}_K(m)$  is defined (cf. Section 3.1), for each prime  $p$ , by its action on  $p$ -torsion. This action depends on the degree  $\varphi_K(p)$  of the  $p$ -th cyclotomic extension of  $K$ , and on a further invariant  $\gamma_K(p) \in \mathbf{N} \cup \infty$  attached to  $K$  and  $p$ , cf. Section 2.2. The main theorem states that if the representation  $\rho$  of a finite group  $G$ , of degree  $m$ , is realizable over  $K$  then

$$\mathcal{E}_K(m)e(\rho) = 0. \quad (*)$$

Moreover  $\mathcal{E}_K(m)$  is best possible in that sense.

We mention here some properties of the operator  $\mathcal{E}_K(m)$ . If  $m$  is not divisible by  $\varphi_K(p)$ , then  $\mathcal{E}_K(m)$  is the identity operator on  $p$ -torsion; thus (\*) just expresses the fact (Proposition 2.1) that in that case the  $p$ -component of  $e(\rho)$  is 0. If  $m$  is divisible by  $\varphi_K(p)$ , one has two different possibilities. Either  $\gamma_K(p) = \infty$ ; then  $\mathcal{E}_K(m)$  annihilates  $p$ -torsion, and (\*) tells nothing about the  $p$ -component of  $e(\rho)$ : in fact, there is, in that case, *no* universal bound for the order of the  $p$ -component of  $e(\rho)$  (Corollary 2.4). Or  $\gamma_K(p) < \infty$ ; then  $\mathcal{E}_K(m)$  is, on  $p$ -torsion, multiplication by  $p^{\gamma_K(p) + \nu_p}$ , where  $\nu_p$  is the exponent of  $p$  in the prime decomposition of  $m$ .

If we assume  $\gamma_K(p) < \infty$  for all primes  $p$ , and if  $\varphi_K(p)$  divides  $m$  for a finite number of primes  $p$  only, then  $\mathcal{E}_K(m)$  can be replaced by multiplication with the integer  $E_K(m) = \text{lcm}\{n \mid m \equiv 0 \pmod{\varphi_K(n)}\}$ . For  $K = \mathbf{Q}$ ,  $E_{\mathbf{Q}}(m) = E_m$  is the integer mentioned above. The assumption is fulfilled for all real number fields  $K$ . Statement (\*) then tells that the order of  $e(\rho)$  divides  $E_K(m)$ , for all finite groups and all  $K$ -representations of degree  $m$ ; and this bound is best possible.

If a representation  $\rho: G \rightarrow GL_m(\mathbf{R})$  is not known to be realizable over a subfield of  $\mathbf{R}$  fixed in advance, we show that (\*) still holds if one takes for  $K$  a field containing the values of the character of  $\rho$  (without assuming  $\rho$  to be defined over  $K \subset \mathbf{R}$ ). In particular we show (Theorem 3.8) that

$$E_{\mathbf{Q}(\chi)}(m)e(\rho) = 0$$

where  $\mathbf{Q}(\chi)$  denotes the field obtained from  $\mathbf{Q}$  by adjoining the values of the character  $\chi$  of  $\rho$ .

We also obtain a bound for the order of  $e(\rho)$  of an arbitrary real representation  $\rho$  in terms of the exponent  $\exp(G)$  of  $G$  (Theorem 3.9):

$$\frac{m}{2} \exp(G) e(\rho) = 0$$

for  $\rho: G \rightarrow GL_m(\mathbf{R})$ ,  $m$  even.

## 1. $K$ -representations of finite $p$ -groups

1.1. Let  $G$  be a finite group, and  $K$  a subfield of the field  $\mathbf{C}$  of complex numbers. For a complex character  $\chi$  of  $G$  we denote by  $K(\chi)$  the Galois field extension obtained by adjoining to  $K$  all values of  $\chi$ . In case  $\chi$  is  $\mathbf{C}$ -irreducible,  $K(\chi)$  is isomorphic to the center of  $A_K(\chi)$ , the unique simple component of the group algebra  $K[G]$  on which  $\chi$  is non-zero. If  $\chi_1$  and  $\chi_2$  are two  $\mathbf{C}$ -irreducible characters of  $G$ , then  $A_K(\chi_1) = A_K(\chi_2)$  if and only if  $\chi_1$  and  $\chi_2$  are *Galois-conjugate over  $K$* , which means that there is a  $\sigma \in \text{Gal}(K(\chi_1)/K)$  such that  $\chi_2(g) = \sigma\chi_1(g)$  for all  $g \in G$ . The  $K$ -irreducible characters of  $K$ -representations of  $G$  are the characters of the form

$$\psi = s_K(\chi) \sum_{\sigma} \sigma\chi$$

where  $\chi$  is  $\mathbf{C}$ -irreducible and the sum is extended over all  $\sigma \in \text{Gal}(K(\chi)/K)$ , and

where  $s_K(\chi)$  denotes the Schur index of  $\chi$  over  $K$  (we recall that  $A_K(\chi)$  is a matrix algebra over a division algebra  $D$ , and that  $s_K(\chi)^2$  is the dimension of  $D$  over its center  $K(\chi)$ ).

1.2. The following result (cf. [E – M], Theorem 1.3) reduces the discussion of  $K$ -representations of finite  $p$ -groups to  $p$ -groups of very special types.

**THEOREM 1.1.** *Let  $G$  be a finite  $p$ -group, and  $\rho: G \rightarrow \text{GL}_m(K)$  an irreducible representation over  $K \subset \mathbb{C}$ . Then either  $\rho$  is induced, or  $\rho$  factors through a faithful representation  $\bar{\rho}: \bar{G} \rightarrow \text{GL}_m(K)$  of a factor group  $\bar{G}$  of  $G$  which is of one of the following types:*

- $C_{p^\alpha}$ ,  $\alpha \geq 0$  (cyclic of order  $p^\alpha$ );
- $Q_{2^\alpha}$ ,  $\alpha \geq 3$  (generalized quaternion group of order  $2^\alpha$ );
- $D_{2^\alpha}$ ,  $\alpha \geq 4$  (dihedral group of order  $2^\alpha$ ); or
- $SD_{2^\alpha}$ ,  $\alpha \geq 4$  (semidihedral group of order  $2^\alpha$ ).

In order to determine the degrees of the faithful irreducible  $K$ -representations of these groups of special type, we use two invariants of  $K$ :

**DEFINITION 1.2.** Let  $K(n)$  denote the “ $n$ -th cyclotomic extension of  $K$ ”; i.e., the field obtained by adjoining to  $K$  the  $n$ -th roots of unity. Then we write  $\varphi_K(n)$  for the dimension of  $K(n)$  over  $K$  and we put

$$\gamma_K(p) = \sup \{ \alpha \mid K(p) = K(p^\alpha) \} \text{ for an odd prime } p,$$

and

$$\gamma_K(2) = \sup \{ \alpha \mid K(4) = K(2^{\alpha+1}) \}.$$

We write sometimes  $\gamma$  for  $\gamma_K(p)$ , if no confusion can arise; there are, of course, cases with  $\gamma = \infty$ .

If  $p$  is an odd prime and  $\alpha \geq 1$  is such that  $K(p^\alpha) \neq K(p^{\alpha+1})$  (i.e.,  $(K(p^{\alpha+1}):K(p^\alpha)) = p$ ) then  $K(p^{\alpha+1}) \neq K(p^{\alpha+2})$ . This follows from the commutative diagram of Galois groups (the maps being induced by restriction)

$$\begin{array}{ccc} \text{Gal}(K(p^{\alpha+2})/K(p^\alpha)) & \rightarrow & \text{Gal}(\mathbf{Q}(p^{\alpha+2})/\mathbf{Q}(p^\alpha)) \cong \mathbf{Z}/p^2\mathbf{Z} \\ \downarrow & & \downarrow \\ \text{Gal}(K(p^{\alpha+1})/K(p^\alpha)) & \rightarrow & \text{Gal}(\mathbf{Q}(p^{\alpha+1})/\mathbf{Q}(p^\alpha)) \cong \mathbf{Z}/p\mathbf{Z} \end{array}$$

Similarly, if  $\alpha \geq 2$ , then  $K(2^\alpha) \neq K(2^{\alpha+1})$  implies  $K(2^{\alpha+1}) \neq K(2^{\alpha+2})$ . Note also that for  $K \subset \mathbf{R}$ ,  $\varphi_K(p)$  is even for  $p$  odd, and  $(K(4):K) = 2$ . The following lemma is now immediate.



LEMMA 1.3. (a) For an odd prime  $p$  one has, for any  $K \subset \mathbf{C}$ ,

$$\varphi_K(p^\alpha) = \begin{cases} \varphi_K(p) & \text{if } 1 \leq \alpha \leq \gamma = \gamma_K(p), \\ \varphi_K(p) \cdot p^{\alpha-\gamma} & \text{if } \alpha \geq \gamma. \end{cases}$$

(b) If  $K \subset \mathbf{R}$  and  $p = 2$ , then

$$\varphi_K(2^\alpha) = \begin{cases} 1 & \text{if } \alpha = 1 \\ 2 & \text{if } 1 < \alpha \leq \gamma + 1 (\gamma = \gamma_K(2)), \\ 2^{\alpha-\gamma} & \text{if } \alpha \geq \gamma + 1. \end{cases}$$

1.3. We now describe the degrees of the faithful irreducible representations of the  $p$ -groups listed in Theorem 1.1, and their orientability.

PROPOSITION 1.4. Let  $K$  be a subfield of  $\mathbf{R}$ , and let  $\rho$  be a faithful irreducible  $K$ -representation of one of the  $p$ -groups  $G$  of special type. Then the degree  $m$  of  $\rho$  is:

$$\begin{aligned} m &= \varphi_K(p^\alpha) && \text{in case } G = C_{p^\alpha} (\alpha \geq 0); \\ m &= 2\varphi_K(2^{\alpha-1}) && \text{in case } G = Q_{2^\alpha} (\alpha \geq 3); \\ m &= \varphi_K(2^{\alpha-1}) && \text{in case } G = D_{2^\alpha} (\alpha \geq 4); \\ m &= \varphi_K(2^{\alpha-1}) \text{ or } 2\varphi_K(2^{\alpha-1}) && \text{in case } G = SD_{2^\alpha} (\alpha \geq 4). \end{aligned}$$

Moreover,  $\rho$  is orientable (i.e., lies in  $SL_m(K)$ ) except for  $G = C_2$ .

*Proof.* The character  $\psi$  of  $\rho$  is of the form  $\psi = s_K(\chi)\Sigma\sigma\chi$ ,  $\sigma \in \text{Gal}(K(\chi)/K)$ , where  $\chi$  is faithful and  $\mathbf{C}$ -irreducible. The faithful and  $\mathbf{C}$ -irreducible representations of the groups of special types were discussed in [E-M]; we will make use of their properties without further reference. The following four cases have to be considered.

$C_{p^\alpha}$ :  $s_K(\chi) = 1$ ,  $\chi$  is of degree one and  $K(\chi) = K(p^\alpha)$ . The degree of  $\psi$  is therefore  $m = |\text{Gal}(K(p^\alpha)/K)| = \varphi_K(p^\alpha)$ .

$Q_{2^\alpha}$ : for any  $K \subset \mathbf{R}$ , one has  $s_K(\chi) = 2$ , and  $\chi$  has degree 2. Since  $K(\chi) = K(2^{\alpha-1}) \cap \mathbf{R}$  and  $\alpha \geq 3$ , we have  $(K(2^{\alpha-1}):K(\chi)) = 2$ . The degree of  $\psi$  is thus given by  $m = 2 \cdot 2 \cdot |\text{Gal}(K(\chi)/K)| = 2|\text{Gal}(K(2^{\alpha-1})/K)| = 2\varphi_K(2^{\alpha-1})$ .

$D_{2^\alpha}$  (or  $SD_{2^\alpha}$  respectively):  $s_K(\chi) = 1$  and  $\chi$  has degree 2. Again we have  $(K(2^{\alpha-1}):K(\chi)) = 2$  (or possibly  $K(2^{\alpha-1}) = K(\chi)$  in the case  $SD_{2^\alpha}$ ) and thus  $m = 2|\text{Gal}(K(\chi)/K)| = \varphi_K(2^{\alpha-1})$  (or possibly  $2\varphi_K(2^{\alpha-1})$  in the case  $SD_{2^\alpha}$ ).

If  $p$  is odd,  $\rho$  is certainly orientable. For  $p=2$  we note that, except for the faithful representation of  $C_2$  of degree 1,  $\psi$  is a sum of an even number of Galois conjugate representations  $\sigma\chi$  which are all orientable in cases  $C_{2^\alpha}$ ,  $\alpha \geq 2$  and  $Q_{2^\alpha}$ ,  $\alpha \geq 3$ ; and which are all non-orientable in the other cases (cf. [E-M]). Hence  $\psi$  is orientable except for  $G = C_2$ .

**COROLLARY 1.5.** *Let  $K$  be a subfield of  $\mathbf{R}$ . The degree of a  $K$ -irreducible representation  $\rho$  of a finite  $p$ -group  $G$  is either 1 or of the form  $\varphi_K(p)p^\beta$ ,  $\beta \geq 0$ .*

*Proof.* We consider the alternative in Theorem 1.1.

If  $\rho$  is induced from a representation  $\tau$  of degree 1, then  $p=2$  and therefore the degree of  $\rho$  is of the form  $2^\beta = \varphi_K(2)2^\beta$  ( $p$  odd would imply that  $\tau$  is a permutation representation, thus reducible). If  $\rho$  is induced from a representation  $\tau$  of degree  $> 1$ , the degree of  $\tau$  is of the form  $\varphi_K(p)p^\beta$ , by induction, and thus the degree of  $\rho$  has the desired form.

If  $\rho$  factors through a faithful representation  $\bar{\rho}$  of  $C_{p^\alpha}$ ,  $Q_{2^\alpha}$ ,  $D_{2^\alpha}$  or  $SD_{2^\alpha}$ , the degree of  $\bar{\rho}$  is  $\varphi_K(p^\alpha)$ ,  $2\varphi_K(2^{\alpha-1})$  or  $\varphi_K(2^{\alpha-1})$ , which is 1 or of the form  $\varphi_K(p)p^\beta$ ,  $\beta \geq 0$ . The assertion of the Corollary thus follows.

## 2. The Euler class of $K$ -representations of $p$ -groups

2.1. For a  $K$ -representation  $\rho: G \rightarrow \mathrm{GL}_m(K)$ , where  $K$  is a subfield of  $\mathbf{R}$ , the Euler class  $e(\rho) \in H^m(G; \mathbf{Z}(\rho))$  is defined as the Euler class of the flat real vector bundle over  $K(G, 1)$ , associated with  $\rho \otimes \mathbf{R}$ ;  $\mathbf{Z}(\rho)$  stands for the  $G$ -module  $\mathbf{Z}$  with  $G$ -action defined by  $g \cdot 1 = \mathrm{sgn} \det \rho(g)$ . The general properties of this (twisted) Euler class were discussed in [E-M].

Our main objective is to find universal bounds, depending on the field  $K \subset \mathbf{R}$  and the degree  $m$  only, for the order of the Euler class of  $K$ -representations of finite groups. We proceed by dealing first with  $p$ -groups and then (Section 3) with arbitrary finite groups.

2.2. We start with the following simple observation.

**PROPOSITION 2.1** *Let  $G$  be a finite  $p$ -group and let  $\rho: G \rightarrow \mathrm{GL}_m(K)$  be a representation of degree  $m \not\equiv 0 \pmod{\varphi_K(p)}$ . Then the Euler class of  $\rho$  is  $= 0$ .*

*Proof.* The assumption implies that  $\varphi_K(p) > 1$  and thus  $p$  odd ( $\varphi_K(2) = 1$ ). Let  $\rho = \bigoplus_{i=1}^n \rho_i$ , with  $\rho_i$  irreducible; then  $e(\rho) = e(\rho_1)e(\rho_2) \cdots e(\rho_n)$ . At least one of the  $\rho_i$  must have degree 1, for otherwise  $m$  would be divisible by  $\varphi_K(p)$  (Corollary 1.5). Thus the corresponding  $e(\rho_i)$  is 0 and whence  $e(\rho) = 0$ .

We may thus, for a  $p$ -group  $G$ , assume that the degree  $m$  of  $\rho$  is  $\equiv 0 \pmod{\varphi_K(p)}$ . It turns out that the situation is quite different according to whether  $\gamma_K(p)$  is finite or infinite.

Let  $m$  be even and  $\equiv 0 \pmod{\varphi_K(p)}$ , and assume  $\gamma_K(p) = \infty$ . Then no uniform bound can exist for the order of the Euler class of  $K$ -representations of  $p$ -groups. This will be illustrated by Corollary 2.4 below. We first prove a lemma concerning the cyclic group  $C_n$ .

**LEMMA 2.3.** *Let  $K \subset \mathbf{R}$  be an arbitrary real field. There exists, for any integer  $l > 0$ , a  $K$ -representation  $\rho$  of  $C_n$  of degree  $l\varphi_K(n)$  and with Euler class  $e(\rho)$  of (maximal possible) order  $n$ .*

*Proof.*  $C_n$  has a faithful irreducible representation  $\tau$  over  $K$  of degree  $m = \varphi_K(n)$  (its character is  $= \sum_{\sigma} \sigma\chi$ , where  $\chi$  is faithful  $\mathbf{C}$ -irreducible and  $\sigma$  varies through  $\text{Gal}(K(n)/K)$ ). For the Euler class  $e(\tau)$  one has  $e(\tau)^2 = \pm c_m(\tau \otimes \mathbf{C})$ , the top Chern class of  $\tau \otimes \mathbf{C}$ ; since  $\tau \otimes \mathbf{C}$  is a sum of  $m$  faithful one-dimensional  $\mathbf{C}$ -representations,  $c_m(\tau \otimes \mathbf{C})$  has order  $n$ , and so has  $e(\tau)$ . If we take for  $\rho$  the  $l$ -fold direct sum of such  $K$ -representations  $\tau$ , the order of  $e(\rho)$  will be  $n$  and the degree  $l \cdot \varphi_K(n)$ .

**COROLLARY 2.4.** *Let  $K \subset \mathbf{R}$ , and let  $p$  be a prime such that  $\gamma_K(p) = \infty$ . If  $m$  is even and  $m \equiv 0 \pmod{\varphi_K(p)}$ , then there exists an  $m$ -dimensional  $K$ -representation of  $C_{p^\alpha}$  with Euler class of order  $p^\alpha$ .*

*Proof.* If  $p$  is odd,  $\gamma_K(p) = \infty$  implies that  $\varphi_K(p) = \varphi_K(p^\alpha)$  for  $\alpha \geq 1$  and the result follows from Lemma 2.3. If  $p = 2$ ,  $\varphi_K(2^\alpha) = 2$  or  $1$  for  $\alpha \geq 1$ . Hence for any even  $m$  one can find a  $K$ -representation of  $C_{2^\alpha}$  of degree  $m$  and Euler class of order  $2^\alpha$  (cf. Lemma 2.3).

2.3. We now turn to the case  $\gamma_K(p) < \infty$ , where the situation is different.

**THEOREM 2.5.** *Let  $K$  be a subfield of  $\mathbf{R}$  and  $p$  a prime with  $\gamma = \gamma_K(p) < \infty$ . For any finite  $p$ -group  $G$  and any  $K$ -representation  $\rho: G \rightarrow \text{GL}_m(K)$  the Euler class  $e(\rho) \in H^m(G; \mathbf{Z}(\rho))$  satisfies*

$$p^\gamma m e(\rho) = 0.$$

*Proof.* We first assume that  $\rho$  is irreducible. According to Theorem 1.2 we distinguish two possibilities.

(a)  $\rho$  factors as  $G \rightarrow \bar{G} \xrightarrow{\bar{\rho}} \text{GL}_m(K)$  where  $\bar{G}$  is one of the  $p$ -groups of special type and  $\bar{\rho}$  faithful. If  $\bar{G}$  is of order  $p^\alpha$ ,  $\alpha \leq \gamma$  then plainly  $p^\gamma m e(\rho) = 0$ ;

thus we may assume  $\alpha > \gamma$ . If  $p$  is odd,  $\rho$  is of degree  $m = \varphi_K(p^\alpha) = \varphi_K(p) \cdot p^{\alpha-\gamma}$ , and hence  $p^\gamma me(\rho) = 0$ . In case  $p = 2$  and  $\alpha = \gamma + 1$ ,  $2^\alpha$  divides  $2^\gamma m$  for  $m$  even; thus  $2me(\rho) = 0$  (the case  $m$  odd is trivial, since then always  $2e(\rho) = 0$ ). It remains to consider the case  $p = 2$ ,  $\alpha \geq \gamma + 2$ . According to Proposition 1.4 the degree of  $\rho$  is then  $2^{\alpha-\gamma}$  for the groups  $C_{2^\alpha}$ ,  $Q_{2^\alpha}$ ; and  $2^{\alpha-\gamma-1}$  for  $D_{2^\alpha}$ ,  $2^{\alpha-\gamma}$  or  $2^{\alpha-\gamma-1}$  for  $SD_{2^\alpha}$ . For the first two groups,  $2^\gamma \cdot 2^{\alpha-\gamma} = 2^\alpha = |\bar{G}|$  annihilates  $e(\rho)$ , and for the latter ones  $2^\gamma \cdot 2^{\alpha-\gamma-1} = 2^{\alpha-1} = |\bar{G}|/2$  annihilates  $e(\rho)$  (since the cohomology of  $D_{2^\alpha}$  and  $SD_{2^\alpha}$  with  $\mathbf{Z}$ -coefficients contains no elements of order  $2^\alpha$ ).

(b)  $\rho$  is induced from  $\tau: H \rightarrow \mathrm{GL}_{m/p}(K)$ , where  $H \subset G$  is of index  $p$ . Let  $tr$  denote the cohomology transfer. The Euler class of the restriction  $\rho_H$  satisfies  $tr e(\rho_H) = pe(\rho)$ . Since we may assume by induction that  $p^\gamma(m/p)e(\tau) = 0$ , and since  $\rho_H$  is of the form  $\tau \oplus \nu$ , we infer  $p^\gamma(m/p)e(\rho_H) = p^\gamma(m/p)e(\tau)e(\nu) = 0$ . It follows that

$$p^\gamma me(\rho) = \mathrm{tr} \left( p^\gamma \frac{m}{p} e(\rho_H) \right) = 0.$$

We now assume that  $\rho$  is reducible,  $\rho = \rho_1 \oplus \rho_2 \oplus \cdots \oplus \rho_k$ , the  $\rho_i$  being  $K$ -irreducible. Then  $e(\rho) = e(\rho_1)e(\rho_2) \cdots e(\rho_k)$ , and

$$p^\gamma me(\rho) = p^\gamma m_1 e(\rho_1)e(\rho_2) \cdots e(\rho_k) + \cdots + p^\gamma m_k e(\rho_1)e(\rho_2) \cdots e(\rho_k)$$

where  $m_i$  is the degree of  $\rho_i$ . Since  $\rho_i$  is irreducible, we have  $p^\gamma m_i e(\rho_i) = 0$ , and thus  $p^\gamma me(\rho) = 0$ .

*Remark 2.6.* If  $m$  is even,  $m = \varphi_K(p)p^\beta \cdot f$  with  $(f, p) = 1$  and  $\gamma_K(p) = \gamma < \infty$ , then there exists a  $K$ -representation of  $C_{p^{\gamma+\beta}}$  of degree  $m$  with Euler class satisfying  $p^{\gamma-1}me(\rho) \neq 0$ . This follows immediately from Lemma 2.3.

### 3. Arbitrary finite groups

3.1. We define for a subfield  $K$  of  $\mathbf{R}$  and an integer  $m > 0$ , an additive operator  $\mathcal{E}_K(m)$  on finite Abelian groups. If  $m$  is odd,  $\mathcal{E}_K(m): A \rightarrow A$  is multiplication by 2. For  $m$  even,  $\mathcal{E}_K(m)$  is given by its action on  $p$ -torsion groups as follows.

- (1)  $\mathcal{E}_K(m)$  is the identity on  $p$ -torsion, if  $m \not\equiv 0 \pmod{\varphi_K(p)}$ .
- (2)  $\mathcal{E}_K(m)$  is zero on  $p$ -torsion if  $m \equiv 0 \pmod{\varphi_K(p)}$  and  $\gamma_K(p) = \infty$ .
- (3)  $\mathcal{E}_K(m)$  is multiplication by  $p^{\gamma+\alpha}$  on  $p$ -torsion, if  $m \equiv 0 \pmod{\varphi_K(p)}$ ,  $\gamma = \gamma_K(p) < \infty$  and  $m = p^\alpha \cdot f$ ,  $f$  prime to  $p$ .

For instance, if  $K = \mathbf{R}$ , then  $\mathcal{E}_{\mathbf{R}}(m)$  is the zero operator for all even  $m$ .

If  $K$  is a field such that  $\gamma_K(p) < \infty$  for all  $p$ , and if only finitely many  $\varphi_K(p)$  divide  $m$ , we define a numerical function by

$$E_K(m) = \text{lcm}\{n \mid m \equiv 0 \pmod{\varphi_K(n)}\}.$$

Note that, for any prime  $p$ ,  $p^\beta$  divides  $E_K(m)$  if and only if  $m \equiv 0 \pmod{\varphi_K(p^\beta)}$ . In one direction this is part of the definition; conversely, if  $p^\beta$  divides  $E_K(m)$  there is an  $n$  divisible by  $p^\beta$  with  $m \equiv 0 \pmod{\varphi_K(n)}$  and thus, since  $\varphi_K(p^\beta)$  divides  $\varphi_K(n)$ , one has  $m \equiv 0 \pmod{\varphi_K(p^\beta)}$ . The prime decomposition of  $E_K(m)$  is now obtained as follows, for  $K \subset \mathbf{R}$  and  $m$  even ( $E_K(m) = 2$  if  $m$  is odd):

Let  $m = \prod p^{\nu_p}$  be the decomposition of  $m$  into powers of different primes. By Lemma 1.3,  $\varphi_K(p^{\beta+\gamma}) = \varphi_K(p)p^\beta$  ( $\gamma = \gamma_K(p)$ ,  $\beta \geq 0$  in case  $p$  odd, and  $\beta \geq 1$  if  $p = 2$ ); thus, for a prime  $p$  with  $m \equiv 0 \pmod{\varphi_K(p)}$ ,  $m$  even, the greatest power dividing  $E_K(m)$  is  $p^{\nu_p+\gamma}$ . We thus have

**PROPOSITION 3.1.** *If for  $K \subset \mathbf{R}$  and  $m = \prod p^{\nu_p}$  the integer  $E_K(m)$  is defined, then*

$$E_K(m) = 2 \text{ if } m \text{ is odd,}$$

$$E_K(m) = \prod' p^{\nu_p+\gamma_K(p)} \text{ if } m \text{ is even, the product } \prod' \text{ being taken over all those primes } p \text{ for which } m \equiv 0 \pmod{\varphi_K(p)}.$$

*Remarks.* (1) If  $K = \mathbf{Q}$ ,  $E_{\mathbf{Q}}(m) = E_m$ , the numerical function considered in [E-M] (which is equal to the denominator of  $B_m/m$ ,  $m$  even).

(2)  $E_K(m)$  is defined for all  $m$  if  $K$  is an algebraic number field.

**COROLLARY 3.2.** *If for  $K \subset \mathbf{R}$  the integer  $E_K(m)$  is defined, then the operator  $\mathcal{E}_K(m): A \rightarrow A$  differs from “multiplication with  $E_K(m)$ ” only by a canonical automorphism of  $A$ . In particular,  $\mathcal{E}_K(m)$  and multiplication by  $E_K(m)$  have the same kernel.*

We will make use later on of the following special case.

**COROLLARY 3.3.** *Let  $K = \mathbf{Q}(4n) \cap \mathbf{R}$  and  $p$  a prime dividing  $4n$ . Then for even  $m$  the operator  $\mathcal{E}_K(m)$  has the same kernel on any  $p$ -torsion group as multiplication by  $2nm$ .*

*Proof.* Let  $m = \prod p^{\nu_p(m)}$  and  $n = \prod p^{\nu_p(n)}$  be the prime decompositions. Since  $p \mid 4n$  we have, for  $p$  odd,  $\varphi_K(p) = 2$ . Further we have  $\gamma_K(p) = \nu_p(n)$  for  $p$  odd and  $\gamma_K(2) = \nu_2(n) + 1$ . Hence, for  $m$  even,  $\mathcal{E}_K(m)$  acts on  $p$ -torsion by multiplication with  $p^{\nu_p(n)+\nu_p(m)}$  if  $p$  is odd, and with  $2^{\nu_2(n)+1+\nu_2(m)}$  if  $p = 2$ . Thus the kernel of  $\mathcal{E}_K(m)$  on  $p$ -torsion is the same as the kernel of multiplication by  $2nm$ .

3.2. We now state and prove our main theorem.

**THEOREM 3.4.** *Let  $K \subset \mathbf{R}$  be a real field and  $\rho: G \rightarrow GL_m(K)$  a  $K$ -representation of degree  $m$  of a finite group  $G$ . Then the Euler class  $e(\rho) \in H^m(G; \mathbf{Z}(\rho))$  satisfies*

$$\mathcal{E}_K(m)e(\rho) = 0.$$

*In particular, if  $E_m(K)$  is defined (e.g., if  $K$  is a number field) the order of  $e(\rho)$  divides  $E_K(m)$ .*

*Proof.* Let  $G(p)$  denote a  $p$ -Sylow subgroup of  $G$ . Since the cohomology restriction from  $G$  to  $G(p)$  is injective on the  $p$ -primary component, it suffices to prove the theorem in the case where  $G$  is a  $p$ -group. If  $m$  is odd,  $2e(\rho) = \mathcal{E}_K(m)e(\rho) = 0$ . If  $m$  is even and  $m \not\equiv 0 \pmod{\varphi_K(p)}$ ,  $e(\rho) = 0$  by Proposition 2.1. It remains to consider the case  $m$  even,  $m \equiv 0 \pmod{\varphi_K(p)}$ : If  $\gamma_K(p) = \infty$ , then  $\mathcal{E}_K(m)e(\rho) = 0$  by definition of  $\mathcal{E}_K(m)$ . If  $\gamma = \gamma_K(p) < \infty$ , we have  $p^\gamma me(\rho) = 0$  by Theorem 2.5; since, for a  $p$ -group  $G$ ,  $p^\gamma me(\rho)$  and  $p^{\gamma + \nu_p(m)}e(\rho)$  have the same order, we infer  $\mathcal{E}_K(m)e(\rho) = 0$ . In case  $E_K(m)$  is defined,  $E_K(m)e(\rho) = 0$  by Corollary 3.2.

**Remark 3.5.** The operator  $\mathcal{E}_K(m)$  in Theorem 3.4 is best possible in the following obvious sense. Suppose  $\mathcal{E}'_K(m)$  is another such operator (i.e., a natural transformation of the identity functor on the category of finite Abelian groups, such that  $\mathcal{E}'_K(m)e(\rho) = 0$  for all  $K$ -representations  $\rho$  of degree  $m$  of finite groups) then

$$\ker(\mathcal{E}_K(m): A \rightarrow A) \subset \ker(\mathcal{E}'_K(m): A \rightarrow A) \quad (*)$$

for all finite Abelian groups  $A$ . In order to prove this we observe that it suffices to check (\*) in case  $A$  is a cyclic  $p$ -group; for that case (\*) is an easy consequence of Lemma 2.3 and Remark 2.6 together with the definition of  $\mathcal{E}_K(m)$ .

In particular, if  $K$  is a number field, we obtain the following.

**COROLLARY 3.6.** *Let  $K \subset \mathbf{R}$  be a number field. Then the least common multiple of the orders of the Euler classes  $e(\rho)$ , where  $\rho$  ranges over all  $K$ -representations of degree  $m$  of finite groups, is equal to  $E_K(m) = \text{lcm}\{n \mid m \equiv 0 \pmod{\varphi_K(n)}\}$ .*

**3.3.** If a representation  $\rho: G \rightarrow GL_m(\mathbf{R})$  is not known to be realizable over some subfield  $K \subset \mathbf{R}$  fixed in advance, one can still obtain a bound on the order of  $e(\rho)$ , depending on the character field  $\mathbf{Q}(\chi)$  (i.e. the field obtained from  $\mathbf{Q}$  by adjoining the values of the character  $\chi$  of  $\rho$ ). We need first the following lemma.

**LEMMA 3.7.** *Let  $\rho: G \rightarrow \mathrm{GL}_m(\mathbf{R})$  be a real representation of a finite  $p$ -group  $G$ . Then  $\rho$  is equivalent to a representation defined over  $\mathbf{Q}(\chi)$ , where  $\chi$  is the character of  $\rho$ .*

*Proof.* If  $p$  is odd, all  $\mathbf{C}$ -irreducible characters  $\psi$  of  $G$  have Schur index 1 over  $\mathbf{Q}$ , and therefore  $\rho$  is defined over  $\mathbf{Q}(\chi)$  (cf. [R]). In case  $p = 2$ , the Schur index  $s_{\mathbf{Q}}(\psi)$  is one or two; by [F; Prop. 4.2],  $s_{\mathbf{Q}}(\psi) = s_{\mathbf{R}}(\psi)$  and therefore  $s_K(\psi) = s_{\mathbf{R}}(\psi)$  for any subfield  $K \subset \mathbf{R}$ . It follows that an  $\mathbf{R}$ -representation of a 2-group whose character takes values in  $K \subset \mathbf{R}$ , is realizable over  $K$ .

**THEOREM 3.8.** *Let  $\rho: G \rightarrow \mathrm{GL}_m(\mathbf{R})$  be a real representation of an arbitrary finite group  $G$ . Then the Euler class  $e(\rho)$  satisfies*

$$E_{\mathbf{Q}(\chi)}(m)e(\rho) = 0$$

where  $\mathbf{Q}(\chi)$  denotes the field obtained from  $\mathbf{Q}$  by adjoining the values of the character  $\chi$  of  $\rho$ .

*Proof.* Let  $\rho'$  denote the restriction of  $\rho$  to a  $p$ -Sylow subgroup of  $G$ , and denote by  $\chi'$  the character of  $\rho'$ . From Theorem 3.4 and Lemma 3.7 we infer that  $E_{\mathbf{Q}(\chi')}e(\rho') = 0$  and, as  $\mathbf{Q}(\chi') \subset \mathbf{Q}(\chi)$ ,  $E_{\mathbf{Q}(\chi)}e(\rho') = 0$ . The assertion of the theorem now follows, since the cohomology restriction from  $G$  to a  $p$ -Sylow subgroup is injective on the  $p$ -primary component.

3.4. Using Corollary 3.3 we can get bounds for the order of Euler classes of arbitrary real representations, in terms of the exponent of  $G$ .

**THEOREM 3.9.** *Let  $G$  be a finite group of exponent  $\exp(G)$  and  $\rho: G \rightarrow \mathrm{GL}_m(\mathbf{R})$  a real representation of even degree  $m$ . Then the Euler class satisfies*

$$\frac{m}{2} \exp(G)e(\rho) = 0.$$

*Proof.* Since the cohomology restriction from  $G$  to a  $p$ -Sylow subgroup is injective on the  $p$ -primary component, we may assume that  $G$  is a  $p$ -group. Then  $\rho$  is realizable over  $\mathbf{Q}(\exp(G)) \cap \mathbf{R}$  since the character of  $\rho$  takes its values in that field (Lemma 3.7). If  $p$  is odd, we apply Corollary 3.3 with  $K = \mathbf{Q}(4 \exp(G)) \cap \mathbf{R}$  and obtain  $2m \exp(G)e(\rho) = 0$ ; thus  $(m/2) \exp(G)e(\rho) = 0$ ,  $e(\rho)$  being a  $p$ -torsion element. If  $p = 2$  and  $\exp(G) \leq 2$ , then  $G$  is an elementary Abelian 2-group and thus even  $\exp(G)e(\rho) = 0$ . If  $p = 2$  and  $\exp(G) = 4n \geq 4$ , we infer from Corollary 3.3 (with  $K = \mathbf{Q}(4n) \cap \mathbf{R}$ ) that  $2nme(\rho) = 0$ , whence the assertion.

## REFERENCES

- [E-M] ECKMANN, B. and MISLIN, G., *Rational representations of finite groups and their Euler class*. Math. Annalen 245 (1979), 45–54.
- [F] FONTAINE, J. M., *Sur la décomposition des algèbres de groupes*. Ann. Scient. Ec. Norm. Sup. 4<sup>e</sup> Série t. 4 (1971) 121–180.
- [R] ROQUETTE, P., *Realisierungen von Darstellungen endlicher nilpotenter Gruppen*. Arch. Math. 9 (1958) 211–250.

*E. T. H. Zürich, Switzerland*

Received September 18, 1979.