

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 55 (1980)

Artikel: Factorisation is not unique for higher dimensional knots.
Autor: Bayer, Eva
DOI: <https://doi.org/10.5169/seals-42397>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Factorisation is not unique for higher dimensional knots

EVA BAYER*

0. Introduction

An n -knot will be a smooth oriented submanifold K of the $(n+2)$ -sphere S^{n+2} , where K is homeomorphic to S^n . A knot is irreducible if it cannot be written as a connected sum of two non-trivial knots. Schubert has shown that every 1-knot can be written uniquely as a connected sum of finitely many irreducible knots (see [S] or [K1, Section 1]). For $n > 2$, Sosinskii has proved that it is still possible to factorise every n -knot into finitely many irreducible knots (cf. [So Theorem 5. 1] or [K1, Section 2]) but Kearton has shown that this factorisation is not necessarily unique for $n = 3$ [K]. In the present note we shall prove the non uniqueness of the factorisation for $(2q-1)$ -knots, $q \geq 3$ and for $(2q)$ -knots, $q \geq 4$.

I would like to thank C. Weber for advising me to consider Levine duality in the even dimensional case. I also thank M. Kervaire for useful conversations.

1. Factorisation is not necessarily unique for $(2q-1)$ -knots, $q \geq 3$

Let $q \geq 3$ be an integer.

DEFINITION. A *Seifert matrix* A is a square matrix of integers such that $\det(A + (-1)^q A^t) = \pm 1$, where A^t is the transpose of A .

Let A be a non-singular Seifert matrix (that is, $\det(A) \neq 0$). We shall say that A is *irreducible* if A is not S -equivalent to the orthogonal sum of two non-singular Seifert matrices. (See [Le] for the definition of S -equivalence. In the examples that we shall construct, the Seifert matrices will be unimodular, and unimodular Seifert matrices are S -equivalent if and only if they are integrally congruent (see [T, Proposition 4.3])). We shall use the notation

$$S = A + (-1)^q A^t, \quad z = S^{-1} A$$

* Supported by a grant from the Alexander von Humboldt Foundation.

LEMMA 1. *Let A_1 and A_2 be Seifert matrices with $z_1 = z_2$. Then*

$$\begin{pmatrix} A_1 & 0 \\ 0 & -A_1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} A_2 & 0 \\ 0 & -A_2 \end{pmatrix}$$

are integrally congruent.

LEMMA 2. *There exist irreducible Seifert matrices A_1 and A_2 such that*

- a) $z_1 = z_2$
- b) A_1 and A_2 are not S -equivalent
 A_1 and $-A_2$ are not S -equivalent.

(We shall give explicit examples of such Seifert matrices after the proof of this lemma.)

The above two lemmas give the desired result. Indeed, let A_1 and A_2 be Seifert matrices as in Lemma 2.

Levine has shown that the S -equivalence classes of non-singular Seifert matrices correspond biunivoguely to the isotopy classes of simple $(2q-1)$ -knots [Le, Theorems 1, 2, 3]. Note that this implies that irreducible Seifert matrices correspond to irreducible knots. Let K_1, L_1, K_2, L_2 be the simple $(2q-1)$ -knots corresponding to $A_1, -A_1, A_2, -A_2$ respectively. These knots are irreducible, because A_1 and A_2 are irreducible. By Lemma 1,

$$\begin{pmatrix} A_1 & 0 \\ 0 & -A_1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} A_2 & 0 \\ 0 & -A_2 \end{pmatrix}$$

are integrally congruent, as $z_1 = z_2$. Therefore they are S -equivalent. So by [Le, Theorem 3] the connected sum of K_1 and L_1 is isotopic to the connected sum of K_2 and L_2 . On the other hand, [Le, Theorem 1] shows that K_1 is not isotopic either to K_2 or to L_2 , as the Seifert matrices are not S -equivalent.

Proof of Lemma 1. Let A be a Seifert matrix, and let

$$M_1 = \begin{pmatrix} A & 0 \\ 0 & -A \end{pmatrix}, \quad M_2 = \begin{pmatrix} 0 & (-1)^q(1-z^t) \\ z & 0 \end{pmatrix}.$$

Then M_1 and M_2 are integrally congruent. Indeed, let

$$X = \begin{pmatrix} 1-z & (-1)^q S^{-1} \\ -z & (-1)^q S^{-1} \end{pmatrix} = \begin{pmatrix} I & I \\ 0 & I \end{pmatrix} \begin{pmatrix} I & 0 \\ 0 & (S^{-1})^t \end{pmatrix} \begin{pmatrix} I & 0 \\ (-1)^{q+1} A & I \end{pmatrix}$$

where I is the identity matrix.

One checks by direct computation that $M_2 = X' M_1 X$ (it is useful to note that $1 - z' = AS^{-1}$, and that $(1 - z')A = Az$). This proves Lemma 1, as $z_1 = z_2 = z$,

$$\begin{pmatrix} A_1 & 0 \\ 0 & -A_1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} A_2 & 0 \\ 0 & -A_2 \end{pmatrix}$$

are both congruent to

$$\begin{pmatrix} 0 & (-1)^q (1 - z') \\ z & 0 \end{pmatrix}.$$

Proof of Lemma 2. Let ϕ be the cyclotomic polynomial corresponding to the 15th roots of unity. Let t be the Jordan matrix associated with ϕ , and let $z = (1 - t)^{-1}$. Note that $\det(z) = 1$: indeed, $\det(1 - t) = \phi(1) = 1$.

Let ζ be a primitive 15th root of unity. Sending ζ to ζ^{-1} induces a non-trivial involution on $\mathbf{Z}[\zeta]$. We shall denote this involution by an overbar.

Let $\Delta = \{x \in \mathbf{Q}(\zeta) \mid \text{Tr}_{\mathbf{Q}(\zeta)/\mathbf{Q}}(x\mathbf{Z}[\zeta]) \subset \mathbf{Z}\}$ be the inverse different of the extension $\mathbf{Q}(\zeta)/\mathbf{Q}$. We have: $\bar{\Delta} = \Delta$.

DEFINITION. Let V be a torsion free $\mathbf{Z}[\zeta]$ -module of finite rank. We shall say that a hermitian or skewhermitian form

$$h: V \times V \rightarrow \Delta$$

is *unimodular*, if

$$\begin{aligned} \text{ad}(h): V &\rightarrow \text{Hom}_{\mathbf{Z}[\zeta]}(V, \Delta) \\ x &\rightarrow h(_, x) \end{aligned}$$

is an isomorphism.

Claim 1. The integral congruence classes of Seifert matrices A such that

$$(A + (-1)^q A^t)^{-1} A = z \tag{1}$$

($z = (1 - t)^{-1}$ as above, fixed; q a fixed integer) are in bijection with the isometry classes of $(-1)^q$ -hermitian unimodular forms

$$h: \mathbf{Z}[\zeta] \times \mathbf{Z}[\zeta] \rightarrow \Delta.$$

Proof of Claim 1. Let A be a Seifert matrix with property (1), and let $S = A + (-1)^q A^t$. Rank $(A) = \text{degree}(\phi) = 8$. Let V be a free \mathbf{Z} -module of rank 8. We can consider S as a $(-1)^q$ -symmetric form $S: V \times V \rightarrow \mathbf{Z}$, and $t = 1 - z^{-1}: V \rightarrow V$ will be an isometry for S .

Setting $\zeta \cdot v = t(v)$ for v in V makes V into a $\mathbf{Z}[\zeta]$ -module. As t corresponds to the Jordan matrix of ϕ , V is isomorphic to $\mathbf{Z}[\zeta]$.

As in [B-M, §1], we associate to S a $(-1)^q$ -hermitian form

$$h: \mathbf{Z}[\zeta] \times \mathbf{Z}[\zeta] \rightarrow \Delta$$

such that

$$\text{Tr}_{\Phi(\zeta)/\Phi} h(\alpha x, y) = S(\alpha x, y) \quad \forall \alpha \in \Phi(\zeta) \quad \forall x, y \in V. \quad (2)$$

It is easy to check that h is unimodular and that congruent Seifert matrices determine isometric $(-1)^q$ -hermitian forms.

Conversely, given a unimodular $(-1)^q$ -hermitian form $h: \mathbf{Z}[\zeta] \times \mathbf{Z}[\zeta] \rightarrow \Delta$, the formula (2) determines a $(-1)^q$ -symmetric matrix S such that $\det(S) = \pm 1$ and t is an isometry for S . Set $A = Sz$. Then $A + (-1)^q A^t = S$, therefore A is a Seifert matrix satisfying (1). Isometric $(-1)^q$ -hermitian forms determine congruent Seifert matrices.

Claim 2. The isometry classes of unimodular $(-1)^q$ -hermitian forms

$$h: \mathbf{Z}[\zeta] \times \mathbf{Z}[\zeta] \rightarrow \Delta$$

are in bijection with $U_0/N(U)$, where U is the group of units of $\mathbf{Z}[\zeta]$, U_0 is the group of units of $\mathbf{Z}[\zeta + \bar{\zeta}]$, and $N: U \rightarrow U_0$, $N(u) = u\bar{u}$, is the norm map.

Proof of Claim 2. Let g be the minimal polynomial of $\zeta + \bar{\zeta}$, and let

$$\alpha_0 = \frac{1}{g'(\zeta + \bar{\zeta})} \frac{1}{\zeta - \bar{\zeta}}.$$

Let Δ_1 be the inverse different of the extension $\mathbf{Q}(\zeta)/\mathbf{Q}(\zeta + \bar{\zeta})$, and Δ_2 the inverse different of $\mathbf{Q}(\zeta + \bar{\zeta})/\mathbf{Q}$. Then $\Delta = \Delta_1 \cdot \Delta_2$ [L, III. §1, Proposition 5] and

$$\Delta_1 = \frac{1}{\zeta - \bar{\zeta}} \mathbf{Z}[\zeta]$$

$$\Delta_2 = \frac{1}{g'(\zeta + \bar{\zeta})} \mathbf{Z}[\zeta + \bar{\zeta}]$$

[L, III. §1 Corollary of Proposition 2]. Therefore $\Delta = a_0 \mathbf{Z}[\zeta]$. Notice that $\bar{a}_0 = -a_0$.

Let $h: \mathbf{Z}[\zeta] \times \mathbf{Z}[\zeta] \rightarrow \Delta$ be a unimodular $(-1)^q$ -hermitian form. We have: $h(x, y) = ax\bar{y}$ for some a in Δ such that $\bar{a} = (-1)^q a$.

As we can identify $\text{Hom}_{\mathbf{Z}[\zeta]}(\mathbf{Z}[\zeta], \Delta)$ with Δ , the unimodularity of h implies that $a\mathbf{Z}[\zeta] = \Delta$. Therefore $a\mathbf{Z}[\zeta] = a_0\mathbf{Z}[\zeta]$. This implies that aa_0^{-1} is a unit. We have $\overline{aa_0^{-1}} = (-1)^{q+1}aa_0^{-1}$.

Set

$$u = \begin{cases} aa_0^{-1} & \text{if } q \text{ is odd} \\ aa_0^{-1}(\zeta - \bar{\zeta}) & \text{if } q \text{ is even} \end{cases} \quad (3)$$

$\zeta - \bar{\zeta}$ is a unit: $(\zeta - \bar{\zeta})^2(\zeta + \bar{\zeta})(-\zeta - \bar{\zeta} + 1) = 1$.

Therefore u is in U_0 in both cases. Conversely, to $u \in U_0$ we associate the $(-1)^q$ -hermitian form $h(x, y) = ax\bar{y}$ where a is given by (3). One checks easily that two $(-1)^q$ -hermitian forms are isometric if and only if the corresponding units are in the same class in $U_0/N(U)$.

Let us determine the cardinality of $U_0/N(U)$. We have

$$[U_0 : N(U)] = \frac{[U_0 : U_0^2]}{[N(U) : U_0^2]}.$$

Using the theorem of Dirichlet on the rank of the group of units, we see that $[U_0 : U_0^2] = 16$.

Let μ be the group of roots of unity in $\mathbf{Q}(\zeta)$. Then

$$[N(U) : U_0^2] = [U : \mu U_0] = Q$$

and $Q = 2$ [L1, Chap. 3, Theorem 4.1]. So $[U_0 : N(U)] = 8$. (We shall actually exhibit 8 distinct classes of $U_0/N(U)$ in the next section.)

Applying Claim 1 and Claim 2, we see that there are 8 non-congruent Seifert matrices A such that

$$(A + (-1)^q A^t)^{-1} A = z. \quad (1)$$

Therefore it is possible to choose A_1 and A_2 satisfying (1), and such that A_1 is not congruent either to A_2 or to $-A_2$. But congruence and S -equivalence are the same in this case, because the Seifert matrices are unimodular (see [T, Proposition 4.3]). A_1 and A_2 are irreducible, as their Alexander polynomial is irreducible.

Explicit examples

Let $\zeta = e^{2i\pi/15}$, and let $u_1 = 1$, $u_2 = \zeta + \zeta^{-1}$. We have $u_2(-u_2^3 + u_2^2 + 4u_2 - 4) = 1$, therefore u_2 is a unit. But u_2 is not in $N(U)$: indeed, u_2 is conjugate to $\zeta^7 + \zeta^{-7}$ which is negative. Clearly $-u_2$ is also negative, therefore not in $N(U)$. Using similar methods for the units $u_3 = \zeta^2 + \zeta^{-2}$, $u_4 = u_2 u_3 = \zeta + \zeta^{-1} + \zeta^3 + \zeta^{-3}$, we see that $u_1, -u_1, u_2, -u_2, u_3, -u_3, u_4, -u_4$ are all in different classes of $U_0/N(U)$. In the proof of Lemma 2 we have seen that the cardinality of $U_0/N(U)$ is 8, therefore we have a complete set of representants of $U_0/N(U)$.

Using the method given in the proof of Lemma 2, let us associate the Seifert matrices A_i to the units u_i , $i = 1 \cdots 4$.

Then the $\begin{pmatrix} A_i & 0 \\ 0 & -A_i \end{pmatrix}$ are all different factorisations of the same Seifert matrix B (see Lemma 1). Moreover, B has no other factorisations than these four. Direct computation gives the following matrices for A_1 and A_2 :

q odd:

$$A_1 = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 0 & -1 & -2 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & -1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ -1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ -2 & -1 & 0 & 1 & 1 & 1 & 1 & 1 \\ -2 & -2 & -1 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

$$A_2 = \begin{pmatrix} 2 & 2 & 2 & 2 & 1 & 0 & -2 & -3 \\ 2 & 2 & 2 & 2 & 2 & 1 & 0 & -2 \\ 2 & 2 & 2 & 2 & 2 & 2 & 1 & 0 \\ 1 & 2 & 2 & 2 & 2 & 2 & 2 & 1 \\ 0 & 1 & 2 & 2 & 2 & 2 & 2 & 2 \\ -2 & 0 & 1 & 2 & 2 & 2 & 2 & 2 \\ -3 & -2 & 0 & 1 & 2 & 2 & 2 & 2 \\ -4 & -3 & -2 & 0 & 1 & 2 & 2 & 2 \end{pmatrix}$$

q even

$$A_1 = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & -2 & -2 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & -2 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & -2 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 2 & 1 & 0 & 0 & 0 \end{pmatrix}$$

$$A_2 = \begin{pmatrix} 0 & 0 & 0 & -1 & -2 & -3 & -3 & -2 \\ 0 & 0 & 0 & 0 & -1 & -2 & -3 & -3 \\ 1 & 0 & 0 & 0 & 0 & -1 & -2 & -3 \\ 2 & 1 & 0 & 0 & 0 & 0 & -1 & -2 \\ 3 & 2 & 1 & 0 & 0 & 0 & 0 & -1 \\ 3 & 3 & 2 & 1 & 0 & 0 & 0 & 0 \\ 2 & 3 & 3 & 2 & 1 & 0 & 0 & 0 \\ 0 & 2 & 3 & 3 & 2 & 1 & 0 & 0 \end{pmatrix}$$

2. Factorisation is not necessarily unique for $(2q)$ -knots, $q \geq 4$

Let $q \geq 4$ be an integer. Let $\Lambda = \mathbf{Z}[t, t^{-1}]$, and let T be a finitely generated \mathbf{Z} -torsion Λ -module such that $(1-t): T \rightarrow T$ is an isomorphism.

DEFINITION. $L: T \times T \rightarrow \mathbf{Q}/\mathbf{Z}$ is a *Levine pairing* if L is \mathbf{Z} -bilinear, non-singular, $(-1)^{q+1}$ -symmetric, such that

$$L(tx, ty) = L(x, y) \quad \text{for } x, y \text{ in } T.$$

In [Le 1] Levine associates to every $(2q)$ -knot K a Levine pairing on the \mathbf{Z} -torsion part T of $H_q(\tilde{X})$, \tilde{X} being the maximal abelian cover of $X = S^{2q+2} \setminus K$. Isotopic knots have isometric pairings. He also shows that every Levine pairing can be realized by a simple $(2q)$ -knot [Le 1, Theorem 13.1]. Conversely, Kojima has shown that if $H_q(\tilde{X})$ is finite and 2-torsion free and if $q \geq 4$, then simple $(2q)$ -knots having isometric Levine pairings are isotopic [Ko, Theorem 1]. Therefore, the following examples determine simple $(2q)$ -knots which factorise in more than one way:

q odd

Let $T = \mathbf{Z}/5$, and let $t(x) = -x$ for x in T . Then $L_1, L_2: T \times T \rightarrow \mathbf{Q}/\mathbf{Z}$ given by $L_1(x, y) = \frac{1}{5}xy$, $L_2(x, y) = \frac{2}{5}xy$ are Levine pairings. Clearly L_1 is not isometric either to L_2 or to $-L_2$. But

$$\begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1/5 & 0 \\ 0 & 4/5 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1/5 \\ 1/5 & 0 \end{pmatrix} = \begin{pmatrix} 4 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2/5 & 0 \\ 0 & 3/5 \end{pmatrix} \begin{pmatrix} 4 & 1 \\ 1 & 1 \end{pmatrix},$$

and the isomorphisms obviously commute with $t \oplus t$.

q even

Let $T = \mathbf{Z}/5 \oplus \mathbf{Z}/5$, $t: T \rightarrow T$ given by the matrix

$$\begin{pmatrix} 0 & -1 \\ 1 & -2 \end{pmatrix}$$

and let $L_1, L_2: T \times T \rightarrow \mathbf{Q}/\mathbf{Z}$ be the Levine pairings given by the matrices

$$\begin{pmatrix} 0 & 1/5 \\ 4/5 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 2/5 \\ 3/5 & 0 \end{pmatrix}$$

respectively.

L_1 is not Λ -isometric either to L_2 or to $-L_2$. Indeed, suppose that L_1 is Λ -isometric to $\varepsilon \cdot L_2$, for $\varepsilon = +1$ or -1 , and let X be the matrix corresponding to this isometry. Then $\det(X) = 2\varepsilon$.

Let

$$X = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

The relation $tX = Xt$ implies $\det(X) = (a+b)^2$. But $(a+b)^2 = 2\varepsilon$ is impossible. $L_1 \oplus -L_1$ and $L_2 \oplus -L_2$ are both Λ -isometric to

$$\begin{pmatrix} 0 & 0 & 0 & 2/5 \\ 0 & 0 & 3/5 & 0 \\ 0 & 2/5 & 0 & 0 \\ 3/5 & 0 & 0 & 0 \end{pmatrix}$$

the isometries are given by

$$\begin{pmatrix} I & I \\ -I & I \end{pmatrix} = \begin{pmatrix} I & I \\ 0 & I \end{pmatrix} \begin{pmatrix} I & 0 \\ 2I & I \end{pmatrix} \begin{pmatrix} 2I & 0 \\ 0 & I \end{pmatrix}$$

and

$$\begin{pmatrix} 3I & I \\ 2I & I \end{pmatrix} = \begin{pmatrix} I & I \\ 0 & I \end{pmatrix} \begin{pmatrix} I & 0 \\ 2I & I \end{pmatrix}$$

respectively, with $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

REFERENCES

- B-M E. BAYER and F. MICHEL, *Finitude du nombre des classes d'isomorphisme des structures isométriques entières*, Comment. Math. Helv. 54 (1979), 378–396.
- K C. KEARTON, *Factorisation is not unique for 3-knots*, Indiana Univ. Math. Jour. 28 (1979), 451–452.
- K1 —, *The factorisation of knots* (preprint).
- Ko S. KOJIMA, *Classification of simple knots by Levine pairings*, Comment. Math. Helv. 54 (1979) 356–367.
- L S. LANG, *Algebraic number theory*, Addison-Wesley Publishing Company
- L1 —, *Cyclotomic fields*, Springer-Verlag (1978).
- Le J. LEVINE, *An algebraic classification of some knots in codimension two*, Comment. Math. Helv. 45 (1970), 185–198.
- Le1 —, *Knot modules*, Trans. of the A.M.S. 229 (1977), 1–50.
- S H. SCHUBERT, *Die eindeutige Zerlegbarkeit eines Knotens in Primknoten*, S.-B. Heidelberger Akad. Wiss. Math.-Natur. Kl. 1, 3 (1949), 57–104.

- So A. B. SOSINSKII, *Decomposition of knots*, Math. USSR Sbornik 10 (1970), 139–150.
T H. F. TROTTER, *Homology of group systems with applications to knot theory*, Ann. of Math. 76 (1962), 464–498.

Mathematisches Institut der Universität Münster
Federal Republic of Germany

Current address:

Université de Genève, Section de Mathématiques
2–4 rue du Lièvre
1211 Genève 24
Switzerland

Received May 15, 1980.