Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 57 (1982)

Artikel: A remark on two dimensional periodic potentials.
Autor: Dahlberg, B.E.J. / Trubowitz, E.
DOI: https://doi.org/10.5169/seals-43877

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 06.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Comment. Math. Helvetici 57 (1982) 130–134 0010–2571/82/001130–05$01.50+0.20/0
© 1982 Birkhäuser Verlag, Basel

A remark on two dimensional periodic potentials

B. E. J. DAHLBERG[1] and E. TRUBOWITZ[1,2]

1. Introduction

Let $A = a_1\mathbf{Z} \oplus a_2\mathbf{Z}$ be the lattice generated by the linearly independent vectors $a_1, a_2 \in \mathbf{R}^2$. Suppose q is a bounded, real-valued function periodic with respect to A. It is known that (see [2] and [5]) the spectrum of the self-adjoint operator $-\Delta + q$ acting on $L^2(\mathbf{R}^2)$ is purely absolutely continuous and is the union of the bands $B_n(q)$ $(n \geqslant 1)$. The nth band $B_n(q) \subset \mathbf{R}^1$ $(n \geqslant 1)$ is by definition the image of the function $\lambda_n(\cdot, q)$ defined on $\mathbf{R}^2$, where $\lambda_n(k, q)$ $(k \in \mathbf{R}^2)$ is the nth eigenvalue (counting with multiplicity) of the boundary value problem

$$(-\Delta + q)f = \lambda f,$$
$$f(x + a_j) = e^{ik \cdot a_j} f(x), \qquad j = 1, 2, \ldots$$

Let $|\mathfrak{F}|$ be the area of a fundamental cell $\mathfrak{F}$ for the lattice B dual to A.

THEOREM. *There exist positive constants $c(q)$, $C(q)$ such that for all $n \geqslant N(q)$*

$$\left[\frac{|\mathfrak{F}|}{\pi} n - cn^{1/4}, \frac{|\mathfrak{F}|}{\pi} n + cn^{1/4}\right] \subset B_n(q) \subset \left[\frac{|\mathfrak{F}|}{\pi} n - Cn^{1/3}, \frac{|\mathfrak{F}|}{\pi} n + Cn^{1/3}\right].$$

Suppose $q(x)$ is a bounded, real-valued, periodic function on $\mathbf{R}^1$. Then the spectrum of $-(d^2/dx^2) + q(x)$ acting on $L^2(\mathbf{R}^1)$ is purely absolutely continuous and its complement generically consists of an infinite number of intervals tending to $+\infty$ [see [1], p. 161]; i.e., there are infinitely many gaps. In two dimensions, however, we have the following immediate consequence of the Theorem.

COROLLARY. *The spectrum of $-\Delta + q$ contains a ray $[\lambda^*, \infty)$. In particular, there are at most a finite number of gaps.*

---

[1]The present work has been completed at the Forschungsinstitut für Mathematik, ETH Zürich.
[2]Research supported by the Sloan Foundation and NSF Grant No. MCS-8002561.

The corollary has been announced in [3] for the special case of rational lattices.

2. Proofs

Let $N_B(r, k)$ be the number of lattice points of B in the *open* disk of radius r centered at $k \in \mathbf{R}^2$. Denote by c, C generic positive constants.

LEMMA 1. *For every $\delta > 0$ there is a constant $c = c(\delta) > 0$ such that*

$$\sup_{k \in \mathbf{R}^2} N_B(r, k) \geq \frac{\pi}{|\mathfrak{F}|} r^2 + cr^{1/2},$$

$$\inf_{k \in \mathbf{R}^2} N_B(r, k) \leq \frac{\pi}{|\mathfrak{F}|} r^2 - cr^{1/2}.$$

for all $r \geq \delta$.

Proof. Let h_r be the characteristic function of the open disk $\{k \in \mathbf{R}^2 : |k| < r\}$ and set $E_B(r, k) = N_B(r, k) - (\pi/|\mathfrak{F}|)r^2$. If $a \in A \setminus \{0\}$ we have

$$\begin{aligned}\int_{\mathfrak{F}} E_B(r, k) e^{-ik.a}\, dk &= \int_{\mathfrak{F}} \left(-\pi \frac{r^2}{|\mathfrak{F}|} + \sum_{b \in B} h_r(k-b)\right) e^{-ik.a}\, dk \\ &= \int_{\mathbf{R}^2} h_r(k) e^{-ik.a}\, dk \\ &= \frac{2\pi r}{|a|} J_1(r\,|a|)\end{aligned}$$

where J_1 is the Bessel function of order 1. A similar calculation yields

$$\int_{\mathfrak{F}} E_B(r, k)\, dk = 0.$$

It follows that for all $a \in A \setminus \{0\}$

$$\begin{aligned}\int_{\mathfrak{F}} |E_B(r, k)|\, dk &\geq \max\left(\left|\int_{\mathfrak{F}} E_B(r, k) e^{-ik\cdot a}\, dk\right|, \left|\int_{\mathfrak{F}} E_B(r, k) e^{-2ik\cdot a}\, dk\right|\right) \\ &\geq 2\pi r \max\left(\frac{|J_1(r\,|a|)|}{|a|}, \frac{|J_1(2r\,|a|)|}{2\,|a|}\right) \\ &\geq (8\pi r)^{1/2} \max\left(\frac{|\sin(r\,|a| - \pi/4)|}{|a|^{3/2}}, \frac{|\sin(2r\,|a| - \pi/4)|}{|2a|^{3/2}}\right) \\ &\quad - Cr^{-1/2}\,|a|^{-5/2} \\ &\geq cr^{1/2}|a|^{-3/2} - Cr^{-1/2}|a|^{-5/2}.\end{aligned}$$

The asymptotic estimate

$$J_1(x)=\left(\frac{2}{\pi x}\right)^{1/2}\sin\left(x-\frac{\pi}{4}\right)+O(x^{-3/2})$$

was used in the third line and

$$\inf_{t\in\mathbf{R}}\max\left(\left|\sin\left(t-\frac{\pi}{4}\right)\right|,\frac{|\sin(2t-\pi/4)|}{\sqrt{8}}\right)>0$$

in the fourth. Consequently, for any fixed $a^*\in A\setminus\{0\}$ there is an r^* such that

$$\int_{\mathfrak{F}}|E_B(r,k)|\,dk\geqslant cr^{1/2}$$

for all $r\geqslant r^*$. Moreover, for all $0<\delta<1$ there is a $c(\delta)$ for which

$$\int_{\mathfrak{F}}|E_B(r,k)|\,dk\geqslant c(\delta)r^{1/2}$$

when $\delta<r<r^*$. This is achieved by picking an $a\in A$ with $|a|$ sufficiently large. Therefore,

$$\int_{\mathfrak{F}}E_B^+(r,k)\,dk=\int_{\mathfrak{F}}E_B^-(r,k)\,dk>cr^{1/2}.$$

Here, $E_B^+=\max(E_B,0)$ and $E_B^-=\min(-E_B,0)$.

The statement of Lemma 1 follows at once. □

Let $[x_n, y_n]$, $n\geqslant 1$, be the nth band corresponding to $q=0$.

LEMMA 2. *For*[1] $n\geqslant 1$

$$\sup_{k\in\mathbf{R}^2}N_B(\sqrt{x_n},k)\leqslant n\leqslant\inf_{k\in\mathbf{R}^2}N_B\left(\sqrt{y_n}+\frac{1}{\sqrt{y_n}},k\right)$$

Proof. Suppose $\sup_k N_B(\sqrt{x_n},k)\geqslant n+1$. Then there is a k such that $\lambda_{n+1}(k,0)<x_n$. This contradicts the definition of $\lambda_{n+1}(k,0)$. Similarly, suppose $\inf_k N_B(\sqrt{y_n}+1/\sqrt{y_n},k)\leqslant n-1$. Then there is a k such that $\lambda_n(k,0)>y_n$. But this contradicts the definition of $\lambda_n(k,0)$. □

[1]Observe that $0\leqslant x_n<y_n$, $n\geqslant 1$.

We can now prove the theorem.

Combining Lemmas 1, 2 with the obvious estimate $(\sqrt{y_n}+1/\sqrt{y_n})^{1/2}>y_n^{1/4}$, we have

$$\frac{\pi}{|\mathfrak{F}|}x_n+cx_n^{1/4}\leq n\leq\frac{\pi}{|\mathfrak{F}|}\left(y_n+2+\frac{1}{y_n}\right)-c\left(\sqrt{y_n}+\frac{1}{\sqrt{y_n}}\right)^{1/2}$$
$$\leq\frac{\pi}{|\mathfrak{F}|}y_n-c'y_n^{1/4}.$$

By similar arguments it is easy to show that

$$n\leq\sup_k N_B\left(\sqrt{x_n}+\frac{1}{\sqrt{x_n}},k\right)\leq cx_n$$

and

$$n\leq\sup_k N_B(\sqrt{y_n},k)\geq cy_n.$$

Consequently,

$$x_n\leq\frac{|\mathfrak{F}|}{\pi}n-cn^{1/4},\qquad\frac{|\mathfrak{F}|}{\pi}n+cn^{1/4}\leq y_n.$$

This proves the left hand inclusion of the theorem for $B_n(0)$. The statement for general q follows from the min-max characterization of $\lambda_n(k,q)$:

$$\lambda_n(k,q)=\sup_{\Phi}\inf_{\substack{\psi\in\Phi^\perp\\ \|\psi\|=1}}\langle\psi,(-\Delta+q)\psi\rangle$$

where $\Phi=\{\Phi_1,\ldots,\Phi_{n-1}\}\subset L^2(\mathbf{R}^2)$, $\Phi^\perp$ is the orthogonal complement of the span of Φ and $\psi(x+a_j)=e^{ik\cdot a_j}\psi(x)$, $j=1,2$. It gives

$$|\lambda_n(k,q)-\lambda_n(k,0)|\leq\|q\|_\infty$$

which yields the results.

The right hand inclusion is verified in almost the same way. The only significant difference is that Lemma 1 is replaced by the well known estimate

$$\left|N_B(r,k)-\frac{\pi}{|\mathfrak{F}|}r^2\right|\leq cr^{2/3}$$

one derives from the Poisson summation formula (c is independent of k). The proof of the theorem is finished.

Remark. The overlap of $B_n(0)$ and $B_{n+1}(0)$ is uniformly bounded away from 0; i.e., $|B_n(0) \cap B_{n+1}(0)| \geq c > 0$ for all $n \geq 1$. For

$$n \geq \sup_k N_B\left(\sqrt{x_{n+1}} - \frac{\varepsilon}{\sqrt{x_{n+1}}}, k\right)$$
$$\geq \frac{\pi}{|\mathfrak{F}|}\left(x_{n+1} - 2\varepsilon + \frac{\varepsilon^2}{x_{n+1}}\right) + c\left(\sqrt{x_{n+1}} - \frac{\varepsilon}{\sqrt{x_{n+1}}}\right)^{1/2}$$

and

$$n \leq \inf_k N\left(\sqrt{y_n} + \frac{\varepsilon}{\sqrt{y_n}}, k\right)$$
$$\leq \frac{\pi}{|\mathfrak{F}|}\left(y_n + 2\varepsilon + \frac{\varepsilon^2}{y_n}\right) - c\left(\sqrt{y_n} + \frac{\varepsilon}{\sqrt{y_n}}\right)^{1/2}.$$

Combining the inequalities and letting ε tend to 0 we find

$$x_{n+1} \leq y_n - c(y_n^{1/4} + x_{n+1}^{1/4}).$$

It is an immediate consequence that the spectrum of $-\Delta + q$ has no gaps at all when $|q|_\infty$ is sufficiently small. this reproves the result of [4].

BIBLIOGRAPHY

[1] McKean, H. P. and Trubowitz, E., *Hill's equation and hyperelliptic function theory in the presence of infinitely many branch points*, Comm. Pure Appl. Math. *29* (1976), 143–226.

[2] Odeh, F. and Keller, J. B., *Partial differential equations with periodic coefficients and Bloch waves in crystals*, J. Math. Phys., *5* (1964), 1499–1504.

[3] Skriganov, M. M., *Proof of the Bethe-Sommerfeld conjecture in dimension two*, Soviet Math., Dokl. *20* (1979), 956–959.

[4] Skriganov, M. M., *On the Bethe-Sommerfeld conjecture*, Soviet Math., Dolk. *20* (1979), 89–90.

[5] Thomas, L. E., *Time dependent approach to scattering from impurities in a crystal*, comm. Math. Phys. *33* (1973), 335–343.

Department of Mathematics
Uppsala University
Thunbergsv. 3
S-75238 Uppsala
Sweden

Courant Institute
New York University
251 *Mercer St.*
New York, NY 10012
U.S.A.

Received January 29, 1982