

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 57 (1982)

**Artikel:** Injectivity of local quasi-isometries.  
**Autor:** Gehring, F.W.  
**DOI:** <https://doi.org/10.5169/seals-43883>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.02.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Injectivity of local quasi-isometries

F. W. GEHRING<sup>(1)</sup>

### 1. Introduction

Suppose that  $E$  is a set in  $\bar{R}^n$ , the one point compactification of euclidean  $n$ -space  $R^n$ ,  $n \geq 2$ , and suppose that  $f$  is a mapping from  $E$  into  $\bar{R}^n$ . We say that  $f$  is an  $L$ -quasi-isometry in  $E$  if

$$\frac{1}{L} \leq \frac{|f(x_1) - f(x_2)|}{|x_1 - x_2|} \leq L \quad (1)$$

for each pair of points  $x_1, x_2 \in E - \{\infty\}$  and if  $f(\infty) = \infty$  whenever  $\infty \in E$ . We say that  $f$  is a local  $L$ -quasi-isometry in  $E$  if for each  $L' > L$  each  $x \in E$  has a neighborhood  $U$  such that  $f$  is an  $L'$ -quasi-isometry in  $E \cap U$ .

Suppose that  $f$  is a local  $L$ -quasi-isometry in a domain  $D$  in  $R^n$ . If  $L = 1$ , then  $f$  is an isometry in  $D$  and hence injective there. (See, for example, Theorem IV in [11].) Simple examples show that  $f$  need not be injective if  $L > 1$ . It was F. John who first noticed that for certain domains  $D$ ,  $f$  will be injective provided  $L$  is close enough to 1.

For each domain  $D \subset R^n$  we let  $L(D)$  denote the supremum of the numbers  $L \geq 1$  with the property that each local  $L$ -quasi-isometry in  $D$  is injective. We say that  $D$  is *rigid* if  $L(D) > 1$ .

John established the following interesting result in 1969 (Theorem A in [12]). See also [7] and [8].

**THEOREM 1.** *If  $D$  is an open ball or half space, then  $L(D) \geq 2^{1/4}$ .*

This result was generalized by John and then extended recently by Martio and Sarvas to a very broad class of domains. We say that  $D \subset R^n$  is a *uniform domain* if there exist constants  $a$  and  $b$  with the following property. Each pair of points

---

<sup>1</sup> This research was supported in part by grants from the U.S. National Science Foundation (Grant MCS 79-01713) and the Finnish Ministry of Education.

$x_1, x_2 \in D$  can be joined by a rectifiable arc  $\alpha$  in  $D$  so that

$$l(\alpha) \leq a |x_1 - x_2| \quad (2)$$

and so that for each  $x \in \alpha$

$$\min_{j=1,2} l(\alpha_j) \leq b d(x, \partial D), \quad (3)$$

where  $\alpha_1, \alpha_2$  denote the components of  $\alpha - \{x\}$ . Here  $l(\alpha)$  denotes the euclidean length of  $\alpha$  and  $d(x, \partial D)$  the distance from  $x$  to  $\partial D$ .

Martio and Sarvas showed that uniform domains are rigid by establishing the following result (Theorem 3.8 in [14]).

**THEOREM 2.** *If  $D$  is a uniform domain, then  $L(D) \geq c > 1$  where  $c$  depends only on the constants  $a$  and  $b$ .*

The present paper is concerned with the problem of identifying the domains in  $R^n$  which are rigid. In particular, we characterize in Section 2 the finitely connected plane domains which have this property. It turns out that each boundary component of such a domain is either a point or a quasicircle, that is, the image of a circle or a line under a quasiconformal mapping of  $\bar{R}^2$ . In Section 3 we establish an extension theorem for quasi-isometries. We then apply this result in Section 4 to show that if  $D$  is a simply connected rigid domain in  $R^2$  and if  $f$  is a local  $L$ -quasi-isometry in  $D$  with  $L < L(D)$ , then  $f$  is not only injective in  $D$  but has an extension as a quasi-isometry to all of  $R^2$ .

## 2. Rigid plane domains

Throughout the remainder of this paper we shall use complex notation to denote points in  $R^2$ . For  $z_0 \in R^2$  and  $0 < r < \infty$  we let  $B(z_0, r)$  denote the open disk with center  $z_0$  and radius  $r$ . Finally for each domain  $D \subset \bar{R}^2$  we let  $D^* = \bar{R}^2 - \bar{D}$ .

In this section we characterize the finitely connected domains in  $R^2$  which are rigid. We begin with a technical lemma concerning a special class of quasi-isometries.

**LEMMA 1.** *Suppose that  $\phi(t)$  is a real valued function defined in  $(0, \infty)$ , that*

$$|\phi(t_1) - \phi(t_2)| \leq a \left| \log \frac{t_1}{t_2} \right| \quad (4)$$

for  $t_1, t_2 \in (0, \infty)$  and that

$$f(z) = \begin{cases} ze^{i\phi(|z|)} & \text{if } 0 < |z| < \infty, \\ 0 & \text{if } z = 0. \end{cases} \quad (5)$$

Then  $f$  is a  $(1+a)$ -quasi-isometry in  $\mathbb{R}^2$ .

*Proof.* Choose distinct points  $z_1, z_2 \in \mathbb{R}^2$  with  $|z_1| \leq |z_2|$ . If  $z_1 \neq 0$ , then

$$\begin{aligned} |f(z_1) - f(z_2)| &\leq |z_1 - z_2| + |z_1| |e^{i\phi(|z_1|)} - e^{i\phi(|z_2|)}| \\ &\leq |z_1 - z_2| + |z_1| |\phi(|z_1|) - \phi(|z_2|)| \\ &\leq |z_1 - z_2| + a |z_1| \left| \log \frac{|z_1|}{|z_2|} \right| \\ &\leq (1+a) |z_1 - z_2| \end{aligned}$$

by (4), while

$$|f(z_1) - f(z_2)| = |z_2| \leq (1+a) |z_1 - z_2|$$

if  $z_1 = 0$ . Since  $f^{-1}$  is given by (5) with  $-\phi$  in place of  $\phi$ , the above argument can be applied to  $f^{-1}$  to complete the proof.

We next use Lemma 1 to obtain a geometric property of plane domains  $D$  with  $L(D) > 1$ .

**LEMMA 2.** *Suppose that  $D$  is a domain in  $\mathbb{R}^2$  with  $L(D) \geq c > 1$ . Then there exists a constant  $b$ , depending only on  $c$ , such that for each  $z_0 \in \mathbb{R}^2$  and  $0 < r < \infty$ ,  $D \cap \partial B(z_0, r)$  lies in component of*

$$G = D \cap (B(z_0, br) - \bar{B}(z_0, r/b)).$$

*Proof.* Choose  $b \in (1, \infty)$  so that

$$1 + \frac{\pi}{\log b} < c, \quad (6)$$

and suppose there exist points  $z_1, z_2 \in D \cap \partial B(z_0, r)$  which belong to different components  $G_1, G_2$  of  $G$ . By making a change of variable we may assume that

$z_0 = 0$ . Choose  $\theta \in [-\pi, \pi]$  so that  $z_2 = z_1 e^{i\theta}$  and let  $f$  be as in (5) with

$$\phi(t) = \begin{cases} 0 & \text{if } 0 < t \leq \frac{r}{b} \text{ or } br \leq t < \infty, \\ \frac{\log \frac{bt}{r}}{\log b} \theta & \text{if } \frac{r}{b} \leq t \leq r, \\ \frac{\log \frac{br}{t}}{\log b} \theta & \text{if } r \leq t \leq br. \end{cases}$$

Then  $\phi$  satisfies (4) with  $a = \pi/\log b$  and  $f$  is a  $(1+a)$ -quasi-isometry in  $\mathbb{R}^2$  by Lemma 1. Set

$$g(z) = \begin{cases} z & \text{if } z \in D - G_1, \\ f(z) & \text{if } z \in G_1. \end{cases} \quad (7)$$

If  $U$  is any open disk in  $D$ , then either  $U \subset D - G_1$ , in which case  $g(z) = z$  in  $U$ , or  $U \subset G_1 \cup (D - G)$ , in which case  $g(z) = f(z)$  in  $U$ . Hence  $g$  is a local  $(1+a)$ -quasi-isometry in  $D$ . Since  $z_2 \notin G_1$ ,

$$g(z_2) = z_2 = z_1 e^{i\theta} = z_1 e^{i\phi(|z_1|)} = g(z_1)$$

and  $g$  is not injective in  $D$ . Thus  $c \leq 1+a$ . This contradicts (6) and establishes the desired conclusion.

We say that  $C \subset \bar{\mathbb{R}}^2$  is a  $K$ -quasicircle if it is the image of a circle or line under a  $K$ -quasiconformal mapping  $f: \bar{\mathbb{R}}^2 \rightarrow \bar{\mathbb{R}}^2$ . Similarly  $D \subset \bar{\mathbb{R}}^2$  is said to be a  $K$ -quasidisk if  $\partial D$  is a  $K$ -quasicircle.

We have next the following information about the boundary of a rigid plane domain.

**LEMMA 3.** *Suppose that  $D$  is a domain in  $\mathbb{R}^2$  with  $L(D) \geq c > 1$ . Then each component  $C$  of  $\partial D$  is either a point or a  $K$ -quasicircle where  $K$  depends only on  $c$ . Moreover if  $C_1$  and  $C_2$  are components of  $\partial D$ , then*

$$\min_{i=1,2} \text{dia}(C_i) \leq a \text{d}(C_1, C_2) \quad (8)$$

where  $a$  is a constant which depends only on  $c$ .

Here  $\text{dia}(C_j)$  denotes the diameter of  $C_j$  and  $d(C_1, C_2)$  the distance between  $C_1$  and  $C_2$ .

*Proof.* Choose  $b \in (1, \infty)$  so that (6) holds and suppose that  $z_0 \in \mathbb{R}^2$ ,  $0 < r < \infty$  and  $z_1, z_2 \in D \cap \bar{B}(z_0, r)$ . Let  $\alpha$  be any arc joining  $z_1$  and  $z_2$  in  $D$ . If  $\alpha$  does not lie in  $\bar{B}(z_0, r)$ , then  $\alpha \cap \bar{B}(z_0, r)$  contains two components  $\alpha_1, \alpha_2$  which join  $z_1, z_2$  to  $w_1, w_2 \in \partial B(z_0, r)$ , respectively. Lemma 2 implies that  $w_1$  and  $w_2$  can be joined by an arc  $\beta$  in  $D \cap \bar{B}(z_0, br)$  and hence  $\alpha_1 \cup \beta \cup \alpha_2$  joins  $z_1$  and  $z_2$  in  $D \cap \bar{B}(z_0, br)$ . A similar argument shows that any pair of points  $z_1, z_2 \in D - B(z_0, r)$  can be joined in  $D - B(z_0, r/b)$ . Hence  $D$  is  $b$ -locally connected and by Lemma 5 in [3], each component  $C$  of  $\partial D$  is either a point or a  $K$ -quasicircle where  $K$  depends only on  $b$ .

Suppose next that  $C_1$  and  $C_2$  are distinct components of  $\partial D$ , choose  $z_1 \in C_1$  and  $z_2 \in C_2$  so that

$$|z_1 - z_2| = d(C_1, C_2) = 2r$$

and let  $z_0 = \frac{1}{2}(z_1 + z_2)$ . We shall use Lemma 2 to show that  $C_1$  or  $C_2$  lies in  $B(z_0, b^2r)$  and hence that

$$\min_{i=1,2} \text{dia}(C_i) \leq 2b^2r.$$

This will establish (8) with  $a = b^2$ .

Suppose that  $C_1$  and  $C_2$  do not lie in  $B(z_0, b^2r)$  and let  $D_0$  denote the component of  $\bar{\mathbb{R}}^2 - (C_1 \cup C_2)$  which contains  $D$ . Then

$$F_1 = \bar{\mathbb{R}}^2 - (D_0 \cap B(z_0, b^2r)), \quad F_2 = \bar{B}(z_0, r)$$

are continua with

$$F_1 \cap F_2 = (C_1 \cup C_2) \cap \bar{B}(z_0, r) = \{z_1, z_2\}. \quad (9)$$

Hence by Theorem V.11.5 in [15], there exist points  $w_1, w_2$  which lie in different components  $G_1, G_2$  of

$$\bar{\mathbb{R}}^2 - (F_1 \cup F_2) = D_0 \cap (B(z_0, b^2r) - \bar{B}(z_0, r))$$

but which can be joined by an arc  $\alpha$  in

$$\bar{\mathbb{R}}^2 - F_1 = D_0 \cap B(z_0, b^2r).$$

Next (9) and Theorem V.16.2 in [15] imply that  $w_1, w_2$  are not separated by  $C_1 \cup C_2 \cup \bar{B}(z_0, r)$  and hence can be joined by an arc  $\beta$  in  $D_0 - \bar{B}(z_0, r)$ . Thus for  $j = 1, 2$ ,  $\alpha \cup \beta$  contains a curve which joins  $\partial B(z_0, r)$  to  $\partial B(z_0, b^2 r)$  in  $G_j$ ; hence

$$H_j = G_j \cap \partial B(z_0, br) \neq \emptyset.$$

Since each component of  $H_j$  is an open arc in  $D_0$  with endpoints in  $C_1 \cup C_2$ ,  $D \cap H_j \neq \emptyset$  and we conclude that  $D \cap \partial B(z_0, br)$  does not lie in a component of  $D \cap (B(z_0, b^2 r) - \bar{B}(z_0, r))$ . This contradicts Lemma 2 and thus establishes the desired conclusion.

Finally we have the following relations between quasidisks and rigid plane domains.

**THEOREM 3.** *If  $D$  is a  $K$ -quasidisk in  $R^2$ , then  $L(D) \geq c > 1$  where  $c$  depends only on  $K$ . Conversely if  $D$  is a simply connected proper subdomain of  $R^2$  with  $L(D) \geq c > 1$ , then  $D$  is a  $K$ -quasidisk where  $K$  depends only on  $c$ .*

*Proof.* If  $D$  is a  $K$ -quasidisk in  $R^2$ , then by Corollary 2.33 in [14],  $D$  is a uniform domain where the constants  $a$  and  $b$  in (2) and (3) depend only on  $K$ . (For an alternative proof see Theorem III.2.3 in [4].) Hence  $L(D) \geq c > 1$  where  $c = c(K)$  by Theorem 2. The converse is a consequence of Lemma 3.

**THEOREM 4.** *A finitely connected domain  $D$  in  $R^2$  is rigid if and only if each component of  $\partial D$  is either a point or a quasicircle.*

*Proof.* If  $D$  is bounded by a finite number of points or quasicircles, then  $D$  is uniform by Theorem 5 in [16] and Theorem 5 in [6]; hence  $D$  is rigid by Theorem 2. The converse follows from Lemma 3.

The problem of characterizing rigid plane domains  $D$  is more difficult when  $D$  is infinitely connected. For example, if  $\mathcal{C}$  denotes the collection of boundary components of a rigid domain  $D$  in  $R^2$ , then

$$\sup_{C, C' \in \mathcal{C}} \frac{\min(\text{dia}(C), \text{dia}(C'))}{d(C, C')} < \infty$$

by Lemma 3. Hence one must take into account not only the shape but the relative size and position of the boundary components when  $D$  has infinite connectivity.

We conclude this section by exhibiting a plane domain  $D$  which is rigid but not uniform; thus the converse of Theorem 2 does not hold. The existence of such a domain is an immediate consequence of the following result.

**THEOREM 5.** *If  $D$  is a rigid domain in  $R^2$  and if  $E$  is a discrete subset of  $D$ , then  $D - E$  is a rigid domain.*

*Proof.* Suppose that  $U$  is an open disk with center at  $z_0$  and let  $U_0 = U - \{z_0\}$ . Then since  $L(D)$  is invariant under similarity mappings, Theorem 4 implies that  $L(U_0)$  is an absolute constant  $c$  which exceeds 1.

Suppose next that  $f$  is a local  $L$ -quasi-isometry in  $D - E$  with  $L < \min(L(D), c)$ . Given  $z_0 \in E$  we can choose an open disk  $U$  centered at  $z_0$  such that

$$U_0 = U - \{z_0\} \subset D - E.$$

If  $z_1, z_2 \in U_0$ , then for each  $\varepsilon > 0$  we can find an arc  $\alpha$  joining  $z_1$  and  $z_2$  in  $U_0$  with

$$l(\alpha) \leq (1 + \varepsilon) |z_1 - z_2|.$$

Since  $f$  is a local  $L$ -quasi-isometry in  $U_0$ ,

$$|f(z_1) - f(z_2)| \leq l(f(\alpha)) \leq Ll(\alpha) \leq L(1 + \varepsilon) |z_1 - z_2|,$$

and letting  $\varepsilon \rightarrow 0$  yields

$$|f(z_1) - f(z_2)| \leq L |z_1 - z_2|. \quad (10)$$

Then (10) implies that  $f$  has a continuous extension in  $U$  which satisfies (10) for  $z_1, z_2 \in U$ . Next since  $L < c$ ,  $f$  is injective in  $U_0$ , and it follows that  $f$  is injective and hence a homeomorphism in  $U$ . Choose an open disk  $V$  about  $f(z_0)$  with  $V \subset f(U)$ . Then  $g = (f|U)^{-1}$  is a local  $L$ -quasi-isometry in  $V_0 = V - \{f(z_0)\}$ , and the above argument applied to  $g$  shows that  $f$  is an  $L$ -quasi-isometry in  $g(V)$ . Thus  $f$  has an extension to  $D$  which is a local  $L$ -quasi-isometry in a neighborhood of each point of  $E$  and hence in  $D$ . Then since  $L < L(D)$ ,  $f$  is injective in  $D$ . Hence  $f$  is injective in  $D - E$ ,

$$L(D - E) \geq \min(L(D), c) > 1$$

and  $D - E$  is a rigid domain.

Now let  $B$  denote the unit disk and let

$$E = \left\{ z = \left(1 - \frac{1}{j}\right) \exp\left(\frac{2\pi i k}{j^2}\right) : k = 1, 2, \dots, j^2, j = 2, 3, \dots \right\}.$$

Then  $D = B - E$  is rigid by Theorem 5. On the other hand if  $z_1 = 0$  and if  $z_2 \in D$  with  $|z_2| \geq 1 - 1/2j$  and  $j \geq 2$ , then each rectifiable arc  $\alpha$  joining  $z_1$  and  $z_2$  in  $D$  must contain a point  $z$  with  $|z| = 1 - 1/j$  and

$$\min_{j=1,2} l(\alpha_j) \geq \frac{j}{2\pi} d(z, \partial D),$$

where  $\alpha_1, \alpha_2$  denote the components of  $\alpha - \{z\}$ . Hence there exists no constant  $b$  for which  $D$  satisfies condition (3) and  $D$  is not a uniform domain.

### 3. Extension of quasi-isometries

We establish here some extension theorems for plane quasi-isometries. Our arguments are based on a reflection principle for quasidisks due to Ahlfors [1] and estimates for the hyperbolic distance.

If  $D$  is a simply connected proper subdomain of  $\mathbb{R}^2$ , then the *hyperbolic metric* with curvature  $-1$  in  $D$  is given by

$$\rho_D(z) = \frac{|g'(z)|}{\operatorname{Im}(g(z))},$$

where  $g$  is any conformal mapping of  $D$  onto the upper half plane  $H$ . From standard distortion theorems it follows that

$$\frac{1}{2} \leq |g'(z)| \frac{d(z, \partial D)}{\operatorname{Im}(g(z))} \leq 2, \quad (11)$$

where  $d(z, \partial D)$  denotes the distance from  $z$  to  $\partial D$ , and hence that

$$\frac{1}{2d(z, \partial D)} \leq \rho_D(z) \leq \frac{2}{d(z, \partial D)}. \quad (12)$$

(See, for example, p. 22 in [17].) Next the *hyperbolic distance* between points  $z_1, z_2 \in D$  is given by

$$h_D(z_1, z_2) = \inf_{\alpha} \int_{\alpha} \rho_D(z) |dz|,$$

where the infimum is taken over all rectifiable arcs  $\alpha$  joining  $z_1$  and  $z_2$  in  $D$ . From

(12) and Lemma 2.1 in [5] it follows that

$$h_D(z_1, z_2) \geq \frac{1}{2} \left| \log \frac{d(z_1, \partial D)}{d(z_2, \partial D)} \right| \quad (13)$$

for  $z_1, z_2 \in D$ . Next if  $D$  is a  $K$ -quasidisk, then by (12), Corollary 2.33 in [14] and Theorem 1 in [6],

$$h_D(z_1, z_2) \leq c \log \left( \frac{|z_1 - z_2|}{d(z_1, \partial D)} + 1 \right) \left( \frac{|z_1 - z_2|}{d(z_2, \partial D)} + 1 \right) + d. \quad (14)$$

for  $z_1, z_2 \in D$ , where  $c$  and  $d$  are constants which depend only on  $K$ . (Cf. pp. 42–44 in [13].)

We begin with a result on a special class of quasi-isometries.

LEMMA 4. *If  $D$  is a Jordan domain in  $R^2$  and if  $z_1, z_2 \in D$  with  $h_D(z_1, z_2) \leq a$ , then there exists an  $L$ -quasi-isometry  $f: \bar{D} \rightarrow \bar{D}$  such that  $f$  is the identity on  $\partial D$ ,  $f(z_1) = z_2$  and  $L$  depends only on  $a$ .*

*Proof.* Choose a conformal mapping  $g: D \rightarrow H$  normalized so that  $g(z_1) = i$  and  $g(z_2) = bi$  where  $b > 1$ . Then

$$\log b = h_D(z_1, z_2) \leq a$$

and  $g$  extends to a homeomorphism which maps  $\bar{D}$  onto  $\bar{H}$ . Set

$$h(w) = \begin{cases} u + iv & \text{if } w = u + iv \in \bar{H} - \{\infty\}, \\ \infty & \text{if } w = \infty. \end{cases}$$

Then  $h$  is continuously differentiable with

$$\left. \begin{aligned} h_H(h(w), w) &= \log b \leq a, \\ \frac{1}{b} \frac{|dw|}{\operatorname{Im}(w)} &\leq \frac{|dh(w)|}{\operatorname{Im}(h(w))} \leq \frac{|dw|}{\operatorname{Im}(w)} \end{aligned} \right\} \quad (15)$$

in  $H$  and  $f = g^{-1} \circ h \circ g$  is a homeomorphism of  $\bar{D}$  onto  $\bar{D}$  which is the identity on  $\partial D$  and maps  $z_1$  onto  $z_2$ .

Fix  $z \in D$  and set  $w = g(z)$ . Then

$$\frac{|df(z)|}{|dz|} = \frac{|dh(w)|}{|dw|} \frac{|g'(z)|}{|g'(f(z))|}, \quad (16)$$

while we obtain

$$\left. \begin{aligned} \frac{1}{2} \leq |g'(z)| \frac{d(z, \partial D)}{\operatorname{Im}(w)} \leq 2 \\ \frac{1}{2} \leq |g'(f(z))| \frac{d(f(z), \partial D)}{\operatorname{Im}(h(w))} \leq 2 \end{aligned} \right\} \quad (17)$$

from (11). Next by (13) and (15),

$$\frac{1}{2} \left| \log \frac{d(f(z), \partial D)}{d(z, \partial D)} \right| \leq h_D(f(z), z) = h_H(h(w), w) \leq a$$

whence

$$e^{-2a} \leq \frac{d(f(z), \partial D)}{d(z, \partial D)} \leq e^{2a}. \quad (18)$$

Combining (15), (16), (17) and (18) yields

$$\frac{1}{L} \leq \frac{|df(z)|}{|dz|} \leq L$$

where  $L = 4e^{3a}$ , and hence  $f$  is a local  $L$ -quasi-isometry in  $D$ . (Cf. p. 395 in [10].) The desired conclusion is now a consequence of the following elementary result.

**LEMMA 5.** *Suppose that  $D_1$  and  $D_2$  are domains in  $R^2$ , that  $f: \bar{D}_1 \rightarrow \bar{D}_2$  is a homeomorphism and that  $f$  is an  $L_1$ -quasi-isometry in  $\partial D_1$  and a local  $L_2$ -quasi-isometry in  $D_1$ . Then  $f$  is an  $L$ -quasi-isometry in  $\bar{D}_1$  where  $L = \max(L_1, L_2)$ .*

*Proof.* Fix  $z_1, z_2 \in D_1$  and let  $\alpha$  be the open segment joining these points in  $R^2$ . If  $\alpha \subset D_1$ , then

$$|f(z_1) - f(z_2)| \leq l(f(\alpha)) \leq L_2 l(\alpha) \leq L |z_1 - z_2|.$$

Otherwise for  $j = 1, 2$  let  $\alpha_j$  denote the component of  $\alpha \cap D_1$  which has  $z_j$  as an endpoint and let  $w_j$  denote the other endpoint of  $\alpha_j$ . Then  $w_j \in \partial D_1$ ,  $\alpha_j \subset D_1$  and

$$\begin{aligned} |f(z_1) - f(z_2)| &\leq |f(z_1) - f(w_1)| + |f(w_1) - f(w_2)| + |f(w_2) - f(z_2)| \\ &\leq L |z_1 - w_1| + L_1 |w_1 - w_2| + L |w_2 - z_2| \\ &\leq L |z_1 - z_2|. \end{aligned}$$

Applying this argument to  $f^{-1}$  shows that  $f$  is an  $L$ -quasi-isometry in  $D_1$ , and hence in  $\bar{D}_1$  by continuity.

Lemma 5 shows that a bijective local quasi-isometry between two domains is a quasi-isometry if the induced boundary correspondence is a quasi-isometry. We can also draw this conclusion without knowledge of the boundary correspondence when the two domains have sufficiently regular boundaries.

**LEMMA 6.** *Suppose that  $D_1$  and  $D_2$  are  $K_1$ - and  $K_2$ -quasidisks in  $R^2$  and that  $f: D_1 \rightarrow D_2$  is a bijective local  $L_1$ -quasi-isometry. Then  $f$  extends to an  $L$ -quasi-isometry of  $\bar{D}_1$  onto  $\bar{D}_2$  where  $L$  depends only on  $K_1$ ,  $K_2$  and  $L_1$ .*

*Proof.* Fix  $z_1, z_2 \in D_1$ . By Corollary 2.33 in [14], there exists a rectifiable arc  $\alpha$  joining  $z_1$  and  $z_2$  in  $D_1$  such that

$$l(\alpha) \leq a_1 |z_1 - z_2|,$$

where  $a_1$  depends only on  $K_1$ . Thus

$$|f(z_1) - f(z_2)| \leq l(f(\alpha)) \leq L_1 l(\alpha) \leq L_1 a_1 |z_1 - z_2|.$$

Next since  $f$  is injective,  $f^{-1}$  is a local  $L_1$ -quasi-isometry in  $D_2$  and arguing as above yields

$$|z_1 - z_2| \leq L_1 a_2 |f(z_1) - f(z_2)|,$$

where  $a_2$  depends only on  $K_2$ . Hence  $f$  is an  $L$ -quasi-isometry in  $D_1$  where  $L = \max(L_1 a_1, L_1 a_2)$ , and we can extend  $f$  to  $\bar{D}_1$  by continuity.

We will require the following version of Lemma 4 for the case where  $D$  is a quasidisk.

**LEMMA 7.** *Suppose that  $D$  is a  $K$ -quasidisk in  $R^2$ , that  $z_1, z_2 \in D$  and that*

$$\frac{1}{b} \leq \frac{|z_1 - z|}{|z_2 - z|} \leq b \tag{19}$$

for all  $z \in \partial D - \{\infty\}$  where  $b$  is a constant. Then there exists an  $L$ -quasi-isometry  $f: \bar{D} \rightarrow \bar{D}$  such that  $f$  is the identity on  $\partial D$ ,  $f(z_1) = z_2$  and  $L$  depends only on  $K$  and  $b$ .

*Proof.* For  $j = 1, 2$  choose  $w_j \in \partial D - \{\infty\}$  so that

$$|z_j - w_j| = d(z_j, \partial D).$$

Then by (19),

$$|z_1 - z_2| \leq |z_1 - w_j| + |z_2 - w_j| \leq (b + 1) d(z_j, \partial D)$$

and hence

$$h_D(z_1, z_2) \leq 2c \log(b + 2) + d = a$$

by (14), where  $c$  and  $d$  depend only on  $K$ . The desired conclusion now follows directly from Lemma 4.

We derive now an extension of Ahlfors' reflection principle for quasidisks. (See, for example, Lemma 3 on p. 80 in [2].)

**THEOREM 6.** Suppose that  $D_1$  is a  $K_1$ -quasidisk with  $\infty \in \partial D_1$ , that  $D_2$  is a Jordan domain in  $\mathbb{R}^2$  with  $\infty \in \partial D_2$  and that  $\phi: \partial D_1 \rightarrow \partial D_2$  is an  $L_1$ -quasi-isometry. Then there exists an  $L$ -quasi-isometry  $f: \bar{D}_1 \rightarrow \bar{D}_2$  such that  $f = \phi$  on  $\partial D_1$  and  $L$  depends only on  $K_1$  and  $L_1$ . Suppose further that  $z_1 \in D_1$ ,  $z_2 \in D_2$  and

$$\frac{1}{b} \leq \frac{|z_1 - z|}{|z_2 - \phi(z)|} \leq b \quad (20)$$

for all  $z \in \partial D_1 - \{\infty\}$  where  $b$  is a constant. Then we can choose  $f$  so that, in addition,  $f(z_1) = z_2$  and  $L$  depends only on  $K_1$ ,  $L_1$  and  $b$ .

If we choose  $D_2 = D_1^*$  and  $\phi(z) = z$ , then the first part of Theorem 6 yields the above mentioned result of Ahlfors.

*Proof.* For  $j = 1, 2$  let  $g_j$  map  $D_j$  conformally onto the upper half plane  $H$ . Then  $g_j$  extends to a homeomorphism of  $\bar{D}_j$  onto  $\bar{H}$  and by performing an additional Möbius transformation we may assume that  $g_j(\infty) = \infty$ . Hence  $\psi(x) = g_2 \circ \phi \circ g_1^{-1}(x)$  is a homeomorphism of  $\partial H$  onto itself with  $\psi(\infty) = \infty$ .

Choose  $-\infty < x < \infty$  and  $t > 0$ , let  $\alpha'_1 = (x, x + t)$  and  $\beta'_1 = (-\infty, x - t)$ , and let  $\alpha_1, \alpha_2, \alpha'_2$  and  $\beta_1, \beta_2, \beta'_2$  denote the images of  $\alpha'_1$  and  $\beta'_1$  under  $g_1^{-1}, \phi \circ g_1^{-1}, \psi$  respectively. If  $\Gamma_1$  is the family of arcs joining  $\alpha_1$  to  $\beta_1$  in  $D_1$ , then the extremal

length  $\lambda(\Gamma_1)$  of  $\Gamma_1$  is equal to 1. Moreover since  $D_1$  is a  $K_1$ -quasidisk,

$$|z_1 - z_2| \leq c_1 |z_1 - z_3| \quad (21)$$

for each ordered triple of points  $z_1, z_2, z_3 \in \partial D_1 - \{\infty\}$  where  $c_1$  is a constant which depends only on  $K_1$ . In particular if we let  $z_1 = g_1^{-1}(x)$  and  $w_1 = g_1^{-1}(x+t)$ , then the argument on pp. 82–83 in [2] shows that

$$\alpha_1 \subset \bar{B}(w_1, r), \quad r = c_1 |z_1 - w_1|$$

and that

$$d(\alpha_1, \beta_1) \geq s = c_1^{-5} e^{-2\pi} r.$$

Since  $\phi$  is an  $L_1$ -quasi-isometry,

$$\alpha_2 \subset \bar{B}(w_2, L_1 r), \quad d(\alpha_2, \beta_2) \geq \frac{s}{L_1}$$

where  $w_2 = \phi(w_1)$ , and arguing again as on p. 83 in [2] we see that

$$\lambda(\Gamma_2) \geq \frac{1}{\pi} \left( \frac{s}{L_1^2 r + s} \right)^2,$$

where  $\Gamma_2$  is the family of arcs joining  $\alpha_2$  to  $\beta_2$  in  $D_2$ . This implies that

$$\frac{\psi(x+t) - \psi(x)}{\psi(x) - \psi(x-t)} \leq c_2, \quad (22)$$

where  $c_2$  is a constant which depends only on  $K_1$  and  $L_1$ . From (22) and the above argument with  $\alpha'_1 = (x-t, x)$  and  $\beta'_1 = (x+t, \infty)$  we conclude that

$$\frac{1}{c_2} \leq \frac{\psi(x+t) - \psi(x)}{\psi(x) - \psi(x-t)} \leq c_2$$

for all such  $x$  and  $t$ . Set

$$h(z) = \frac{1}{2} \int_0^1 (\psi(x+ty) + \psi(x-ty)) dt + \frac{i}{2} \int_0^1 |\psi(x+ty) - \psi(x-ty)| dt$$

for  $z = x + iy \in \bar{H} - \{\infty\}$  and  $h(\infty) = \infty$ . Then  $h$  maps  $\bar{H}$  homeomorphically onto  $\bar{H}$  and  $h$  is continuously differentiable and  $K$ -quasiconformal in  $H$  with

$$\frac{1}{c_3} \frac{|dz|}{\operatorname{Im}(z)} \leq \frac{|dh(z)|}{\operatorname{Im}(h(z))} \leq c_3 \frac{|dz|}{\operatorname{Im}(z)},$$

where  $K$  and  $c_3$  depend only on  $c_2$ , and hence on  $K_1$  and  $L_1$ . (See pp. 69–74 in [2] for the case where  $\psi(x)$  is increasing in  $x$ .) Thus  $f_1 = g_2^{-1} \circ h \circ g_1$  is a homeomorphism of  $\bar{D}_1$  onto  $\bar{D}_2$ ,  $f = \phi$  on  $\partial D_1$ ,  $f_1$  is  $K$ -quasiconformal in  $D_1$  and

$$\frac{|df_1(z)|}{|dz|} = \frac{|dh(w)|}{|dw|} \frac{|g'_1(z)|}{|g'_2(f_1(z))|}$$

for  $z \in D_1$ ,  $w = g_1(z)$ . From (11) applied to  $g_1$  and  $g_2$  we obtain

$$\frac{1}{2} \leq |g'_1(z)| \frac{d(z, \partial D_1)}{\operatorname{Im}(w)} \leq 2,$$

$$\frac{1}{2} \leq |g'_2(f_1(z))| \frac{d(f_1(z), \partial D_2)}{\operatorname{Im}(h(w))} \leq 2.$$

Thus

$$\frac{1}{4c_3} \frac{d(f_1(z), \partial D_2)}{d(z, \partial D_1)} \leq \frac{|df_1(z)|}{|dz|} \leq 4c_3 \frac{d(f_1(z), \partial D_2)}{d(z, \partial D_1)} \quad (23)$$

and it remains to bound the ratio on the left and right sides of (23).

If  $w_1, w_2, w_3$  is an ordered triple of points in  $\partial D_2 - \{\infty\}$ , then

$$|w_1 - w_2| \leq c_1 L_1^2 |w_1 - w_3|$$

by (21) and  $D_2$  is a  $K_2$ -quasidisk where  $K_2$  depends only on  $K_1$  and  $L_1$ . Hence  $f_1$  can be extended by quasiconformal reflection in  $\partial D_1$  and  $\partial D_2$  to yield a  $K_3$ -quasiconformal mapping of  $\bar{R}^2$  onto itself with  $K_3 = KK_1^2K_2^2$ . Fix  $z_1 \in D_1$  and  $z_2 \in \partial D_1 - \{\infty\}$ , and choose  $z_3 \in \partial D_1$  so that  $|z_3 - z_2| = |z_1 - z_2|$ . Since  $f_1$  is  $K_3$ -quasiconformal in  $\bar{R}^2$  with  $f_1(\infty) = \infty$ ,

$$|f_1(z_1) - f_1(z_2)| \leq c |f_1(z_3) - f_1(z_2)| \leq c_4 |z_3 - z_2| = c_4 |z_1 - z_2|$$

where  $c$  and  $c_4 = cL_1$  depend only on  $K_1$  and  $L_1$ . We thus obtain

$$\frac{1}{c_4} |z_1 - z_2| \leq |f_1(z_1) - f_1(z_2)| \leq c_4 |z_1 - z_2| \quad (24)$$

for all  $z_1 \in D_1$  and  $z_2 \in \partial D_1 - \{\infty\}$ . In particular, (24) implies that

$$\frac{1}{c_4} d(z, \partial D_1) \leq d(f_1(z), \partial D_2) \leq c_4 d(z, \partial D_1)$$

for all  $z \in D_1$ , and we conclude from (23) that  $f_1$  is a local  $L_2$ -quasi-isometry in  $D_1$  with  $L_2 = 4c_3c_4$ . Lemma 5 then implies that  $f_1$  is an  $L_3$ -quasi-isometry in  $\bar{D}_1$  where  $L_3 = \max(L_1, L_2)$ , and choosing  $f = f_1$  completes the proof of the first part of Theorem 6.

Finally suppose that  $z_1 \in D_1$ ,  $z_2 \in D_2$  and that (20) holds for all  $z \in \partial D_1 - \{\infty\}$ . If  $w \in \partial D_2 - \{\infty\}$ , then  $z = f_1^{-1}(w) \in \partial D_1 - \{\infty\}$  and

$$\frac{|f_1(z_1) - w|}{|z_2 - w|} = \frac{|f_1(z_1) - f_1(z)|}{|z_1 - z|} \frac{|z_1 - z|}{|z_2 - \phi(z)|}$$

lies between  $(bc_4)^{-1}$  and  $bc_4$  by (20) and (24). By Lemma 7 there exists an  $L_4$ -quasi-isometry  $f_2: \bar{D}_2 \rightarrow \bar{D}_2$  such that  $f_2$  is the identity on  $\partial D_2$ ,  $f_2(f_1(z_1)) = z_2$  and  $L_4$  depends only on  $K_2$  and  $b$ . Thus  $f = f_2 \circ f_1$  has all the properties required in the second part of Theorem 6.

Finally we require the following result which shows that a certain class of quasi-isometries is invariant under conjugation by inversion.

**LEMMA 8.** *Suppose that  $f$  is an  $L$ -quasi-isometry in  $E \subset \bar{R}^2$ , that*

$$\frac{1}{L} \leq \frac{|f(z)|}{|z|} \leq L \quad (25)$$

*for  $z \in E - \{0, \infty\}$  and that  $f(0) = 0$  if  $0 \in E$ . Then  $g = T \circ f \circ T^{-1}$  is an  $L^3$ -quasi-isometry in  $T(E)$  where  $T(z) = 1/z$ .*

*Proof.* Choose distinct points  $w_1, w_2 \in T(E) - \{\infty\}$  and let  $z_i = 1/w_i$ . If  $w_1, w_2 \neq 0$ , then  $z_1, z_2 \in E - \{0, \infty\}$ ,

$$\frac{|g(w_1) - g(w_2)|}{|w_1 - w_2|} = \frac{|f(z_1) - f(z_2)|}{|z_1 - z_2|} \frac{|z_1|}{|f(z_1)|} \frac{|z_2|}{|f(z_2)|}$$

and hence

$$\frac{1}{L^3} \leq \frac{|g(w_1) - g(w_2)|}{|w_1 - w_2|} \leq L^3 \quad (26)$$

by (25). If  $w_1 = 0$ , then  $g(w_1) = 0$ ,

$$\frac{|g(w_1) - g(w_2)|}{|w_1 - w_2|} = \frac{|z_2|}{|f(z_2)|}$$

and again (26) holds. Finally if  $\infty \in T(E)$  then  $g(\infty) = \infty$  and thus  $g$  is an  $L^3$ -quasi-isometry in  $T(E)$ .

We now obtain the main result of this section from combining Lemma 8 and Theorem 6.

**THEOREM 7.** *Suppose that  $D_1$  is a  $K_1$ -quasidisk in  $\mathbb{R}^2$ , that  $D_2$  is a Jordan domain in  $\mathbb{R}^2$  and that  $\phi: \partial D_1 \rightarrow \partial D_2$  is an  $L_1$ -quasi-isometry. Then there exist  $L$ -quasi-isometries  $f: \bar{D}_1 \rightarrow \bar{D}_2$  and  $f^*: \bar{D}_1^* \rightarrow \bar{D}_2^*$  such that  $f = f^* = \phi$  on  $\partial D_1$  and  $L$  depends only on  $K_1$  and  $L_1$ .*

*Proof.* Suppose that  $\infty \in \partial D_1$ . Then  $D_1$  and  $D_1^*$  are  $K_1$ -quasidisks with  $\partial D_1 = \partial D_1^*$  and the existence of  $f$  and  $f^*$  is an immediate consequence of Theorem 6.

Suppose next that  $\infty \notin \partial D_1$ . Then  $\infty \in D_1^*$  and  $\infty \in D_2^*$ . By making a preliminary change of variables we may assume that  $0 \in \partial D_1$  and that  $\phi(0) = 0$ . For  $j = 1, 2$  let  $G_j$  denote the image of  $D_j$  under  $T(z) = 1/z$  and set  $\psi = T \circ \phi \circ T^{-1}$ . Then  $G_1$  is a  $K_1$ -quasidisk with  $\infty \in \partial G_1$ ,

$$\frac{1}{L_1} \leq \frac{|\phi(z)|}{|z|} \leq L_1 \quad (27)$$

for  $z \in \partial D_1 - \{0\}$  and hence  $\psi: \partial G_1 \rightarrow \partial G_2$  is an  $L_1^3$ -quasi-isometry by Lemma 8. Theorem 6 then yields  $L_2$ -quasi-isometries  $g: \bar{G}_1 \rightarrow \bar{G}_2$  and  $g^*: \bar{G}_1^* \rightarrow \bar{G}_2^*$  such that  $g = g^* = \psi$  on  $\partial G_1$  and  $L_2$  depends only on  $K_1$  and  $L_1$ . In addition, since  $0 \in G_1^*$ ,  $0 \in G_2^*$  and

$$\frac{1}{L_1} \leq \frac{|0 - z|}{|0 - \psi(z)|} \leq L_1$$

for  $z \in \partial G_1 - \{\infty\}$  by (27), we can choose  $g^*$  so that  $g^*(0) = 0$ . Fix  $z_1 \in \bar{G}_1 - \{\infty\}$  and

let  $z_0$  be a point where the segment joining 0 to  $z_1$  meets  $\partial G_1$ . Then

$$\begin{aligned} |g(z_1)| &\leq |g(z_1) - g(z_0)| + |g(z_0)| \leq L_2 |z_1 - z_0| + |\psi(z_0)| \\ &\leq L_2 |z_1 - z_0| + L_1 |z_0| \leq L_2 |z_1| \end{aligned}$$

since  $L_2 \geq L_1$ . Thus by symmetry

$$\frac{1}{L_2} \leq \frac{|g(z)|}{|z|} \leq L_2$$

for  $z \in \bar{G}_1 - \{\infty\}$ . Next

$$\frac{1}{L_2} \leq \frac{|g^*(z)|}{|z|} \leq L_2$$

for  $z \in \bar{G}_1^* - \{0, \infty\}$  since  $g^*$  is an  $L_2$ -quasi-isometry in  $\bar{G}_1^*$ . Thus  $f = T^{-1} \circ g \circ T$  and  $f^* = T^{-1} \circ g^* \circ T$  have the required properties by Lemma 8.

**COROLLARY 1.** *Suppose that  $D_1$  and  $D_2$  are  $K_1$ - and  $K_2$ -quasidisks in  $\mathbb{R}^2$  and that  $f: D_1 \rightarrow D_2$  is a bijective local  $L_1$ -quasi-isometry. Then there exists an  $L$ -quasi-isometry  $g: \bar{R}^2 \rightarrow \bar{R}^2$  such that  $g = f$  in  $D_1$  and  $L$  depends only on  $K_1$ ,  $K_2$  and  $L_1$ .*

*Proof.* By Lemma 6,  $f$  extends to an  $L_2$ -quasi-isometry of  $\bar{D}_1$  onto  $\bar{D}_2$  where  $L_2$  depends only on  $K_1$ ,  $K_2$  and  $L_1$ . Next Theorem 7 with  $\phi = f|_{\partial D_1}$  yields an  $L_2^*$ -quasi-isometry  $f^*: \bar{D}_1^* \rightarrow \bar{D}_2^*$  such that  $f^* = f$  on  $\partial D_1$  and  $L_2^*$  depends only on  $K_1$ ,  $K_2$  and  $L_1$ . Then

$$g = \begin{cases} f & \text{in } \bar{D}_1 \\ f^* & \text{in } \bar{D}_1^*, \end{cases} \quad (28)$$

is the desired extension of  $f$ .

**COROLLARY 2.** *Suppose that  $C_1$  is a  $K_1$ -quasicircle and that  $\phi$  is an  $L_1$ -quasi-isometry in  $C_1$ . Then there exists an  $L$ -quasi-isometry  $g: \bar{R}^2 \rightarrow \bar{R}^2$  such that  $g = \phi$  on  $C_1$  and  $L$  depends only on  $K_1$  and  $L_1$ .*

*Proof.* Let  $C_2 = \phi(C_1)$  and for  $j = 1, 2$  let  $D_j$  be a component of  $\bar{R}^2 - C_j$  chosen so that  $D_j \subset \mathbb{R}^2$ . If  $f$  and  $f^*$  are the  $L$ -quasi-isometries given by Theorem 7, then  $g$  defined in (28) is the required extension.

Corollary 2 extends recent results of Jerison and Kenig [9] and of Tukia [18] who consider the cases where  $C_1$  is a line and a circle, respectively.

#### 4. An application

If  $f$  is a local  $L$ -quasi-isometry in a plane domain  $D$  with  $L < L(D)$ , then  $f$  is injective. The following result shows that one can say more whenever  $D$  is simply connected.

**THEOREM 8.** *Suppose that  $D_1$  is a simply connected proper subdomain of  $R^2$  and that  $f$  is a local  $L_1$ -quasi-isometry in  $D_1$  with  $L < L(D_1)$ . Then there exists an  $L$ -quasi-isometry  $g: R^2 \rightarrow R^2$  such that  $g = f$  in  $D_1$  and  $L$  depends only on  $L(D_1)$  and  $L_1$ .*

*Proof.* Let  $D_2 = f(D_1)$  and let  $g$  denote any local  $L_2$ -quasi-isometry in  $D_2$  with  $L_2 < L(D_1)/L_1$ . Then  $g \circ f$  is a local  $L_1 L_2$ -quasi-isometry in  $D_1$ ,  $g \circ f$  is injective in  $D_1$  since  $L_1 L_2 < L(D_1)$  and hence  $g$  is injective in  $D_2$ . Thus

$$L(D_2) \geq \frac{L(D_1)}{L_1} > 1. \quad (29)$$

Since  $f$  is injective in  $D_1$ ,  $f$  is an  $L_1^2$ -quasiconformal mapping of  $D_1$  and hence  $D_2$  is a simply connected proper subdomain of  $R^2$ . Then by Theorem 3 and (29)  $D_1$  and  $D_2$  are  $K_1$ - and  $K_2$ -quasidisks, where  $K_1$  and  $K_2$  depend only on  $L(D_1)$  and  $L(D_1)/L_1$  respectively, and the existence of  $g$  follows from Corollary 1.

Theorem 8 can be interpreted physically if we think of  $D_1$  as a homogeneous elastic body and  $f$  as the distortion of  $D_1$  due to a force field. In this case  $L(D_1)$  measures the maximum permissible strain in  $D_1$  before  $D_1$  buckles. Theorem 8 asserts that if the strain in  $D_1$  is less than  $L(D_1)$ , then the shape of  $D_1$  is not substantially changed under the force field.

#### REFERENCES

- [1] L. V. AHLFORS, *Quasiconformal reflections*, Acta Math. 109 (1963) 291–301.
- [2] L. V. AHLFORS, *Lectures on quasiconformal mappings*, Van Nostrand Mathematical Studies 10, Princeton 1966.
- [3] F. W. GEHRING, *Univalent functions and the Schwarzian derivative*, Comment. Math. Helv. 52 (1977) 561–572.
- [4] F. W. GEHRING, *Characteristic properties of quasidisks*, Les Presses de l’Université de Montréal 1982.

- [5] F. W. GEHRING and B. P. PALKA, *Quasiconformally homogeneous domains*, J. d'Analyse Math. 30 (1976) 172–199.
- [6] F. W. GEHRING and B. G. OSGOOD, *Uniform domains and the quasi-hyperbolic metric*, J. d'Analyse Math. 36 (1979) 50–74.
- [7] J. GEVIRTZ, *On the univalence of quasi-isometric mappings*, Rev. Colombiana Mat. 7 (1973) 125–132.
- [8] J. GEVIRTZ, *A sharp condition for univalence in euclidean spaces*, Proc. Amer. Math. Soc. 57 (1976) 261–265.
- [9] D. S. JERISON and C. E. KENIG, *Hardy spaces,  $A_\infty$ , and singular integrals on chord-arc domains*, Math. Scand. (to appear).
- [10] F. JOHN, *Rotation and strain*, Comm. Pure Appl. Math. 14 (1961) 391–413.
- [11] F. JOHN, *On quasi-isometric mappings, I*, Comm. Pure Appl. Math. 21 (1968) 77–110.
- [12] F. JOHN, *On quasi-isometric mappings, II*, Comm. Pure Appl. Math. 22 (1969) 265–278.
- [13] P. W. JONES, *Extension theorems for BMO*, Indiana Univ. Math. J. 29 (1980) 41–66.
- [14] O. MARTIO and J. SARVAS, *Injectivity theorems in plane and space*, Ann. Acad. Sci. Fenn. 4 (1978–79) 383–401.
- [15] M. H. A. NEWMAN, *Elements of the topology of plane sets of points*, Camb. Univ. Press 1954.
- [16] B. G. OSGOOD, *Univalence criteria in multiply-connected domains*, Trans. Amer. Math. Soc. 260 (1980) 459–473.
- [17] C. POMMERENKE, *Univalent functions*, Vandenhoeck and Ruprecht, Göttingen 1975.
- [18] P. TUKIA, *The planar Schönflies theorem for Lipschitz maps*, Ann. Acad. Sci. Fenn. 5 (1980) 49–72.

University of Helsinki  
Helsinki, Finland

University of Michigan  
Ann Arbor, Michigan

Received January 14, 1982