

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 58 (1983)

Artikel: Poincaré duality groups of dimension two, II.
Autor: Eckmann, Beno / Linnell, Peter
DOI: <https://doi.org/10.5169/seals-44592>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Poincaré duality groups of dimension two, II

BENO ECKMANN and PETER LINNELL

1. Introduction

A Poincaré duality group of dimension n , in short a PD^n -group, is a group G acting on \mathbf{Z} such that one has natural isomorphisms

$$H^k(G; A) \cong H_{n-k}(G; \mathbf{Z} \otimes A)$$

for all integers k and all $\mathbf{Z}G$ -modules A (where $\mathbf{Z} \otimes A$ is the tensor product over \mathbf{Z} with diagonal G -action). G is called orientable or not according to whether or not \mathbf{Z} is trivial as a $\mathbf{Z}G$ -module. All “surface groups”, i.e., fundamental groups of closed surfaces of genus ≥ 1 are well-known to be PD^2 -groups. In Eckmann–Müller [4] it was proved that a PD^2 -group with *positive* first Betti number β_1 is isomorphic to a surface group. The purpose of the present paper is to show that the condition on β_1 is automatically fulfilled:

THEOREM 1. *The first Betti number β_1 of a PD^2 -group is positive.*

As a consequence we thus have a complete classification of PD^2 -groups.

THEOREM 2. *A group G is a PD^2 -group if and only if it is isomorphic to a surface group.*

For notations and properties concerning PD^n -groups, not explicitly mentioned here, we refer to [4] where also several (algebraic and topological) consequences are discussed.

2. Finitely generated projective $\mathbf{Z}G$ -modules

For the proof of Theorem 1 we need the following fact, which may be of interest in connection with the conjectures of Bass (4.4 and 4.5 of [2]).

If B is an abelian group, we let $\text{rank } B$ denote the dimension of the \mathbf{Q} -vector space $B \otimes \mathbf{Q}$.

PROPOSITION 3. *Let G be a PD^2 -group, $M \neq 0$ a finitely generated projective $\mathbf{Z}G$ -module, and \mathbf{Z} the trivial $\mathbf{Z}G$ -module. Then $\text{rank}(\mathbf{Z} \otimes_G M) \neq 0$.*

Proof. Let r_M denote the Hattori-Stallings trace of the identity endomorphism of M as defined, e.g., in [1] and [2]. It is a finite linear combination with integral coefficients of the conjugacy classes τ in G ,

$$r_M = \sum_{\tau} r_M(\tau) \tau.$$

For $x \in G$ let $r_M(x)$ be the coefficient of the conjugacy class of x . Suppose that $r_M(x) \neq 0$ for an element $x \in G$, $x \neq 1$. Then there exists, by Proposition 6.2 of [2], a prime p and an integer $n > 0$ such that x is conjugate to x^{p^n} . It follows (see the remark on p. 12 of [2]) that x is contained in a subgroup $H \cong \mathbf{Z}[1/p]$ of G . By Strebel's theorem [5] all subgroups of infinite index in G are of cohomological dimension 1 and thus free. Therefore H has finite index in G ; since G is finitely generated so is H and we have a contradiction. Hence $r_M(x) = 0$ for all $x \in G \setminus 1$ and it follows that $r_M(1) = \text{rank}(\mathbf{Z} \otimes_G M)$.

We now consider the nonzero finitely generated projective $\mathbf{C}G$ -module $M \otimes \mathbf{C}$. We have $r_M(1) = r_{M \otimes \mathbf{C}}(1)$ which is positive by Kaplansky's theorem (see [1], Theorem 8.9), and the result follows.

3. Proof of Theorem 1. Euler characteristic

The completion of the proof is now in the same spirit as [3]. We first note that we can restrict attention to orientable PD^2 -groups. Indeed (see [4], p. 511), if G is non-orientable and G_1 the orientable subgroup of index 2 in G then $\beta_1(G_1) > 0$ implies $\beta_1(G) > 0$.

So let G be an orientable PD^2 -group, and

$$0 \rightarrow P \rightarrow \mathbf{Z}G^d \rightarrow \mathbf{Z}G \xrightarrow{\epsilon} \mathbf{Z} \tag{1}$$

a projective resolution of the trivial $\mathbf{Z}G$ -module \mathbf{Z} . Since PD^n -groups are of type (FP) , the module P is finitely generated projective. Since $H^0(G; \mathbf{Z}G) = H^1(G; \mathbf{Z}G) = 0$ and $H^2(G; \mathbf{Z}G) = \mathbf{Z}$ with trivial G -action for any orientable

PD^2 -group, applying $\text{Hom}_G(-, \mathbf{Z}G)$ to (1) yields an exact sequence

$$\mathbf{Z} \xleftarrow{\gamma} P^* \leftarrow \mathbf{Z}G^d \leftarrow \mathbf{Z}G \leftarrow 0 \quad (2)$$

where $P^* = \text{Hom}_G(P, \mathbf{Z}G)$ is finitely generated projective. Let IG be the kernel of ε (the augmentation ideal) and L the kernel of γ . Applying Schanuel's lemma to (1) and (2) gives

$$P^* \oplus IG \cong \mathbf{Z}G \oplus L.$$

There is a surjection $\mathbf{Z}G^d \twoheadrightarrow L$, and we obtain a surjection $\mathbf{Z}G^{d+1} \twoheadrightarrow P^* \oplus IG$ and hence a surjection $\mathbf{Z}G^{d+1} \twoheadrightarrow P^*$, with kernel $K \neq 0$. Obviously K is a finitely generated projective $\mathbf{Z}G$ -module, and we see from Proposition 3 that $\text{rank}(\mathbf{Z} \otimes_G K) \neq 0$. It follows that $\text{rank}(\mathbf{Z} \otimes_G P^*) \leq d$.

The Euler characteristic $\chi(G)$ of G can be obtained by applying $\mathbf{Z} \otimes_G -$ to the resolution (2) and taking the alternating sum of the ranks:

$$\chi(G) = \text{rank}(\mathbf{Z} \otimes_G P^*) - d + 1 \leq 1.$$

On the other hand $\chi(G) = \beta_0 - \beta_1 + \beta_2 = 2 - \beta_1$ since the Betti numbers β_0 and β_2 of an orientable PD^2 -group are $= 1$. Thus $2 - \beta_1 \leq 1$, i.e., $\beta_1 \geq 1$.

4. Poincaré 2-complexes

As a corollary of the above group-theoretic results the topological application mentioned in [4], Section 2 can be given an improved version.

We recall that a Poincaré n -complex is a CW -complex dominated by a finite complex and fulfilling Poincaré duality of formal dimension n for arbitrary local coefficients. By results of Wall [6] a Poincaré 2-complex X with finite fundamental group $\pi_1(X)$ is homotopy equivalent to the 2-sphere or to the real projective plane; if $\pi_1(X)$ is infinite, then X is aspherical, i.e., an Eilenberg-MacLane complex $K(G, 1)$ for $G = \pi_1(X)$. In the latter case G is a PD^2 -group, and thus by our Theorem 2 isomorphic to $\pi_1(Y)$ where Y is a closed surface of genus ≥ 1 . The isomorphism $\pi_1(X) \cong \pi_1(Y)$ yields a homotopy equivalence between X and Y . In summary we have

THEOREM 4. *A CW -complex is a Poincaré 2-complex if and only if it is homotopy equivalent to a closed surface of genus ≥ 0 .*

REFERENCES

- [1] BASS, H., *Euler characteristics and characters of discrete groups*, *Inventiones Math.* 35 (1976) 155–196.
- [2] BASS, H., *Traces and Euler characteristics*, in: *Homological Group Theory* (Cambridge Univ. Press 1979), 1–26.
- [3] COHEN, J. M., *Poincaré 2-complexes I*, *Topology* 11 (1972) 417–419.
- [4] ECKMANN, B. and MÜLLER, H., *Poincaré duality groups of dimension two*, *Comment. Math. Helvetici* 55 (1980) 510–520.
- [5] STREBEL, R., *A remark on subgroups of infinite index in Poincaré duality groups*, *Comment. Math. Helvetici* 52 (1977) 317–324.
- [6] WALL, C. T. C., *Poincaré complexes I*, *Ann. Math.* 86 (1967) 213–245.

ETH Zurich
8092 Zurich, Switzerland
and
Dept. of Pure Mathematics
and Mathematical Statistics
16 Mill Lane
Cambridge CB2 1SB
England

Received October 4, 1982