

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 58 (1983)

Artikel: Wall's obstructions and Whitehead torsion.
Autor: Kwasik, Slawomir
DOI: <https://doi.org/10.5169/seals-44610>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 24.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Wall's obstructions and Whitehead torsion

SŁAWOMIR KWASIK

In this note we show that the Wall-type obstruction defined by S. Ferry in [4] is in fact the original Wall's one. As a consequence we obtain the geometric proof of the Product Formula (see [5]) for the Wall finiteness obstructions.

1. Introduction

Let X be a topological space which is homotopy dominated by a finite CW complex. In [9] C. T. C. Wall introduced the obstruction $w(X)$ which is an element of $\tilde{K}_0(Z(\pi_1(X)))$ to decide when X has the homotopy type of some finite CW complex. Alternatively in [4] S. Ferry has found, in a geometric manner, an analogous obstruction $\sigma(X)$ in $Wh(X \times S^1)$. The natural question about the relation between these two obstructions was not considered in [4] (note that this question was explicitly asked by H. J. Munkholm in [10]). The purpose of this note is to fill this gap. We prove a rather expected result that these two obstructions are the same. To be more precise; we prove that $w(X)$ is the image of $\sigma(X)$ under the Bass–Heller–Swan isomorphism, thus answering the question from [10].

As a consequence we obtain the geometric proof of the Product Formula for the Wall finiteness obstructions. Originally the Product Formula was proved by S. Gersten in [5] in a purely algebraic manner. This note does not pretend to the originality, but we hope that it will a little bit clarify the geometry of the Wall finiteness obstruction.

We will assume some familiarity with the simple homotopy theory. An excellent reference is [3].

2. Wall's obstruction and simple types

In our note we will consider the Whitehead group of an arbitrary topological space following [8].

Let us recall the construction of the obstruction to the finiteness given by S. Ferry in [4].

Let X be a topological space which is homotopy dominated by a finite CW complex K , i.e. there exist maps $g : X \rightarrow K$, $f : K \rightarrow X$ such that $fg = id_X$. By the theorem of M. Mather (see [6]) $X \times S^1$ has a homotopy type of a finite CW complex. To see it we repeat his beautiful geometric argument. Namely, consider the mapping torus $T(\alpha)$ of the map $\alpha = gf : K \rightarrow K$; recall that $T(\alpha)$ is the space obtained from the mapping cylinder $M(\alpha)$ by identifying the top and bottom of $M(\alpha)$ using the identity map. Of course we can assume that up to homotopy type $T(\alpha)$ is a finite CW complex. Now the following picture shows that $X \times S^1 \simeq T(\alpha)$.




Figure 1

We will denote this homotopy equivalence by $\Phi : T(\alpha) \rightarrow X \times S^1$ and its inverse by $\Phi^{-1} : X \times S^1 \rightarrow T(\alpha)$.

DEFINITION 2.1 (S. Ferry [4]). Let $T : X \times S^1 \rightarrow X \times S^1$ be a homeomorphism given by $T(x, \theta) = (x, \bar{\theta})$. We define $\sigma(X) = \Phi_*(\tau(\Phi^{-1}T\Phi)) \in Wh(X \times S^1)$, where $\tau(\Phi^{-1}T\Phi)$ is a torsion of the homotopy equivalence $\Phi^{-1}T\Phi : T(\alpha) \rightarrow T(\alpha)$.

It turns out (see [4]) that $\sigma(X)$ is well-defined (does not depend from f , g and K) and $\sigma(X) = 0$ if and only if X is a homotopy equivalent to some finite CW complex.

The crucial role in our considerations plays the following Bass-Heller-Swan decomposition of the Wh functor (see [1], [2]).

Let X be a topological space. Then there exists a functorial direct sum decomposition

$$Wh(X \times S^1) = Wh(X) \oplus Nil(X) \oplus Nil(X) \oplus \tilde{K}_0(X)$$

where by $Nil(X)$, $\tilde{K}_0(X)$ we mean $Nil(Z(\pi_1(X)))$, $\tilde{K}_0(Z(\pi_1(X)))$ respectively. Using this we prove:

THEOREM 2.2. *Let X be a topological space which is homotopy dominated by a finite CW complex. Then the Wall finiteness obstruction $w(X)$ is a image of $\sigma(X)$ under the Bass-Heller-Swan decomposition of $\text{Wh}(X \times S^1)$.*

Proof. Let K be a finite CW complex and let $g : X \rightarrow K$, $f : K \rightarrow X$ be maps such that $fg \simeq id_X$. As previous by $T(\alpha)$ we denote the mapping torus of the map $\alpha = gf : K \rightarrow K$.

Let $\Phi : T(\alpha) \rightarrow X \times S^1$ be a homotopy equivalence. The natural infinite cyclic covering of $X \times S^1$ induces an infinite cyclic covering $I(\alpha)$ of $T(\alpha)$.




Figure 2

The space $I(\alpha)$ is an infinite CW complex with two ends ϵ_+ , ϵ_- which correspond to the two ends of the real line.

Observe that the homotopy equivalence $h = \Phi^{-1}T\Phi : T(\alpha) \rightarrow T(\alpha)$ induces a proper homotopy equivalence \tilde{h} between $I(\alpha)$ and its reversed copy $\overline{I(\alpha)}$ (reversed with respect to the ends).




Figure 3

Without loss of generality we can assume that \tilde{h} is a strong proper deformation retraction of $I(\alpha)$.

Now we proceed as in [7]. In $I(\alpha)$ consider a subcomplex L such that L is a neighborhood of ϵ_+ and $(I(\alpha) - L) \cup \overline{I(\alpha)}$ is a neighborhood of ϵ_- . Put $L_1 = \overline{I(\alpha)} \cap L$ and consider the pair (L, L_1) . It can be easily proved (see Lemma 4.5 in [7]) that the pair (L, L_1) is homotopy dominated by a pair $(L_0 \cup L_1, L_1)$, where L_0 is a finite subcomplex of L . Then the cellular chain complex $C_*(\tilde{L}, \tilde{L}_1)$ of the universal covering $p : \tilde{L} \rightarrow L$ of the pair (L, L_1) , which is a free $Z(\pi_1(I(\alpha)))$ -complex is chain homotopy dominated by the free $Z(\pi_1(I(\alpha)))$ -complex $C_*(\tilde{L}_0 \cup \tilde{L}_1, \tilde{L}_1)$; we used the notation: for every $B \subset L$, $\tilde{B} = p^{-1}(B)$. Hence we can define

$w(I(\alpha), \overline{I(\alpha)}, \epsilon_+) = w(C_*(\tilde{L}, \tilde{L}_1) \in \tilde{K}_0(Z(\pi_1(I(\alpha))))$, where $w(C_*(\tilde{L}, \tilde{L}_1))$ is the Wall obstruction. It is not difficult to see that $w(I(\alpha), \overline{I(\alpha)}, \epsilon_+)$ is well-defined i.e. does not depend of the choice of L_1 .

Now let L_- , L_+ be a neighborhoods of ϵ_- , ϵ_+ so that $I(\alpha) - L_+$, $I(\alpha) - L_-$ are again neighborhoods of ϵ_- , ϵ_+ respectively and $L_- \cup L_+ = I(\alpha)$. Then $L_- \cap L_+$ is a finite CW complex and since $I(\alpha)$ is homotopy dominated by a finite CW complex (in fact by K) then from the Mayer-Vietoris sequence

$$0 \rightarrow C_*(\tilde{L}_- \cap \tilde{L}_+) \rightarrow C_*(\tilde{L}_-) \oplus C_*(\tilde{L}_+) \rightarrow C_*(I(\alpha)) \rightarrow 0$$

we infer that $C_*(\tilde{L}_+)$ is chain homotopy dominated by a finitely generated free complex. This gives us the well defined element $w(I(\alpha), \epsilon_+) = w(C_*(\tilde{L}_+)) \in \tilde{K}_0(Z(\pi_1(I(\alpha))))$. Analogously we can define $w(\overline{I(\alpha)}, \epsilon_+) \in \tilde{K}_0(Z(\pi_1(I(\alpha))))$. An elementary property of the Wall obstructions yields:

$$w(I(\alpha), \epsilon_+) = w(I(\alpha), \overline{I(\alpha)}, \epsilon_+) + w(\overline{I(\alpha)}, \epsilon_+)$$

Observe (see Fig. 4) that in our situation $w(\overline{I(\alpha)}, \epsilon_+) = 0$ and $\tilde{h}_*(w(I(\alpha))) = w(I(\alpha), \epsilon_+)$ by a homotopy type invariance of the Wall obstruction.




Figure 4

Hence $w(I(\alpha), \overline{I(\alpha)}, \epsilon_+) = \tilde{h}_*(w(I(\alpha)))$. The Bass-Heller-Swan projection (B-H-S) : $Wh(X \times S^1) \rightarrow \tilde{K}_0(X)$ induces a natural projection $p : Wh(T(\alpha)) \rightarrow \tilde{K}_0(I(\alpha))$.

This gives the following commutative diagram:

$$\begin{array}{ccc}
 Wh(T(\alpha)) & \xrightarrow{\Phi_*} & Wh(X \times S^1) \\
 \downarrow p & & \downarrow (B-H-S) \\
 \tilde{K}_0(I(\alpha)) & \xrightarrow{\tilde{\Phi}_*} & \tilde{K}_0(X)
 \end{array}$$

where the map $\tilde{\Phi} : I(\alpha) \rightarrow X \simeq X \times R$ is induced by Φ . So we have:

$$\tilde{\Phi}_* p(\tau(\Phi^{-1}T\Phi)) = (B-H-S)\Phi_*(\tau(\Phi^{-1}T\Phi)) = (B-H-S)(\sigma(X)).$$

But $\tilde{\Phi}_* p(\tau(\Phi^{-1}T\Phi)) = \tilde{\Phi}_*(w(I(\alpha), \overline{I(\alpha)}, \epsilon_+))$ by the Proposition 4.7 in [7], hence:

$$(B-H-S)(\sigma(X)) = \tilde{\Phi}_*(w(I(\alpha))) = w(X)$$

by the homotopy type invariance of the Wall obstruction.

COROLLARY 2.3 (Product Formula). *Let X be a topological space which is homotopy dominated by a finite CW complex, and let L be a finite CW complex. Then:*

$$w(L \times X) = \chi(L) \cdot i_*(w(X))$$

where $i : X \rightarrow L \times X$ is given by $i(x) = (1_0, x)$ for some $1_0 \in L$, and $\chi(L)$ denotes the Euler characteristic of L .

Proof. Let K be a finite CW complex and $g : X \rightarrow K$, $f : K \rightarrow X$ be maps such that $fg \simeq id_X$. Let $T(\alpha)$ be the mapping torus of the map $\alpha = gf : K \rightarrow K$ and let $\Phi : T(\alpha) \rightarrow X \times S^1$ be a homotopy equivalence. The space $L \times X$ is a homotopy dominated by the finite CW complex $L \times K$ using the maps $id \times g : L \times X \rightarrow L \times K$, $id \times f : L \times K \rightarrow L \times X$. Hence we have the homotopy equivalence $\tilde{\Phi} : T(id \times \alpha) \rightarrow L \times X \times S^1$. But $T(id \times \alpha) = L \times T(\alpha)$ and without loss of the generality we can write $\tilde{\Phi} = id \times \Phi : L \times T(\alpha) \rightarrow L \times X \times S^1$. Now our finiteness obstruction is given by:

$$\sigma(L \times X) = (id \times \Phi)_*(\tau(id \times \Phi^{-1}T\Phi)) \in Wh(L \times X \times S^1).$$

By the product theorem for Whitehead torsion (see [3] for the nice and short

geometric proof) we have:

$$\tau(id \times \Phi^{-1}T\Phi) = \chi(L) \cdot j_*(\tau(\Phi^{-1}T\Phi))$$

where $j : T(\alpha) \rightarrow L \times T(\alpha)$ is given by $j(t) = (1_0, t)$, for $t \in T(\alpha)$. Hence $\sigma(L \times X) = \chi(L) \cdot i_*(\sigma(X))$, where $i_* : Wh(X \times S^1) \rightarrow Wh(L \times X \times S^1)$. Now the formula $w(L \times X) = \chi(L) \cdot i_*(w(X))$ follows from the naturality of the Bass–Heller–Swan decomposition of $Wh(X \times S^1)$.

This work was done while the author was visiting the University of Heidelberg. I am grateful to Professor Dieter Puppe for the opportunity to work there.

Note added in proof:

In fact $\sigma(X) \in \tilde{K}_0(X)$. This can be deduced from T. Chapman, Approximation results in Hilbert cube manifolds, Trans. Amer. Math. Soc. 262 (1980), 303–334, in particular, see p. 321 of this paper.

REFERENCES

- [1] H. BASS, *Algebraic K-theory*, W. A. Benjamin, New York, 1969.
- [2] H. BASS, A. HELLER and R. SWAN, *The Whitehead group of a polynomial extension*, Publ. I.H.E.S. Paris 22 (1964), 67–79.
- [3] M. COHEN, *A course in a simple homotopy theory*, Springer-Verlag, New York, 1970.
- [4] S. FERRY, *A simple-homotopy approach to the finiteness obstruction*, In Proceedings: Shape Theory and Geometric Topology, Dubrownik 1981, Lect. Not. in Math. 870 (1981), 73–81.
- [5] S. GERSTEN, *A product formula for Wall's obstruction*, Amer. J. Math. 88 (1966), 337–346.
- [6] M. MATHER, *Counting homotopy types of manifolds*, Topology 4 (1965), 93–94.
- [7] L. C. SIEBENMANN, *A total Whitehead torsion obstruction*, Comment. Math. Helv. 45 (1970), 1–48.
- [8] R. STÖCKER, *Whideheadgruppe topologischer Räume*, Inventiones Math. 9 (1970), 271–278.
- [9] C. T. C. WALL, *Finiteness conditions for CW complexes*, Ann. of Math. (2) (1965), 55–64.
- [10] Zentralblatt für Mathematik, Mathematics Abstracts, 472 (1982) 57014, p. 318.

Mathematisches Institut,
Universität Heidelberg,
Im Neuenheimer Feld 288,
D-6900 Heidelberg,
West Germany

Institute of Mathematics,
University of Gdańsk,
Wita Stwosza 57,
PL-80-952 Gdańsk,
Poland.

Received November 17, 1982