

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 58 (1983)

Artikel: A reciprocity law for K2-traces.
Autor: Rosset, Shmuel / Tate, John
DOI: <https://doi.org/10.5169/seals-44587>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A reciprocity law for K_2 -traces

SHMUEL ROSSET and JOHN TATE

Suppose $E \subset F$ is a finite field extension and let

$$\text{Tr}: K_2(F) \rightarrow K_2(E)$$

be the trace map (also called transfer, see [5, §14]). If $x, y \in F^*$ and $\{x, y\}$ is the corresponding symbol in $K_2(F)$ then we know, since $K_2(E)$ is generated by symbols, that $\text{Tr}_{F/E}(\{x, y\})$ can be expressed as a sum of symbols. In this paper we give an algorithm for computing such an expression explicitly (cf. the proposition in §3). The algorithm is based on a reciprocity law (§2) and involves repeated polynomial division with remainder, like the Euclidean algorithm. The proof works not only for Milnor's K_2 , but for functors sufficiently like K_2 , which we define in §1 and call Milnor functors. This abstraction is useful for it yields as a corollary (§3) the fact that the canonical map from K_2 to any Milnor functor commutes with traces. Another corollary is that, if $(F:E) = n$, then $\text{Tr}_{F/E}(\{x, y\})$ can be written as a sum of n symbols (or less). On the other hand this is also the best bound: in §4 we give an example, using division algebras, of a symbol whose trace is not a sum of less than n symbols.

One of us (S.R) would like to thank David Saltman for a conversation which helped realize the example in section 4.

1. Milnor functors

Let K be a fixed base field and let \mathfrak{C} be the category of commutative finite dimensional k -algebras.

DEFINITION. A *Milnor functor over k* is a functor $M: \mathfrak{C} \rightarrow$ (Abelian groups) together with

- (i) For each $A \in \mathfrak{C}$ a bilinear map $\varphi = \varphi_A: A^* \times A^* \rightarrow M(A)$;
- (ii) For each extension $A \rightarrow B$ in \mathfrak{C} such that B is a projective A -module, a homomorphism $\text{Tr}_{B/A}: M(B) \rightarrow M(A)$; such that the following properties hold.

(φ) The maps φ are functorial, i.e., induce a morphism of functors from the functor $A \mapsto A^* \times A^*$ to the functor $A \mapsto M(A)$, and satisfy

$$\varphi_A(a, 1-a) = 0, \quad \text{if } a \in A^* \text{ and } 1-a \in A^*,$$

$$\varphi_A(a, -a) = 0, \quad \text{if } a \in A^*.$$

(Tr) if $A \rightarrow B \rightarrow C$ are \mathfrak{C} -morphisms such that C is projective over B and B over A , then

$$\text{Tr}_{C/A} = \text{Tr}_{B/A} \circ \text{Tr}_{C/B}$$

(Tr- φ) If $A \rightarrow B$ is a \mathfrak{C} -morphism with B projective as A -module, and if $x \in A^*$, $y \in B^*$ then

$$\text{Tr}_{B/A} \varphi_B(x, y) = \varphi_A(x, N_{B/A}y),$$

where $N_{B/A} : B^* \rightarrow A^*$ is the usual norm:

$$N_{B/A}(y) = \det \quad (\text{multiplication by } y).$$

EXAMPLE 1. Milnor's K_2 ; see [5] and [6].

EXAMPLE 2. Assume that the characteristic of k does not divide a given integer n and let μ_n denote the sheaf on n -th roots of 1 on the étale site over $\text{Spec } A$; here A is a given element in $\text{Ob}(\mathfrak{C})$. By Kummer theory

$$H^1(\text{Spec } A, \mu_n) = A^*/(A^*)^n.$$

The cup product

$$H^1(\text{Spec } A, \mu_n) \times H^1(\text{Spec } A, \mu_n) \rightarrow H^2(\text{Spec } A, \mu_n^{\otimes 2}) = M(A)$$

provides us with a context satisfying (i) and (ii). We refer to Milne's book [4] for details. The existence of a trace can probably be extracted from [7, exp. XVII]. However, this Milnor functor can be expressed entirely in terms of Galois cohomology and the trace in terms of corestriction, as follows. For $A \in \mathfrak{C}$, $\alpha \in M(A)$, and $x \in \text{Spec } A$, let $\alpha(x) \in M(A/x)$ be the image of α under the residue class map $A \rightarrow A/x$. then the map

$$\alpha \mapsto (\alpha(x))_{x \in \text{Spec } A}$$

gives an isomorphism

$$M(A) \xrightarrow{\sim} \prod_{x \in \text{Spec } A} M(A/x). \quad (*)$$

For each $x \in \text{Spec } A$, A/x is a finite extension field of k . If E is a finite extension field of k , then

$$M(E) = H^2(\text{Gal}(E_s/E), \mu_n(E_s) \otimes \mu_n(E_s)),$$

where E_s is a separable algebraic closure of E . The map φ_A is characterized in terms of the isomorphism $(*)$ by

$$(\varphi_A(a, b))(x) = \varphi_{A/x}(a(x), b(x))$$

for each $x \in \text{Spec } A$, where $a(x)$ (resp. $b(x)$) is the residue mod x of a (resp. b), and for a field E the map

$$\varphi_E : E^* \times E^* \rightarrow M(E)$$

is the Galois cohomology symbol (cf. [8]) characterized by $\varphi(a, b) = da \cup db$, where $d : E^* \rightarrow H^1(\text{Gal}(E_s/E), \mu_n(E_s))$ is the connecting homomorphism in the exact cohomology sequence associated with

$$0 \rightarrow \mu_n(E_s) \rightarrow E_s^* \xrightarrow{n} E_s^* \rightarrow 0.$$

Let $A \rightarrow B$ be an extension in \mathfrak{C} such that B is a projective A -module. Then for each $x \in \text{Spec } A$ and each $y \in \text{Spec } B$ lying over x , the local ring B_y is a free A_x -module; let $r(y/x)$ denote its rank. Let $E_x = A/x$ and let F_y be the field between E_x and B/y such that F_y/E_x is separable and $(B/y)/F_y$ purely inseparable. Then the ratio

$$q(y/x) \stackrel{\text{defn}}{=} \frac{r(y/x)}{[F_y : E_x]}$$

is an integer, and the M -trace from B to A is characterized in terms of the isomorphism $(*)$ by

$$(\text{Tr}_{B/A}\beta)(x) = \sum_{y|x} q(y/x) \text{cor}_{F_y/E_x}(\beta(y)),$$

where cor is the corestriction in Galois cohomology, and we identify $M(B/y)$ with $M(F_y)$ via the isomorphism induced by the inclusion $F_y \hookrightarrow B/y$.

In case $E \in \mathbb{C}$ is a field containing a primitive n -th root of unity ζ , we can identify $M(E)$ with the group $\text{Br}_n(E)$ of elements of order n in the Brauer group of E in such a way that

$$(a, b)_M = \text{the Brauer class of } A_\zeta(a, b)$$

where $A_\zeta(a, b)$ denote the cyclic algebra generated over E by elements X and Y subject to the relations

$$X^n = a, \quad Y^n = b, \quad XY = \zeta YX;$$

(cf. [5], p. 143).

EXAMPLE 3. The dlog symbol, see [1]. If A is a k algebra in \mathbb{C} let $\Omega_{A/k}^1$ be the A -module of Kähler differentials of A over k , and let $\Omega_{A/k}^2$ be its second exterior power. Define

$$\text{dlog}: A^* \rightarrow \Omega_{A/k}^1$$

by $\text{dlog}(f) = f^{-1} \cdot df$. It is simple to verify that Ω^2 and $\text{dlog} \wedge \text{dlog}$ satisfy axioms (i), (ii) above. The existence of a good trace is a non-trivial fact [2].

2. Reciprocity

Let M be a Milnor functor over k . In this section we shall write the M -symbol $\varphi_E(x, y)$ by

$$(x, y)_E, \quad \text{or} \quad (x, y)$$

if E is evident.

Let K be a field of finite degree over k . For relatively prime non-zero polynomials $f(T), g(T)$ in $K[T]$ we define a new kind of symbol (f/g) . Its values are in the group $M(K)$ and it is defined by the following requirements.

1) It is additive in g , i.e. if g_1, g_2 are both prime to f then

$$\left(\frac{f}{g_1 g_2} \right) = \left(\frac{f}{g_1} \right) + \left(\frac{f}{g_2} \right)$$

2) It is 0 if g is a constant or $g = T$.

3) If g is monic irreducible $\neq T$ and x is a root of $g(T)$ then

$$\left(\frac{f}{g}\right) = \text{Tr}_{K(x)/K}(x, f(x))_{K(x)}.$$

It is clear that, thus defined, the symbol (f/g) is additive in f , as well as in g , and it depends only on the residue class of f modulo (g) . As function of g it depends only on the ideal generated by g in the ring $K[T, T^{-1}]$.

To formulate the reciprocity law satisfied by (f/g) we introduce some notation: if

$$p(T) = a_n T^n + a_{n-1} T^{n-1} + \cdots + a_m T^m$$

with $a_m a_n \neq 0$, let

$$p^*(T) = (a_m T^m)^{-1} p(T)$$

$$c(p) = (-1)^n a_n.$$

Reciprocity law

$$\left(\frac{f}{g}\right) = \left(\frac{g^*}{f}\right) - (c(g^*), c(f)). \quad (**)$$

Proof. We first dispose of a few trivial cases. If g is a constant or T it is easily checked that both sides are 0, so we assume henceforth that $g(T)$ is monic irreducible $\neq T$. Let x be a root of $g(T)$. If $f(T)$ is a constant c then the left side of $(**)$ is

$$\begin{aligned} \text{Tr}_{K(x)/K}(x, c)_{K(x)} &= (N_{K(x)/K}x, c)_k \\ &= ((-1)^{\deg(g)} \cdot g(0), c) = -((-1)^{\deg(g)} \cdot g(0)^{-1}, c) \\ &= -(c(g^*), c(f)) \end{aligned}$$

which is equal to the right hand side since $(g^*/f) = 0$, by definition.

A similar computation using $(x, -x) = 0$ works when $f(T) = T$ so we now assume that both f and g are monic irreducible, and not T .

Let x be a root of g and y a root of f . Let

$$A = K(x) \otimes_K K(y).$$

$K(x)$ and $K(y)$ are naturally imbedded in A and we identify them as such. Then

the elements $x, y, x - y$ are invertible in A , indeed the norm

$$N_{A/K(x)}(x - y) = f(x)$$

is invertible, so $x - y$ is.

The identity

$$(x, x - y) = \left(y, \frac{y - x}{-x} \right) + (x, -1)$$

follows from a little computation with the relations $(u, 1 - u) = (u, -u) = 0$. We use it to compute the same thing in two ways

$$\begin{aligned} \text{Tr}_{A/K}(x, x - y) &= \text{Tr}_{K(x)/K} \text{Tr}_{A/K(x)}(x, x - y) \\ &= \text{Tr}_{K(x)/K}(x, N_{A/K(x)}(x - y)) \\ &= \text{Tr}_{K(x)/K}(x, f(x)) = \left(\frac{f}{g} \right). \end{aligned}$$

$$\begin{aligned} \text{Tr}_{A/K} \left(y, \frac{y - x}{-x} \right) &= \text{Tr}_{K(y)/K} \text{Tr}_{A/K(y)} \left(y, \frac{y - x}{-x} \right) \\ &= \text{Tr}_{K(y)/K} \left(y, \frac{N_{A/K(y)}(y - x)}{N_{A/K(y)}(-x)} \right) \\ &= \text{Tr}_{K(y)/K} \left(y, \frac{g(y)}{g(0)} \right) \\ &= \text{Tr}_{K(y)/K}(y, g^*(y)) = \left(\frac{g^*}{f} \right). \end{aligned}$$

Finally

$$\begin{aligned} \text{Tr}_{A/K}(x, -1) &= \text{Tr}_{K(y)/K} \text{Tr}_{A/K(y)}(x, -1)_A \\ &= \text{Tr}_{K(y)/K}(N_{A/K(y)}x, -1)_{K(y)} \\ &= \text{Tr}_{K(y)/K}(N_{K(x)/K}x, -1)_{K(y)} \\ &= (c(g^*)^{-1}, (-1)^{\deg(f)}) = -(c(g^*), c(f)). \end{aligned}$$

Here we used the obvious fact that

$$N_{A/K(y)}(x) = N_{K(x)/K}(x).$$

This completes the proof of the reciprocity law.

3. Consequences

Let $E \subset F$ be a finite extension of fields finite over k , and let $x, y \in F^*$. Then

$$\text{Tr}_{F/E}(x, y) = \left(\frac{f}{g} \right)$$

where $g(T) \in E[T]$ is the monic irreducible polynomial with root x and $f(T) \in E[T]$ is the polynomial of smallest degree such that $N_{F/E(x)}y = f(x)$.

PROPOSITION. *Let $g_0, g_1, \dots, g_m \neq 0, g_{m+1} = 0$ be the sequence of polynomials defined by:*

$$g_0 = g, \quad g_1 = f,$$

and for $i \geq 1$

$$g_{i+1} = \text{the remainder of the division of } g_{i-1}^* \text{ by } g_i,$$

as long as $g_i \neq 0$. We have then

$$1 \leq m \leq \deg g = [E(x) : E] \leq [F : E]$$

and

$$\text{Tr}_{F/E}(x, y) = - \sum_{i=1}^m (c(g_{i-1}^*), c(g_i)).$$

By the reciprocity law, we find by induction on j , using $(g_{i-1}^* / g_i) = (g_{i+1} / g_i)$:

$$\left(\frac{g_1}{g_0} \right) = - \sum_{i=1}^j (c(g_{i-1}^*), c(g_i)) + \left(\frac{g_{j-1}^*}{g_j} \right)$$

for $1 \leq j \leq m$. But the last non-zero polynomial g_m is a constant because it divides the relatively prime polynomials g_0 and g_1 . Hence $(g_{m-1}^* / g_m) = 0$, and the proposition follows on putting $j = m$; We have $m \leq \deg g$ because the degrees of the polynomials in the sequence are strictly decreasing, and $m \geq 1$ because $f \neq 0$.

COROLLARY 1. *If $[F : E] = r$ and $x, y \in F^*$, then $\text{Tr}_{F/E}(x, y)$ is a sum of at most r symbols.*

The sequence of polynomials in the proposition depends only on F , E , x , and y , not on the Milnor functor M . Thus the trace of a symbol $(x, y)_M$ has an expression as a sum of symbols which is *independent of the Milnor functor M* ; on symbols, the trace is uniquely determined. Any morphism $M_1 \rightarrow M_2$ of Milnor functors which carries each symbol $(a, b) \in M_1(A)$ to the “same” symbol $(a, b) \in M_2(A)$ must therefore commute with $\text{Tr}_{F/E}$ on symbols. In particular, letting $R_F : K_2(F) \rightarrow M(F)$ be the homomorphism (whose existence and unicity are guaranteed by Matsumoto’s theorem) such that $R_F(\{a, b\}) = (a, b)_M$ for $a, b \in F^*$, and similarly R_E , we have

COROLLARY 2. *The diagram*

$$\begin{array}{ccc} K_2 & \xrightarrow{R_F} & M(F) \\ \text{Tr}_{F/E} \downarrow & & \downarrow \text{Tr}_{F/E} \\ K_2(E) & \xrightarrow{R_E} & M(E) \end{array}$$

is commutative.

4. An example

We have just proved that if $[F:E]=r$ and $x, y \in F^*$ then $\text{Tr}_{F/E}(x, y)$ is a sum of r symbols. Yet it is known that in some cases, e.g. global or local fields, every element of K_2 (say) is a symbol [8, 3], so it is well to give an example where $\text{Tr}(x, y)$ cannot be written as a sum of fewer than r symbols. For this it will suffice to work with the functor of Example of Section 1.

Let $n \geq 2$ and $r \geq 1$ be integers. Let k_0 be a field containing a primitive n -th root of unity, ζ . Let $u_1, v_1, \dots, u_r, v_r$ be $2r$ independent variable over k_0 and let

$$F = k_0(u_1, v_1; u_2, v_2; \dots; u_r, v_r)$$

be the field they generate. Let M be the Milnor functor of Example 2.

LEMMA. *The element $\beta = \sum_{i=1}^r (u_i, v_i)$ in $M(F)$ is not a sum of fewer than r symbols.*

Proof. We use the identification $M(F) \xrightarrow{\sim} \text{Br}_n(F)$ discussed at the end of

Example 2. For $1 \leq i \leq r$ let B_i by the cyclic algebra over F generated by elements X_i and Y_i subject to the relations

$$X_i^n = u_i, \quad Y_i^n = v_i, \quad X_i Y_i = \zeta Y_i X_i,$$

so that (u_i, v_i) is the Brauer class of B_i . Then β is the Brauer class of $B = \bigotimes_{i=1}^r B_i$, an algebra of dimension n^{2r} over F . We will show B is a division algebra. This will prove the lemma, for it shows that β cannot be the Brauer class of an algebra of dimension less than n^{2r} , and consequently cannot be a sum of fewer than r symbols.

If B were not a division algebra it would have zero divisors, and multiplying these zero divisors by a common denominator of their coefficients in F relative to the basis

$$\{X_1^{l_1} Y_1^{m_1} \cdots X_r^{l_r} Y_r^{m_r}\} \quad (0 \leq l_i, m_i < n)$$

for B over F , we would find zero divisors in the ring

$$R = k_0[u_1, v_1, \dots, u_r, v_r][X_1, Y_1, \dots, X_r, Y_r] = k_0[X_1, y_1, \dots, X_r, Y_r].$$

But this ring has no zero divisors, for it has a basis over k_0 consisting of the monomials

$$X_1^{l_1} Y_1^{m_1} \cdots X_r^{l_r} Y_r^{m_r}$$

with l_i, m_i integers ≥ 0 , and the product of two such monomials is a power of ζ times the monomial obtained by adding exponents. Hence, if we order the monomials by the lexicographical order of their exponent sequences, the product of two non-zero polynomials will contain the product of the highest terms in the two factors with a non-zero coefficient, so will not be 0. This proves the lemma.

Let σ be the automorphism of F which is identity on k_0 and acts on the variables by

$$\begin{aligned} \sigma u_i &= u_{i+1}, & 1 \leq i \leq r; \quad u_{r+1} &= u_1, \\ \sigma v_i &= v_{i+1}, & 1 \leq i \leq r; \quad v_{r+1} &= v_1. \end{aligned}$$

Let G be the cyclic group of order r generated by σ , and let $E = F^G$.

PROPOSITION. *The image of $\{u_1, v_1\}$ under $\text{Tr}_{F/E} : K_2 F \rightarrow K_2 E$ is not a sum of fewer than r symbols.*

Proof. We use the commutativity of

$$\begin{array}{ccc} K_2(F)/nK_2(F) & \xrightarrow{R_F} & \text{Br}_n(F) \\ \text{Tr} \downarrow & & \downarrow \text{Tr} \\ K_2(E)/nK_2(E) & \xrightarrow{R_E} & \text{Br}_n(E). \end{array}$$

and the rule

$$\text{res}_{E/F} \text{Tr}_{F/E} \alpha = \sum_{\tau \in G} \tau \alpha$$

for $\alpha \in \text{Br } F$. If $\text{Tr } \{u_1, v_1\}$ were a sum of $s < r$ symbols so also would be

$$\begin{aligned} \text{res } R_E \text{Tr } \{u_1, v_1\} &= \text{res } \text{Tr } R_F \{u_1, v_1\} = \text{res } \text{Tr } (u_1, v_1) \\ &= \sum_{\tau \in G} \tau(u_1, v_1) = \sum_{i=1}^r (u_i, v_i) = \beta, \end{aligned}$$

contradicting the lemma.

REFERENCES

- [1] BLOCH, S., *K_2 and algebraic cycles*. Ann. Math. 99 (1974), 349–379.
- [2] HARTSHORNE, R., *Residues and Duality*, Springer Lecture notes #20 (1966).
- [3] LENSTRA, H. W., *K_2 of global fields consists of symbols*, in *Algebraic K-theory* Evanston 1976, Springer Lecture notes #551, pp. 69–73.
- [4] MILNE, J. S., *Etale Cohomology*, Princeton Univ. Press, 1980.
- [5] MILNOR, J., *Introduction to algebraic K-theory*, Ann. Math. Studies #72, Princeton, 1971.
- [6] QUILLEN, D., *Higher algebraic K-theory I*, in *Algebraic K-theory I*, Springer Lecture notes #341, 1973.
- [7] SGA 4, Tome 3, Springer Lecture notes #305, 1972.
- [8] TATE, J., *Relations between K_2 and Galois cohomology*, Inv. Math. 36 (1976), 257–274.

Dept. of Mathematical Sciences

Tel-Aviv University

Tel-Aviv, Israel

and

Dept. of Mathematics

Harvard University

Cambridge, MA 02138, USA

Received August 18, 1982