Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 59 (1984)

Artikel: A sharp four dimensional isoperimetric inequality.
Autor: Croke, Christopher B.
DOI: https://doi.org/10.5169/seals-45390

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 06.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Comment. Math. Helvetici **59** (1984) 187–192 0010–2571/84/020187–06$01.50 + 0.20/0
© 1984 Birkhäuser Verlag, Basel

A sharp four dimensional isoperimetric inequality

CHRISTOPHER B. CROKE*

Introduction

Let $(M, \partial M)$ be a compact n-dimensional Riemannian manifold (with boundary) of non positive sectional curvature. Assume further that every geodesic ray in M minimizes length up to the point it hits the boundary. In this paper we show:

THEOREM. *If* $(M, \partial M)$ *is as above* $(n \geq 3)$ *then* $\mathrm{Vol}\,(\partial M)^n \geq C(n)\,\mathrm{Vol}\,(M)^{n-1}$ *where*

$$C(n) = \frac{\alpha(n-1)^{n-1}}{\alpha(n-2)^{n-2}\left\{\int_0^{\pi/2} \cos(t)^{n/n-2} \sin(t)^{n-2}\, dt\right\}^{n-2}}$$

and $\alpha(n)$ *represents the volume of the unit n sphere. If* $n \neq 4$ *equality never holds. If* $n = 4$ *equality holds if and only if M is isometric to a flat ball.*

This answers a long standing conjecture in dimension 4. The conjecture states that for $(M, \partial M)$ a compact domain in a complete simply connected manifold of non-positive curvature (which implies the condition of the theorem) we have $\mathrm{Vol}\,(\partial M)^n \geq \bar{C}(n)\,\mathrm{Vol}\,(M)^{m-1}$ for $\bar{C}(n) = n^{n-1}\alpha(n-1)$, with equality holding if and only if M is isometric to a flat ball. It is an easy computation to see that $C(4) = \bar{C}(4)$. The conjecture was proved in dimension 2 by Beckenbach and Radó (see [B–R]) in 1933, and is open in all dimensions except 2, and now 4.

This conjecture is a special case of a more general conjecture (see [A]) where an upper bound K (not necessarily 0) on the curvature is assumed. The more general conjecture was proved in dimension 2 by Aubin (see [A]), and for constant curvature in all dimensions by Schmidt (see [Sc]).

The isoperimetric constants are related to Sobolev constants (see [A], [Bo], and [FF] among other). In particular we see that for a domain D in a simply

* Supported by NSF Grant #MCS-01780

connected n-dimensional Riemannian manifold of non-positive curvature and for any $g \in H_0^1(M)$ we have

$$\int_D \|dg\| \geq C(n)\left\{\int_D |g|^{n/n-1}\right\}^{(n-1)/n}$$

where $C(4)$ is the flat constant.

Such isoperimetric inequalities are interesting even with non-sharp constants. Previous non-sharp versions of the theorem are consequences of results in [H–S] and [C]. The constants $C(n)$ given here are the best known to the author in all dimensions (greater than 2). In particular $C(3) = 32\pi$ while $\bar{C}(3) = 36\pi$.

Notation and definitions

We will use the notation of [C]. Let $UM \xrightarrow{\pi} M$ represent the unit sphere bundle with the canonical (local product) measure. For $v \in UM$, let γ_v be the geodesic with $\gamma_v'(0) = v$ and let $\xi^t(v)$ represent the geodesic flow (i.e. $\xi^t(v) = \gamma_v'(t)$). For $v \in UM$ we let $l(v) = \max\{t \mid \gamma_v(t) \in M\}$. Note $\xi^t(v)$ is defined for $t \leq l(v)$ and $\gamma_v(l(v)) \in \partial M$.

For $p \in \partial M$ let N_p be the inwardly pointing unit normal vector to ∂M at p. Let $U^+\partial M \xrightarrow{\pi} \partial M$ be the bundle of inwardly pointing unit vectors (i.e. $U^+\partial M = \{u \in UM \mid \pi(u) \in \partial M$ and $\langle u, N_{\pi(u)}\rangle > 0\}$). We let $U_p^+\partial M$ represent $\pi^{-1}(p)$. For $u \in U^+\partial M$ we will use $\cos(u)$ to represent $\langle u, N_{\pi(u)}\rangle$. The measure on $U^+\partial M$ is the local product measure du where the measure of the fibre is that of the unit upper hemisphere.

The proof

The main tool in the proof is a formula due to Santalo:

$$\text{(i)} \quad \int_{UM} f(v)\, dv = \int_{U^+\partial M} \int_0^{l(u)} f(\xi^t(u)) \cos(u)\, dt\, du$$

for all integrable functions f. The formula takes this form in our case since all geodesics in M hit ∂M. For a proof see [Sa] pp. 336–338 or [B] p. 286.

From this we derive:

LEMMA 1. a) $\mathrm{Vol}\,(M) = 1/\alpha(n-1) \int_{U^+\partial M} l(u) \cos(u)\, du$
b) *For all integrable functions* g

$$\int_{U^+\partial M} g(u) \cos(u)\, du = \int_{U^+\partial M} g(\mathrm{ant}\,(u)) \cos(u)\, du$$

where $\mathrm{ant}\,(u) = -\gamma'_u(l(u))$.

Proof. Part a) follows directly from (i) by letting $f(v) \equiv 1$ and integrating the t. That is

$$\alpha(n-1)\,\mathrm{Vol}\,(M) = \int_{UM} dv = \int_{U^+\partial M} \int_0^{l(u)} \cos(u)\, dt\, du = \int_{U^+\partial M} l(u) \cos(u)\, du.$$

To prove part b) we first note that (i) says that the geodesic flow ξ is a measure preserving map from Q to UM where $Q = \{(u, t) \mid u \in U^+\partial M \text{ and } 0 \le t \le l(u)\}$ is given the measure $\cos(u)\, dt\, du$. ξ has an inverse (smooth almost everywhere) ξ^{-1} which is also measure preserving, for $v \in UM$ $\xi^{-1}(v) = (-\gamma'_{-v}(l(-v)), l(-v))$. Since the antipodal map $-1 : UM \to UM$ is also measure preserving we have $\xi^{-1} \circ (-1) \circ \xi : Q \to Q$ is measure preserving. Since $\xi^{-1} \circ (-1) \circ \xi(u, t) = (\mathrm{ant}\,(u), l(u) - t)$ we see that for every integrable $G : Q \to \mathbb{R}$ we have:

$$\int_{U^+\partial M} \int_0^{l(u)} G(u, t) \cos(u)\, dt\, du = \int_{U^+\partial M} \int_0^{l(u)} G(\mathrm{ant}\,(u), l(u) - t) \cos(u)\, dt\, du$$

To complete the proof of part b) simply take $G(u, t) = g(u)/l(u)$ and integrate the t (note: $l(\mathrm{ant}\, u) = l(u)$).

LEMMA 2. a) $\int_{U^+\partial M} l(u)^{n-1}/\cos(\mathrm{ant}\, u)\, du \le \mathrm{Vol}\,(\partial M)^2$ *with equality holding if and only if M is flat and convex.*

b) $\int_{U^+\partial M} (\cos(\mathrm{ant}\, u))^{1/n-2} (\cos(u))^{n-1/n-2}\, du \le \mathrm{Vol}\,(\partial M) \cdot C_2(n)$ *where* $C_2(n) = \alpha(n-2) \int_0^{\pi/2} \cos^{n/n-2}(t) \sin^{n-2}(t)\, dt$. *Equality holds if and only if* $\cos(u) = \cos(\mathrm{ant}\, u)$ *almost everywhere.*

Remark. "almost everywhere" above can be replaced with "everywhere" but it is not worth going into.

Proof. Let dx be the volume form on M and dp the volume form on ∂M. Let $q \in \partial M$. In normal polar coordinates (u, r) about q in the region $\operatorname{Exp}\{tu \mid u \in U_q^+ \partial M \text{ and } 0 \leq t \leq l(u)\}$ we have $dx = F(u, r)\, du\, dr$ for some function $F(u, r)$. Let $A = \operatorname{Exp}\{tu \mid t = l(u)\}$. Then $A \subset \partial M$ and dp on A is precisely $(F(u, l(u))/\cos(\operatorname{ant} u))\, du$. Thus we see

$$\int_{U_q^+\partial M} \frac{F(u, l(u))}{\cos(\operatorname{ant} u)}\, du = \operatorname{Vol}(A) \leq \operatorname{Vol}(\partial M).$$

Equality holds if and only if $A = \partial M$. That is, M is (geodesically) star shaped from q.

Integrating over q we get

$$\int_{U^+\partial M} \frac{F(u, l(u))}{\cos(\operatorname{ant} u)}\, du \leq \operatorname{Vol}(\partial M)^2$$

with equality holding if and only if M is convex. Part a) now follows since M having non-positive curvature implies $F(u, l(u)) \geq l(u)^{n-1}$ with equality if and only if the sectional curvatures of all sections containing $\gamma_u'(t)$ for some t, are 0 (see [B–C] Section 11.10).

To prove part b) we apply a Schwarz inequality and Lemma 1b.

$$\begin{aligned}
&\int_{U^+\partial M} (\cos(\operatorname{ant} u))^{1/n-2}(\cos(u))^{n-1/n-2}\, du \\
&= \int_{U^+\partial M} (\cos(\operatorname{ant} u))^{1/n-2}(\cos(u))^{1/n-2} \cos(u)\, du \\
&\leq \left\{\int_{U^+\partial M} (\cos(\operatorname{ant} u))^{2/n-2} \cos(u) du\right\}^{1/2} \cdot \left\{\int_{U^+\partial M} (\cos(u)^{2/n-2} \cos(u)\, du\right\}^{1/2} \\
&= \int_{U^+\partial M} (\cos(u))^{n/n-2}\, du = \int_{\partial M} \left(\int_{U^+\partial M} (\cos(u)^{n/n-2}\, du\right) dq \\
&= \operatorname{Vol}(\partial M) \cdot C_2(n)
\end{aligned}$$

In order for equality to hold we need to have equality in the inequality, i.e. $\cos(\operatorname{ant} u) = K \cos(u)$ almost everywhere for some constant K. Since the maximum values of both $\cos(\operatorname{ant} u)$ and $\cos(u)$ are 1 it is clear that K must be 1.

Proof of the theorem. By Lemma 1a and a Hölder inequality we have

$$\begin{aligned}\mathrm{Vol}\,(M) &= \frac{1}{\alpha(n-1)}\int_{U^+\partial M} l(u)\cos(u)\,du \\ &= \frac{1}{\alpha(n-1)}\int_{U^+\partial M} \frac{l(u)}{(\cos(\mathrm{ant}\,u))^{1/n-1}}(\cos(\mathrm{ant}\,u))^{1/n-1}\cos(u)\,du \\ &\le \frac{1}{a(n-1)}\left\{\int_{U^+\partial M}\frac{l(u)^{n-1}}{\cos(\mathrm{ant}\,u)}\,du\right\}^{1/n-1} \\ &\quad\cdot\left\{\int_{U^+\partial M}(\cos(\mathrm{ant}\,u))^{1/n-2}(\cos(u))^{n-1/n-2}\,du\right\}^{n-2/n-1}.\end{aligned}$$

Applying Lemmas 2a and 2b we get

$$\mathrm{Vol}\,(M) \le \frac{1}{\alpha(n-1)}(\mathrm{Vol}\,(\partial M))^{2/n-1}\cdot(\mathrm{Vol}\,(\partial M))^{n-2/n-1}C_2(n)^{n-2/n-1}.$$

hence

$$C(n)\,\mathrm{Vol}\,(M)^{n-1} = \frac{\alpha(n-1)^{n-1}}{C_2(n)^{n-2}}(\mathrm{Vol}\,(M))^{n-1} \le (\mathrm{Vol}\,(\partial M))^n.$$

In order for equality to hold we must have equality in Lemmas 2a and 2b as well as the above Hölder inequality. By Lemma 2a we see that M must be flat and hence the theorem follows from the classical result in $\mathbb{R}^n$, since $C(4)$ is sharp and $C(n)$ for $n \neq 4$ is not. One can also see the $n=4$ case directly. Equality in the Hölder inequality gives

$$\frac{l^3(u)}{\cos(\mathrm{ant}\,u)} = K\cos(\mathrm{ant}\,u)^{1/2}\cos(u)^{3/2}$$

almost everywhere for some constant K. By the equality condition in Lemma 2b we see $l(u) = 2r\cos(u)$ for some constant r. It is now easy to see (since M is flat) that M is a ball of radius r.

Remark. A similar equality analysis for $n \neq 4$ would require:

1) M flat and convex
2) $\cos(\mathrm{ant}\,u) = \cos(u)$
3) $l(u) = K\cos^{2/n-2}(u)$.

No such M exists.

REFERENCES

[A] T. AUBIN, *Problèmes isopérimétriques et espaces de Sobolev*, J. Differential Geometry *11* (1976), 573–598.

[B] M. BERGER, *Lectures on geodesics in Riemannian geometry*, Tata Institute, Bombay, 1965.

[B–R] E. F. BECKENBACH and T. RADÓ, *Subharmonic functions and surfaces of negative curvature*, Trans, Amer. Math. Soc. *35* (1933), 662–674.

[B–C] R. BISHOP and R. CRITTENDEN, *Geometry of manifolds*, Academic Press, 1964.

[Bo] E. BOMBIERI, *Theory of minimal surfaces, and a counterexample to the Bernstein conjecture in high dimension*, Lecture Notes, Courant Inst., 1970.

[C] C. CROKE, *Some isoperimetric inequalities and eigenvalue estimates*, Ann. Scient. Éc. Norm. Sup., 4e série, t. *13*, 1980, 419–435.

[F–F] H. FEDERER and W. FLEMMING, *Normal integral currents*, Ann. of Math. *72* (1960), 458–520.

[H–S] D. HOFFMAN and J. SPRUCK, *Sobolev and isoperimetric inequalities for Riemannian submanifolds*, Communications Pure Appl. Math. *27* (1974), 715–727; correction, ibid. *28* (1975), 765–766.

[Sa] L. A. SANTALÓ, *Integral geometry and geometric probability*, (Encyclopedia of Mathematics and Its Applications), Addison-Wesley, London–Amsterdam–Don Mills, Ontario–Sydney–Tokyo, 1976.

[Sc] E. SCHMIDT, *Beweis der isoperimetrischen Eigenschaft der Kugel in hyperbolischen und spharischen Raum jeder Dimensionenzahl*, Math. Z. *49* (1943/44), 1–109.

University of Pennsylvania
Philadelphia, PA 19104 USA

Received March 28, 1983