

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 59 (1984)

Artikel: On the homotopy groups of a finite dimensional space.
Autor: McGibbon, C.A. / Neisendorfer, J.A.
DOI: <https://doi.org/10.5169/seals-45395>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On the homotopy groups of a finite dimensional space

C. A. McGIBBON and J. A. NEISENDORFER

The purpose of this note is to prove the following.

THEOREM 1. *Let X be a 1-connected space and p a prime number such that*

- (i) $H_n(X; \mathbb{Z}/p) \neq 0$ for some $n > 0$, and
- (ii) $H_n(X; \mathbb{Z}/p) = 0$ for all n sufficiently large.

Then for infinitely many n , $\pi_n X$ contains a subgroup of order p . \square

Thirty years ago, J.-P. Serre conjectured such a result for $p = 2$ [3, page 219]. He arrived at this conjecture after having proved the 2-primary part of the following result.

THEOREM 2. *Let X and p be as in Theorem 1. Moreover, assume that $H_*(X; \mathbb{Z})$ is of finite type. Then for infinitely many n , $\pi_n X$ contains an element whose order either equals p or is infinite. \square*

Serre's proof in this case used, among other things, Poincaré series and methods of analytic number theory. Later Y. Umeda, [5], showed that these methods could be modified to work for odd primes as well.

Notice that Theorem 1 represents an improvement over Theorem 2 in two respects. First, of course, it establishes the existence of torsion in $\pi_* X$ in infinitely many dimensions and, second, it does so without the hypothesis of finite type.

The key ingredient in our proof is the following recent result of Haynes Miller, [1].

THEOREM 3. *Let X and p be as in Theorem 1. Let $B = B\mathbb{Z}/p$, the classifying space for the group \mathbb{Z}/p . Then the space of pointed maps from B to X is weakly contractible; that is, $\pi_n(\text{map}_*(B, X)) = 0$ for all $n \geq 0$. \square*

Of course, in this theorem, B may also be regarded as the Eilenberg–MacLane space $K(\mathbb{Z}/p, 1)$ or, in the case when $p = 2$, as the infinite real projective space RP^∞ . We should add that we have not stated Miller's result in its most general form. However, for our purposes the statement above is sufficient.

Theorem 3 indicates a remarkable property of the iterated loop spaces, $\Omega^n X$, of such a space X . In more detail, notice that if $\text{map}_*(B, X)$ is weakly contractible then so is its iterated loop space $\Omega^n(\text{map}_*(B, X))$. This latter space, however, is easily seen to be homeomorphic to $\text{map}_*(B, \Omega^n X)$. Hence Theorem 3 implies that for all $n \geq 0$, the space $\text{map}_*(B, \Omega^n X)$ is weakly contractible, or equivalently, for all $n \geq 0$, every map from B to $\Omega^n X$ is null homotopic.

To begin the proof, let X and p satisfy the hypothesis of Theorem 1. Without loss of generality, we may assume that X has been localized at p . Notice that the conditions on X do not rule out the possibility that some of the groups $\pi_n X$ may contain rational vector spaces.

Our first goal is to establish that for infinitely many n , the mod p homotopy groups $\pi_n(X; \mathbb{Z}/p) \neq 0$. Recall that these groups are defined for $n \geq 2$, as

$$\pi_n(X; \mathbb{Z}/p) = \pi_0(\text{map}_*(S^{n-1} \cup_p e^n, X)).$$

They are related to the ordinary homotopy groups of X by a short exact sequence

$$0 \rightarrow \pi_n X \otimes \mathbb{Z}/p \rightarrow \pi_n(X; \mathbb{Z}/p) \rightarrow \text{Tor}(\pi_{n-1} X, \mathbb{Z}/p) \rightarrow 0.$$

For more details, see [2].

Suppose that at most a finite number of the mod p homotopy groups of X are nontrivial. Then by condition (i) and the mod p Hurewicz theorem we can choose a largest integer, say m , such that $\pi_m(X; \mathbb{Z}/p) \neq 0$.

What does this supposition imply about the ordinary homotopy groups of X ? By the universal coefficient sequence, mentioned earlier, it follows that there are just two possibilities; either

Case 1. $\pi_m X \otimes \mathbb{Z}/p \neq 0$, or

Case 2. $\pi_m X \otimes \mathbb{Z}/p = 0$ and $\text{Tor}(\pi_{m-1} X, \mathbb{Z}/p) \neq 0$

Moreover in both cases, if $\pi = \pi_n X$, then $\pi \otimes \mathbb{Z}/p = 0$ if $n > m$ and $\text{Tor}(\pi, \mathbb{Z}/p) = 0$ if $n \geq m$.

The second case is the easier to handle. In it we see that \mathbb{Z}/p is a subgroup of $\pi_{m-1} X = \pi_1 \Omega^{m-2} X$. Hence there is an essential map

$$f_1: K(\mathbb{Z}/p, 1) \rightarrow K(\pi_{m-1} X, 1)$$

Consider the obstructions to lifting this map up the Postnikov tower of $\Omega^{m-2}X$ to a map

$$f_\infty: K(\mathbb{Z}/p, 1) \rightarrow \Omega^{m-2}X$$

These obstructions take values in $\tilde{H}^*(K(\mathbb{Z}/p, 1); \pi)$ where $\pi = \pi_n X$ and $n > m$. By the universal coefficient theorem for cohomology [4, page 246], these obstruction groups are trivial since $\pi \otimes \mathbb{Z}/p = \text{Tor}(\pi, \mathbb{Z}/p) = 0$.

Hence Case 2 implies the existence of the essential map, f_∞ , which in turn contradicts Theorem 3. That leaves us with Case 1. In it, we see that $\mathbb{Z}_{(p)}$ is a subgroup of $\pi_m X = \pi_2 \Omega^{m-2}X$. More precisely we see that there is a monomorphism

$$g: \mathbb{Z}_{(p)} \rightarrow \pi_m X$$

which, when tensored with \mathbb{Z}/p , is still injective. This, in turn, implies that the following composition

$$g_2: K(\mathbb{Z}/p, 1) \rightarrow K(\mathbb{Z}_{(p)}, 2) \rightarrow K(\pi_m X, 2)$$

is essential. Here the first map represents a generator of $H^2(K(\mathbb{Z}/p, 1); \mathbb{Z}_{(p)}) = \mathbb{Z}/p$, and the second map is determined by g .

Let $\Omega^{m-2}X\langle 1 \rangle$ denote the 1-connective cover of $\Omega^{m-2}X$. The map g_2 can be taken to be a map into the first stage of the Postnikov tower for this cover. The obstructions to lifting g_2 up to a map

$$g_\infty: K(\mathbb{Z}/p, 1) \rightarrow \Omega^{m-2}X\langle 1 \rangle,$$

are zero for the same reasons as before. Thus g_∞ exists and is essential. The composition of g_∞ with the covering projection back into $\Omega^{m-2}X$ would likewise be essential. Once again we have reached a contradiction of Theorem 3. We therefore conclude that $\pi_n(X; \mathbb{Z}/p) \neq 0$ for infinitely many n . Notice that Theorem 2 is an immediate consequence of this fact.

To complete the proof of Theorem 1, suppose that $\text{Tor}(\pi_n X, \mathbb{Z}/p) \neq 0$ for at most a finite number of n . Then we may choose $m > 0$ large enough so that

- (i) $\text{Tor}(\pi_q \Omega^m X, \mathbb{Z}/p) = 0$ for all $q > 0$, and
- (ii) $\pi_2 \Omega^m X \otimes \mathbb{Z}/p \neq 0$.

These conditions on $\pi_2\Omega^m X$, in particular, are the same as those in the case just considered. Hence, as before there is a commutative diagram of essential maps

$$\begin{array}{ccc} K(\mathbb{Z}_{(p)}, 2) & \xrightarrow{h} & K(\pi_2\Omega^m X, 2) \\ & \nwarrow \quad \nearrow j & \\ & K(\mathbb{Z}/p, 1) & \end{array}$$

This time, however, we will consider the lifting problem for h , rather than working directly with map j .

We want to lift h up through the Postnikov tower for $\Omega^m X\langle 1 \rangle$. At the n -th stage this involves the diagram

$$\begin{array}{ccccc} & & E_{n+1} & & \\ & \nearrow h_{n+1} & \downarrow & & \\ K(\mathbb{Z}_{(p)}, 2) & \xrightarrow{h_n} & E_n & \xrightarrow{k} & K(\pi', q) \end{array}$$

where h_n is some lift of h . As usual, the next lift, h_{n+1} , exists if and only if the composition kh_n is trivial. With this in mind, note that under rationalization the k -invariant (and hence kh_n) is taken to zero. This follows because $\Omega^m X\langle 1 \rangle$ is an H -space. On the other hand, since π' is torsion-free, a simple calculation shows that

$$H^*(K(\mathbb{Z}_{(p)}, 2), \pi') \rightarrow H^*(K(\mathbb{Z}_{(p)}, 2), \pi' \otimes Q)$$

is injective. We conclude that kh_n must therefore represent the zero class in the first group. Thus kh_n is null homotopic and there is a solution, h_{n+1} , to the lifting problem.

In summary, the map h has been shown to lift to a map into $\Omega^m X\langle 1 \rangle$. Composing this lift with maps previously considered we obtain an essential map $K(\mathbb{Z}/p, 1) \rightarrow \Omega^m X$. This third and final contradiction of Theorem 3, completes the proof of Theorem 1.

We thank Dan Burghilea and Clarence Wilkerson for useful comments on this paper.

REFERENCES

- [1] H. MILLER, *The Sullivan conjecture*, Bull. AMS 9, 1983, 75–78.
- [2] J. A. NEISENDORFER, *Primary Homotopy Theory*, Memoirs AMS 232, 1980.
- [3] J.-P. SERRE, *Cohomologie modulo 2 des complexes d'Eilenberg–MacLane*, Comm. Math. Helv. 27 (1953), 198–232.

- [4] E. SPANIER, *Algebraic Topology*, McGraw-Hill, 1966.
- [5] Y. UMEDA, *A remark on a theorem of J.-P. Serre*, Proc. Japan Acad. 35 (1959), 563–566.

Wayne State University
Detroit, Michigan

Ohio State University
Columbus, Ohio

Received September 8, 1983