

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 60 (1985)

Artikel: The Euler and Pontrjagin numbers of an n -manifold in ... n .
Autor: Webster, S.M.
DOI: <https://doi.org/10.5169/seals-46309>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Euler and Pontrjagin numbers of an n -manifold in \mathbb{C}^n

S. M. WEBSTER⁽¹⁾

Introduction

According to a theorem of H. Whitney, every smooth n -dimensional manifold M^n can be smoothly embedded in the Euclidean space \mathbb{R}^{2n} . Viewing \mathbb{R}^{2n} as \mathbb{C}^n , one may ask for embeddings which have nice properties relative to the complex structure. The simplest properties relate to complex tangents. If M has no complex tangents, the embedding is said to be totally real. In general there are global obstructions to finding totally real embeddings. For example, if M is compact, orientable and totally real, then its Euler number and Pontrjagin classes must vanish, a result due to R. Wells [11].

In this paper we shall give an explicit formula for the Euler number of a compact real n -manifold M suitably immersed in a complex n -manifold. (The requirements on M hold generically if $n \leq 5$.) We shall also give a formula for the Pontrjagin number of a compact, orientable M^4 generically immersed in \mathbb{C}^4 . We must assume that M has only one-dimensional complex tangents which are non-degenerate in a certain sense, and occur along a smooth, compact, codimension two submanifold $N \subset M$. There is a smooth invariant function γ on N , $0 \leq \gamma \leq +\infty$. In section 5 we derive a relation among the Euler numbers $\chi(M)$, $e = \chi[\gamma < \frac{1}{2}]$, $h = (-1)^n \chi[\gamma > \frac{1}{2}]$, $\chi(M^\perp)$ (normal bundle), and the parabolic index p , which is described in section 1. As a special case we have the following.

THEOREM (0.1). *Let the compact, orientable n -manifold M be embedded in \mathbb{C}^n as just described. Then its Euler number satisfies*

$$\chi(M) = e - h + p. \quad (0.1)$$

When $n = 2$ we have $p = 0$, since there are no parabolic points. In this case the

¹ Partially supported by N.S.F. grant No. MCS-8300245.

theorem is due to E. Bishop [1], who reduced it to a theorem of Chern and Spanier [3]. Our method of proof is different, being based on the Poincaré–Hopf formula for the Euler number. If M is totally real, (0.1) implies $\chi(M) = 0$. For a direct proof see [11]. If the 2-sphere is embedded in \mathbb{C}^2 as above, then it must have at least 2 elliptic points by (0.1). Elliptic points are of interest since they contribute to the local hull of holomorphy of M . They have been studied most recently in [6], [7], and [10]. It is not known whether a 4-sphere generically embedded in \mathbb{C}^4 must have an elliptic point. If not then (0.1) gives $h = -2$, which puts some restriction on the topology of $N = N_h$.

In some cases the Pontrjagin numbers give more information about the complex-tangent structure. Let M be immersed in \mathbb{C}^n , and J denote the real operator corresponding to multiplication by $\sqrt{-1}$. Then $H_m = T_m(M) \cap JT_m(M)$ for $m \in N$, defines a complex line bundle H over N . N inherits a natural orientation from M (section 1).

THEOREM (0.2). *Suppose the compact, orientable 4-manifold M is generically immersed in \mathbb{C}^4 . Then its Pontrjagin number satisfies*

$$p_1(M) = \chi(H). \quad (0.2)$$

This is proved in section 3, where we also show that $\chi(H) = 0$ if M has no parabolic points. Thus if $p_1(M) \neq 0$, then M has a parabolic point, hence nearby elliptic and hyperbolic points. A generic embedding of \mathbb{CP}_2 in \mathbb{C}^4 therefore has a non-trivial hull of holomorphy.

In [8] H. F. Lai has given general formulas for the Euler and Pontrjagin classes of a wider class of submanifolds of \mathbb{C}^n . The relation of his work to the present paper is not clear. His formulas do not yield (0.1) or (0.2). In section 1 we describe the local properties of $M^n \subset \mathbb{C}^n$ near a complex tangent. In section 2 we study the real Grassmannian and give a transversality argument. Section 3 contains a general result about the intersection properties of Schubert varieties needed for Theorem (0.2). Section 4 is devoted to deriving suitable local equations for M near a complex tangent.

1. Complex tangents and the parabolic index

Let M^n be a smooth real n -manifold immersed in the complex n -manifold \tilde{M} . In a local holomorphic coordinate system $z = x + iy = (z^1, \dots, z^n)$, M is given by

$$\begin{aligned} M: R = (r^1, \dots, r^n) = 0, \quad R = \bar{R}, \quad dr^1 \wedge \dots \wedge dr^n \neq 0, \\ \partial r^1 \wedge \dots \wedge \partial r^n = B dz^1 \wedge \dots \wedge dz^n. \end{aligned} \quad (1.1)$$

Under a change of holomorphic coordinates $z \rightarrow z'$ and defining function $R \rightarrow R'$, the factor B changes by

$$B \rightarrow B' = \frac{\partial R'}{\partial R} \left(\frac{\partial z'}{\partial z} \right)^{-1} B. \quad (1.2)$$

Let F denote the normal bundle of M in \tilde{M} , and F^* its dual. Also, let K denote the canonical line bundle of \tilde{M} , and L and L^* the real line bundles $\Lambda^n F$ and $\Lambda^n F^*$, respectively. Then (1.2) says that the collection $\{B, B', \dots\}$, which we denote simply by B , defines a section of the complex line bundle $K \otimes L^{*-1} = K \otimes L$ over M . In particular, the set

$$N = \{m \in M : B(m) = 0\} \quad (1.3)$$

is well defined and is precisely the set of points m at which M has a (non-trivial) complex tangent space H_m . If J_m denotes the real linear operator on the real tangent space $T_m \tilde{M}$ corresponding to multiplication by $\sqrt{-1}$, then $H_m = T_m \cap J T_m$, where $T = T(M)$ is the real tangent bundle of M .

We assume that $\dim_{\mathbb{C}} H_m = 1$, so that $H_m \otimes \mathbb{C} = H'_m \oplus H''_m$, $H''_m = \bar{H}'_m$, and H'_m is spanned by

$$X = \sum \xi^i \partial / \partial z^i, \quad XR = \sum \xi^i (\partial R / \partial z^i)|_m = 0. \quad (1.4)$$

We further assume that the complex tangent H_m is non-degenerate, in that

$$\text{either } XB \neq 0 \quad \text{or} \quad \bar{X}B \neq 0. \quad (1.5)$$

Since X is determined up to $X \rightarrow cX$, c a non-zero complex constant, it follows that

$$\gamma(m) = \frac{1}{2} |XB / \bar{X}B| \in [0, \infty] \quad (1.6)$$

is a well defined biholomorphic invariant of M at m . It was first found by Bishop [1] in the following form.

We suppose that m is the origin of a coordinate system $(z_1, z^\alpha, 2 \leq \alpha \leq n-1, z_n)$ in which T_m is the (z_1, x^α) -space. M is given locally as the graph $R = 0$,

$$\begin{aligned} r &\equiv -z_n + F(z_1, x), & F &= q + x \cdot O(1) + O(2), \\ r^\alpha &\equiv -y^\alpha + f^\alpha(z_1, x), & f^\alpha &= \bar{f}^\alpha = O(2), & 2 \leq \alpha \leq n-1, \\ x &= (x^\alpha), & q &= az_1^2 + bz_1\bar{z}_1 + c\bar{z}_1^2. \end{aligned} \quad (1.7)$$

Here and elsewhere $O(k)$ indicates a term which vanishes to order k at the origin. Then, up to a constant, $X = \partial/\partial z_1$, and

$$B = \frac{\partial(r, \bar{r}, r^\alpha)}{\partial(z_1, z_n, z^\beta)} = (1/2)^{n-2} \partial \bar{F} / \partial z_1 + O(2), \quad XB = 2c, \quad \bar{X}B = \bar{b},$$

$$\gamma(m) = |c/b|. \quad (1.8)$$

We also introduce

$$\Delta = |XB|^2 - |\bar{X}B|^2. \quad (1.9)$$

N is partitioned into

$$N_e = [\gamma < \tfrac{1}{2}], \quad N_p = [\gamma = \tfrac{1}{2}], \quad N_h = [\gamma > \tfrac{1}{2}], \quad (1.10)$$

the sets of elliptic, parabolic, and hyperbolic points. These sets correspond to $\Delta < 0$, $\Delta = 0$, $\Delta > 0$, respectively.

Next we examine more closely a parabolic point m . Since the determinant (1.9) vanishes, (1.5) implies that the system

$$aXB + b\bar{X}B = 0, \quad aX\bar{B} + b\bar{X}\bar{B} = 0,$$

has a non-trivial solution (a, b) unique up to $(a, b) \rightarrow (a', b') = (\mu a, \mu b)$. Complex conjugation of these equations shows that $\bar{b} = \lambda a$, $\bar{a} = \lambda b$, for some λ , $|\lambda| = 1$. The factor μ can be adjusted so that $b' = \bar{a}'$. Hence, there is an a , unique up to $a \rightarrow \rho a$, $\rho \in \mathbb{R}$, for which

$$YB = 0, \quad Y = aX + \bar{a}\bar{X}. \quad (1.11)$$

It's easily to be seen that the changes $X \rightarrow cX$ and (1.2) can change Y by at most a real factor. Thus Y spans an intrinsic real line $l_m \subset H_m$, which we call the *parabolic line* at m .

Now we assume that

$$dB \wedge d\bar{B} \neq 0 \quad \text{on} \quad B = 0, \quad (1.12)$$

so that N is a real submanifold of M of codimension 2. The conormal bundle S^* of N in M is spanned by the coframes dB along N , and hence is the restriction of $(K \otimes L)^*$ to N . So the *normal bundle* S of N in M is the complex line bundle $K \otimes L$ restricted to N . The non-degeneracy condition (1.5) precludes H_m being contained in $T_m(N)$. It follows from (1.11) and (1.9) that m is a parabolic point

precisely when $T_m N \cap H_m = l_m$ is one dimensional. We denote by φ the composite vector bundle mapping

$$\varphi : H \hookrightarrow T(M)|_N \rightarrow (T(M)|_N)/T(N) \equiv S. \quad (1.13)$$

Since dB is local coframe for S and X is a $(1, 0)$ -frame for H_m , consideration of the sign of (1.9) shows that φ_m is orientation reversing if m is elliptic and orientation preserving if m is hyperbolic. It is singular of real rank one if m is parabolic.

We assume still further that $d\gamma \neq 0$ when $\gamma = \frac{1}{2}$, so that N_p is a smooth $(n-3)$ -dimensional manifold. We may now proceed to define the parabolic index. For each $m \in N_p$ we have two lines in $T_m(N)$, the parabolic line l_m and the line k_m determined by the normal vector $\nabla\gamma$ (gradient relative to any convenient metric on N). This gives us two sections l, k of the projective bundle $P \rightarrow N_p$, which has as fiber over $m \in N_p$ all lines through the origin in $T_m(N)$.

We next describe an orientation on the open subset of P consisting of lines not tangent to N_p . Let x^α , $3 \leq \alpha \leq n-1$, be local coordinates on N_p . Let y be a local defining function for N_p : $y=0$, with $\partial\gamma/\partial y > 0$. Then (x, y) are local coordinates on N , and any line $L \in P$, not tangent to N_p , is spanned by a unique vector

$$\frac{\partial}{\partial y} + \sum_{\alpha=3}^{n-1} w^\alpha \frac{\partial}{\partial x^\alpha} \in L. \quad (1.14)$$

Thus (x, w) are coordinates for L , and

$$\Omega_P = dx^3 \wedge \cdots \wedge dx^{n-1} \wedge dw^3 \wedge \cdots \wedge dw^{n-1} \quad (1.15)$$

defines a local volume form on the $(2n-6)$ -dimensional manifold P . If (\tilde{x}, \tilde{y}) is another such coordinate system, then one easily sees that

$$\tilde{\Omega}_P = (\det \partial \tilde{x} / \partial x)^2 (\partial \tilde{y} / \partial y)^{-n+3} \Omega_P.$$

Since $\partial \tilde{y} / \partial y > 0$, we have a well defined orientation.

The *parabolic index* is defined as the intersection number of $l(N_p)$ and $k(N_p)$ relative to this orientation. This is possible since k_m is never tangent to N_p . More precisely, we take a slight perturbation of k , if necessary, so that $k(N_p)$ and $l(N_p)$ intersect transversely in P at a finite number of points. At a point $m \in N_p$ where $l_m = k_m$, we choose local coordinates as in the previous paragraph. Then $\partial l / \partial x^\alpha$, $\partial k / \partial x^\alpha$, $3 \leq \alpha \leq n-1$, give frames for $l(N_p)$ and $k(N_p)$ at m . The intersection index

at m is given by the sign

$$\text{ind}_{P,m}(l, k) = \text{sgn } \Omega_P(\partial l / \partial x, \partial k / \partial x). \quad (1.16)$$

This is well defined since a change of orientation of the local coordinates x changes the orientations on *both* $l(N_p)$ and $k(N_p)$, so that (1.16) remains unchanged. The parabolic index is

$$p = \sum \{\text{ind}_{P,m}(l, k) : m \in N_p, l_m = k_m\}. \quad (1.17)$$

2. The Grassmann manifold and transversality

In this section we consider an n -manifold M immersed in \mathbb{C}^n . Its Gauss map g associates to each $m \in M$ the real tangent plane $T_m(M)$. It is a smooth mapping of M into $Gr(n; n) \equiv Gr$, the n^2 -dimensional Grassmann manifold of real n -planes through the origin of $\mathbb{C}^n \cong \mathbb{R}^{2n}$. We define

$$\begin{aligned} C_k &= \{V \in Gr : \dim_{\mathbb{C}} V \cap JV = k\}, \quad 0 \leq k \leq n/2; \\ C &= C_1 \cup \cdots \cup C_{[n/2]}. \end{aligned} \quad (2.1)$$

C_0 is the dense open subset of totally real planes, and $Gr = C_0 \cup \cdots \cup C_{[n/2]}$ is a disjoint union. For each k , $0 < k \leq n/2$, C_k fibers over the complex Grassmannian $Gc(k, n-k)$ of complex k -planes in \mathbb{C}^n . In fact, for V in C_k , $V \cap JV \cong \mathbb{C}^k$ and $V \cap (V \cap JV)^\perp$ is a totally real $(n-2k)$ -plane in \mathbb{C}^{n-k} . So the fiber is an open subset of $Gr(n-2k; n-k)$, which has dimension $n(n-2k)$, while the base has real dimension $2k(n-k)$. Thus C_k is a real submanifold of codimension $2k^2$. If $n = 2m$ is even, $C_m = Gc(m; n)$; while if $n = 2m+1$, C_m is a bundle over $Gc(m; n)$ with fiber the real projective space $\mathbb{RP}(2m+1)$.

We repeat some of the constructions of section one in this “universal” setting. For $V \in Gr$ there are two natural vector spaces; V itself and $F_V = \mathbb{C}^n / V$. We also have the two real line bundles $L = \bigwedge^n F$, $L^* = \bigwedge^n F^*$, and the complex line bundle $K = \bigwedge^n (Gr \times \mathbb{C}^{n*}) \cong Gr \times \mathbb{C}$. We refer back to (1.1) where now the dr^i are n independent real linear forms on \mathbb{C}^n annihilating a fixed V in Gr , and the dz^i are a basis of complex linear functions on \mathbb{C}^n . As before it follows that B is a section of $K \otimes L$, having as zero set precisely C . We restrict this bundle to C_1 , where it is identified with the normal bundle S of C_1 in Gr . Its dual bundle has local frames dB restricted to C_1 . Also, we have the complex line bundle $H \rightarrow C_1$, $H_V = V \cap JV$, $H \otimes \mathbb{C} = H' \oplus H''$.

In addition we have a real 4-plane bundle $E \rightarrow C_1$ defined by $E_V = \text{Hom}_{\mathbb{R}}(H_V, S_V)$. Each element of E_V may be described by an equation

$$dB = XB\theta + \bar{X}B\bar{\theta},$$

where X is a frame vector for H' and θ is its dual. If we denote by E_0 the zero section of E , which corresponds to $XB = \bar{X}B = 0$, then we have a well defined function $\gamma : (E - E_0) \rightarrow [0, \infty]$ given by (1.6). Parallel to (1.10) we have the disjoint union $E = E_0 \cup E_e \cup E_p \cup E_h$, where E_e, E_p, E_h are the sets where $\gamma < \frac{1}{2}$, $\gamma = \frac{1}{2}$, $\gamma > \frac{1}{2}$, respectively.

For M immersed in \mathbb{C}^n with at most one dimensional complex tangents, it is clear that g^*H and g^*S are the corresponding bundles of section 1. For each $m \in N$ we have

$$dg_m : H_m \cong H_{g(m)} \rightarrow T_{g(m)}Gr \rightarrow T_{g(m)}Gr / T_{g(m)}C_1 \cong S_{g(m)},$$

which defines a map $dg : N \rightarrow E$. The degenerate points form the set $dg^{-1}(E_0)$. Clearly, $\gamma \circ dg$ is the invariant (1.6).

PROPOSITION (2.1). *Let $f : M^n \rightarrow \mathbb{C}^n$ be a smooth immersion. a) A generic small perturbation of f results in an immersion with the following properties: M has no complex k -dimensional tangents if $2k^2 > n$, while if $2k^2 \leq n$, the points with such tangents form a submanifold of codimension $2k^2$. b) Suppose in addition that the immersion f has only one dimensional complex tangents which occur along the set N . After a generic small perturbation, N is a compact smooth $(n-2)$ -manifold, and the set N_0 of degenerate points is a smooth $(n-6)$ -manifold. c) Assume further that N_0 is empty. Then after a generic small perturbation the parabolic set N_p forms a compact smooth $(n-3)$ -manifold along which $d\gamma \neq 0$.*

Remark. As a consequence a generic M^n in \mathbb{C}^n has the following characteristics:

- i) $n = 2$ —isolated elliptic or hyperbolic points;
 - ii) $n = 3, 4, 5$ —at most one-dimensional non-degenerate complex tangents with N_p smooth;
 - iii) $n = 6$ —at most one-dimensional complex tangents and at most isolated degenerate points;
 - iv) $n = 8$ —at most isolated 2-dimensional complex tangents.
- Case i) is due to Hunt and Wells [5].

The main ingredient in the proof is the parametric transversality theorem (see

e.g. Hirsch [4], p. 79). Let f be any immersion of M with Gauss map $g = f_*$. Let Q denote the set of all real affine transformations $A(z) = A'(z) + a$, $A' \in Gl(2n, \mathbb{R})$, $a \in \mathbb{C}^n$. Then $G(A, z) \equiv g_{Af}(z) = A' \circ g_f(z)$ defines a smooth map

$$G : Q \times M \rightarrow Gr. \quad (2.2)$$

We claim that the mapping G is transverse to C_k for every $k \leq [n/2]$. This means that $TC_k + T\mathcal{J}mG = TGr$, at every point $G(A, z) \in C_k$, where $T\mathcal{J}mG \equiv DG(T(Q \times M))$. This will follow from $T\mathcal{J}mG = TGr$, which has nothing to do with complex tangents. We let $\mathbb{C}^n = V \oplus V^\perp$ with coordinates (x_1, x_2) , and restrict to the submanifold of Q of maps of the form $(x_1, x_2) \rightarrow (x_1, x_2 - Bx_1)$, $B \in \text{Hom}_{\mathbb{R}}(V, V^\perp)$. It is clear that DG maps the tangent space of this submanifold at $B = 0$ onto $T_V Gr$.

By the parametric transversality theorem, the set Q_k of A 's for which $z \rightarrow G(A, z)$ is transverse to C_k is residual. It follows that $Q_1 \cap \dots \cap Q_{[n/2]}$ is also residual and therefore dense. Thus there are affine (or even linear) mappings A , arbitrarily close to the identity, for which $A \circ f$ is transverse to every C_k . Part a) follows by transversality. Under the additional assumption of b) N is a compact smooth $(n-2)$ -manifold.

For parts b) and c) we must use mappings quadratic in (z, \bar{z}) : $A(z) = a + A'(z) + A''(z^2)$. We first replace f by a perturbation as in a), so that the Gaussian image $g(M)$ remains disjoint from C_k , $k > 1$, and intersects C_1 transversely. We then restrict to quadratic mappings A with A' unitary and A'' so small that this situation is preserved. We let Q denote this set of maps. For A'' small enough, perturbation by $A \in Q$ will result in a new manifold N_A close to the original N in the following sense. N_A will lie in a small tubular neighborhood $u = U\{D_m : m \in N\}$ and will intersect each normal 2-disc D_m in a unique point $\eta_A(m)$, giving a diffeomorphism $\eta_A : N \rightarrow N_A$. We consider the composite map.

$$G'(A, m) = dg_{Af} \circ \eta_A(m), \quad G' : Q \times N \rightarrow E.$$

We claim that the map G' is transverse to E_0 under the assumptions of b) and transverse to E_p under those of c). Assume $G'(A, m) \in E_0$. After an affine unitary coordinate change we may assume that $m = 0$ and that M is given as in (1.7). We then restrict G' to the submanifold of Q consisting of mappings of the special form

$$A : \begin{cases} z_1 \rightarrow z_1, z^\alpha \rightarrow z^\alpha, & 2 \leq \alpha \leq n-1 \\ z_n \rightarrow z_n + Bz_1\bar{z}_1 + C\bar{z}_1^2, \end{cases}$$

which result in $(b, c) \rightarrow (b-B, c-C)$ in (1.7). Locally $E \cong C_1 \times \mathbb{R}^4$ and $E_0 \cong C_1 \times \{0\}$. The normal space to E_0 at m is $\{m\} \times \mathbb{R}^4 \cong \{m\} \times \mathbb{C}^2$ with coordinates (b, c) . If we restrict G' further to (A, m) with A as described and $m = 0$, then it is clear that DG' at $(A, m) = (I, 0)$ maps onto the normal space. Thus G' is transverse to E_0 . A similar argument shows that G' is transverse to $E_p \cong C_1 \times \text{Cone}$. Now b) and c) follow by the parametric transversality theorem since E_0 has codimension 4 and E_p codimension one in E .

3. The Pontrjagin number of a 4-manifold in \mathbb{C}^4

By a well-known theorem [9] the Pontrjagin class $p_k(T(M))$ equals $(-1)^k c_{2k}(T(M) \otimes \mathbb{C})$, where c denotes the Chern class. For $M^n \subset \mathbb{C}^n$ with Gauss map g , $p_k(T(M)) = g^* p_k(VGr)$, where $VGr \rightarrow Gr$ is the universal bundle. $Gr \subset Gc(n, n)$ and $(VGr) \otimes \mathbb{C}$ is the restriction of the complex universal bundle $VGc \rightarrow Gc$. Thus we must consider the Chern classes $c_k(VGc)$.

We begin by recalling some facts [2] about $Gc(n, r)$, the space of all complex n -planes $Z^n \subset \mathbb{C}^{n+r}$. For $0 \leq k_1 \leq \dots \leq k_n \leq r$ and linear spaces $L_1 \subset \dots \subset L_n$, $\dim L_j = k_j + j$, the Schubert variety is defined by

$$Z(k_1, \dots, k_n) = \{Z \in Gc(n, r) \mid \dim Z \cap L_j \geq j\},$$

and has complex dimension $k_1 + \dots + k_n$. The Chern class $c_k(VGc(n, r))$ is dual to $Z_k(n, r) \equiv Z(r-1, \dots, r-1, r, \dots, r)$, where $r-1$ appears k -times. It may also be defined as

$$Z_k(n, r) = \{Z \in Gc(n, r) \mid \dim Z \cap L \geq k\}, \quad \dim L = r-1+k, \quad (3.1)$$

and decomposes into the disjoint union of

$$Z_k^0(n, r) = \{Z : \dim Z \cap L = k\},$$

and

$$Z_k^s(n, r) = \{Z : \dim Z \cap L > k\}.$$

$Z_k^0(n, r)$ is a complex manifold of complex codimension k , fibering over $Gc(k, r-1)$, and $Z_k^s(n, r) = Z(r-2, \dots, r-2, r, \dots, r)$ ($k+1(r-2)$'s) has codimension $2(k+1)$. Thus a generic compact orientable real $2k$ -manifold in $Gc(n, r)$, which we also denote by M , will be disjoint from $Z_k^s(n, r)$ and intersect $Z_k^0(n, r)$

transversely in finitely many points. $c_k(\text{VGc}(n, r))[M]$ is the sum of the intersection indices at these points.

We fix k and $L_{r-2+k} \subset L_{r-1+k}$, subscripts denoting dimension, and consider the corresponding varieties $Z_{k-1}(n, r) \supset Z_k(n, r)$, $Z_{k-1}^s(n, r) \supset Z_k^s(n, r)$. A generic M^{2k} will miss $Z_{k-1}^s(n, r)$, since it has real codimension $4k$. Thus

$$M \cap Z_{k-1}(n, r) = M \cap Z_{k-1}^0(n, r) \equiv N$$

is a smooth compact oriented 2-manifold containing the finite set $Z_k(n, r) \cap M$. For $Z \in Z_{k-1}^0(n, r)$, we set

$$A_Z = Z \cap L_{r-2+k} \quad \text{and} \quad B_Z = Z \cap A_Z^\perp,$$

so that $Z = A_Z \oplus B_Z$ is an orthogonal direct sum relative to the standard hermitian inner product on \mathbb{C}^{n+r} . We define a smooth map β by

$$\beta: Z_{k-1}^0(n, r) \rightarrow \text{Gc}(n-k+1, r+k-1), \quad \beta(Z) = B_Z. \quad (3.2)$$

For $Z \in Z_{k-1}^0(n, r)$, $Z \in Z_k^0(n, r)$ if and only if $B_Z \in Z_1(n-k+1, r+k-1)$; i.e.

$$Z_k(n, r) \cap Z_{k-1}^0(n, r) = \beta^{-1}(Z_1(n-k+1, r+k-1)).$$

LEMMA (3.1)

$$c_k(\text{VGc}(n, r))[M] = c_1(\text{VGc}(n-k+1, r+k-1))[\beta N]. \quad (3.3)$$

Proof. This comes down to comparing two intersection indices. First, we have the equality of oriented vector spaces at $m \in M \cap Z_k^0(n, r)$

$$T_m Z_k^0(n, r) \oplus T_m M = c_k T_m \text{Gc}(n, r),$$

where $c_k = \pm 1$. If S is the normal bundle of $Z_{k-1}^0(n, r)$ in $\text{Gc}(n, r)$, then its restriction to N is that of N in M , so

$$T_m \text{Gc} = T_m Z_{k-1}^0 \oplus S_m \quad \text{and} \quad T_m M = T_m N \oplus S_m.$$

It follows that

$$T_m Z_k^0(n, r) \oplus T_m N = c_k T_m Z_{k-1}^0(n, r). \quad (3.4)$$

The map β is not holomorphic since it involves the orthogonal complement A_Z^\perp . However, a slight local deformation of it is. For all $Z \in Z_{k-1}^0(n, r)$ sufficiently near m we may replace A_Z^\perp by A_m^\perp , then $\alpha(z) = Z \cap A_m^\perp$ is holomorphic and approximates β near m . Clearly, $\alpha^{-1}(Z_1^0(n-k+1, r+k-1)) = Z_k^0(n, r)$. By (3.4)

$$T_{\alpha m}(\alpha Z_k^0(n, r)) \oplus T_{\alpha m}(\alpha N) = c_k T_{\alpha m}(\alpha Z_{k-1}^0(n, r)), \quad (3.5)$$

where the orientations agree with those from $Gc(n-k+1, r+k-1)$, since α is holomorphic. If \tilde{S} is the normal bundle of $\alpha Z_{k-1}^0(n, r)$ in $Gc(n-k+1, r+k-1)$ it is also the normal bundle of $\alpha Z_k^0(n, r)$ in $Z_1^0(n-k+1, r+k-1)$. Adding $\tilde{S}_{\alpha m}$ to both sides of (3.5) gives

$$T_{\alpha m} Z_1^0(n-k+1, r+k-1) \oplus T_{\alpha m}(\alpha N) = c_k T_{\alpha m} Gc(n-k+1, r+k-1).$$

Now $\alpha m = \beta m$, and if we continuously deform α back to β , we see that $c_k = c_1$, where c_1 is the intersection index at βm entering into the right hand side of (3.3). Summing over all such m in $M \cap Z_k^0(n, r)$ gives (3.3).

Note. The same argument gives

$$c_k(VGc(n, r)[M^{2k}]) = c_{k-l}(VGc(n-l, r+l))[M \cap Z_l^0(n, r)],$$

when the intersections are nice, which is generically so when $2l+2 > k$.

We return to the study of $Gr(n; n)$. We set $\mathbb{C}^n = (W, J)$, $W \otimes \mathbb{C} \equiv W^c \equiv W' \oplus W''$, $\bar{W}'' = W' = \{w \in W^c : Jw = iw\}$. For a subspace $V \subset W$, $V^c \subset W^c$, $V^c \supset V' \oplus V''$, where $V' = V^c \cap W'$, $V'' = V^c \cap W''$. The map $V \rightarrow V^c$ embeds $Gr(n; n)$ in $Gc(n, n)$ as a totally real submanifold.

LEMMA (3.2). a) $V^c = V' \oplus V''$ if and only if $JV = V$.

b) $V' = H'$ if $H = V \cap JV$.

c) $C = Gr(n; n) \cap Z_1(n, n)$.

Proof. a) If $JV = V$, then $JV^c = V^c$, and any $w \in V^c$ is the sum $\frac{1}{2}(w - iJw) + \frac{1}{2}(w + iJw) \in V' \oplus V''$. If $V^c = V' \oplus V''$, then $JV^c = JV' + JV'' = V' \oplus V'' = V^c$; so $JV = V$. b) $V' \supset H'$ is clear since $V^c \supset H^c$. If $w \in V'$, then $Jw = iw$. So if $w = u + iv$, $u, v \in V$, then $u = Jv$, $v = -Ju$. Hence, $u, v \in H$ and b) holds. If we apply a) to H , then it follows that $V \in C$ if and only if $\dim_{\mathbb{C}} V^c \cap W' \geq 1$. So c) follows by taking $n = r$, $k = 1$, and $L_n = W'$ in (3.1).

We now turn to the proof of Theorem (0.2) of the introduction. Since M^4 is generically immersed in \mathbb{C}^4 it has the properties of Remark (ii) following Proposi-

tion (2.1). We have

$$\begin{aligned} p_1(M) &= p_1(T(M))[M] = p_1(VGr(4; 4))[gM] \\ &= -c_2(VGc(4, 4))[gM] = -c_1(VGc(3, 5))[\beta gN], \end{aligned}$$

by (3.3) with $n=r=4$, $k=2$. Now over the surface N (or rather βgN) $VGc(4, 4) = A \oplus B = H' \oplus VGc(3, 5)$ by Lemma 3.2). Thus, the total Chern class $c = 1 + c_1$ satisfies $[9] \quad c(VGc(4, 4)) = c(H')c(VGc(3, 5))$ or $c_1(VGc(4, 4)) = c_1(H') + c_1(VGc(3, 5))$. Over $Gr(4; 4)$ $VGc(4, 4) = VGr(4; 4) \otimes \mathbb{C}$, hence its first Chern class is a 2-torsion element. When pulled back to the compact orientable surface N it vanishes; thus $c_1(VGc(3, 5)) = -c_1(H') = -c(H)$. Hence, $p_1(M) = \chi(H)$, since $\chi(H) = c_1(H)[N]$.

The bundle mapping φ (1.13) can be used to get a formula for $\chi(H)$. Since $\tilde{M} = \mathbb{C}^n$, the canonical bundle K is trivial. Since M is orientable, so is its normal bundle, hence the line bundle L is trivial. It follows that $S = K \otimes L$ is trivial. φ is a bundle isomorphism over N_h and an anti-isomorphism over N_e . Therefore H is trivial over any connected component of N which does not meet N_p . If we let N^0 be the union of the components of N which meet N_p , and H^0 the restriction of H to N^0 , then $\chi(H) = \chi(H^0)$. We choose a section v of H^0 which does not vanish on $N_e^0 \cup N_p$ and has only isolated non-degenerate zeros in N_h^0 . v gives a trivialization of H over N_p ; hence, the parabolic line l gives a map $\tilde{l}: N_p \rightarrow \mathbb{RP}_1$, where N_p and \mathbb{RP}_1 have naturally induced orientations. We define the H -parabolic index p_H to be the degree of this mapping \tilde{l} . If w is a piecewise smooth section of H over N_p which spans l at each point, then $w = \mu v$, $\mu \neq 0$ and piecewise smooth. We have

$$p_H = \operatorname{Re} \frac{1}{\pi i} \int_{N_p} \frac{d\mu}{\mu}. \quad (3.6)$$

Note that w is determined up to $w \rightarrow \rho w$, with $\rho \neq 0$, real and piecewise smooth. It follows that (3.6) is not affected by this change. Also, v may be changed by $v \rightarrow \xi v$, $\xi \neq 0$ and smooth on $N_p \cup N_e^0$. Applying Stokes's theorem to $d\xi/\xi$ on N_e^0 shows that the integral in (3.6) remains unchanged. Thus p_H is well defined.

LEMMA (3.3). $\chi(H) = -p_H$.

Proof. This follows by comparing the index sums for v and $\varphi(v)$. We assume that v has been chosen so that l_m contains v_m at only a finite number of points m in N_p and that l crosses v transversely at such points. In otherwords $1 \in \mathbb{RP}_1$ is a regular value of \tilde{l} . At such a point m we choose local coordinates (x, y) on N so that $m = (0, 0)$, N_p is given by $y = 0$, N_h by $y > 0$, and N_e by $y < 0$. We let $\zeta = \xi + i\eta$

be a local fiber coordinate on H^0 relative to v and choose a local frame v' and related coordinate $\zeta' = \xi' + i\eta'$ for S near m . We may assume that $\varphi(v)$ is a positive multiple of iv' at m . Then

$$\begin{aligned} \varphi: \quad \xi' &= a\xi + b\eta, & a(0) &= b(0) = c(0) = 0, \\ \eta' &= c\xi + d\eta, & d(0) &> 0. \end{aligned}$$

$\Delta = ad - bc$, $\Delta(x, 0) = 0$, $\Delta_y(x, 0) > 0$. We let $(\xi, \eta) = (1, \lambda(x))$, $\lambda(0) = 0$, span l along N_p ; then the sign of $\lambda_x(0)$ gives the intersection index of l with respect to v . Since $\varphi(l) = 0$, we have $c + d\lambda = 0$, so $\lambda_x(0) = -c_x(0)/d(0)$. Also, $a(x, 0) = (bc/d)(x, 0)$, so $a_x(0) = 0$, and $\Delta_y(0, 0) = a_y(0)d(0)$, so $a_y(0) > 0$. Finally, $\varphi(v) = \varphi(1, 0) = (a, c)$ has index at $m = (0, 0)$ given by the sign of

$$\frac{\partial(a, c)}{\partial(x, y)}(0, 0) = -(a_y c_x)(0, 0).$$

Thus the index of $\varphi(v)$ at m is the same as the H -parabolic index at m . Since $\varphi(v)$ has the same index as v at any zero of v (in N_h), we have $\chi(S) = \chi(H) + p_H$. But $\chi(S) = 0$, since S is trivial, and the lemma follows.

Theorem (0.2) and Lemma (3.3) give

COROLLARY (3.4). *If M^4 is compact, orientable and generically immersed in \mathbb{C}^4 , then $p_1(M) = -p_H$. If $p_1(M) \neq 0$, then M must have elliptic, parabolic, and hyperbolic points.*

4. Local equations for M

To facilitate the study of M near a complex tangent, we shall simplify the presentation (1.7) by means of a local holomorphic coordinate change. In this section we prove the following.

PROPOSITION (4.1). *Suppose M has a non-degenerate one-dimensional complex tangent at a point m . Then holomorphic coordinates $z = (z_1, z^\alpha, 2 \leq \alpha \leq n-1, z_n)$ can be chosen so that $m = 0$ and M is given locally by*

$$\begin{aligned} M: \quad z_n &= F(z_1, x), & x &= (x^2, \dots, x^{n-1}), \\ y^\alpha &= f^\alpha(z_1, x), & f^\alpha &= \bar{f}^\alpha, 2 \leq \alpha \leq n-1. \end{aligned} \tag{4.1}$$

If m is an elliptic or hyperbolic point, then

$$\begin{aligned} F &= q + H, \quad q = az_1^2 + bz_1\bar{z}_1 + a\bar{z}_1^2, \quad H = O(3), \\ f^\alpha &= b^\alpha z_1\bar{z}_1 + h^\alpha, \quad h^\alpha = O(3), \end{aligned} \quad (4.2)$$

where $a \geq 0$, and b, b^α are either 0 or 1. If m is a parabolic point, then

$$\begin{aligned} F &= Q + H, \quad f^\alpha = O(4) \\ Q &= \frac{1}{2}(z_1 + \bar{z}_1)^2 + i(z_1 - \bar{z}_1)c(x), \quad c(x) = c_\beta x^\beta, \\ H &= (-i\eta(z_1 + \bar{z}_1) + \eta_\beta x^\beta)z_1\bar{z}_1 + O(4), \end{aligned} \quad (4.3)$$

where β is summed from 2 to $n-1$, c_β, η_β are real, and η is either 0 or 1. If the transversality condition $dB \wedge d\bar{B} \neq 0$ holds, then $c(x) \neq 0$. In this case the parabolic line at m , which is the y_1 -axis, is tangent to N_p if and only if $\eta = 0$.

We remark that (4.2) is already known [1], [10].

We begin with M in the form (1.7). If $b \neq 0$, we replace z_n by bz_n to make $b = 1$. By a rotation $z_1 \rightarrow \mu z_1$, $\mu\bar{\mu} = 1$, we can make $c \geq 0$. Then by a change of the form

$$\begin{aligned} z_n &\rightarrow z_n + (c - a)z_1^2 + e_\alpha z_1 z^\alpha + f_{\beta\alpha} z^\alpha z^\beta, \\ z^\alpha &\rightarrow z^\alpha + 2i(a^\alpha z_1^2 + d_\beta^\alpha z_1 z^\beta + f_{\beta\gamma}^\alpha z^\beta z^\gamma), \end{aligned} \quad (4.4)$$

we can achieve (4.2) but with

$$H = c(x)z_1 + \bar{c}(x)\bar{z}_1 + O(3), \quad c(x) = c_\beta x^\beta. \quad (4.5)$$

The b^α are either 0 or can be made 1 by $z^\alpha \rightarrow b^\alpha z^\alpha$.

We make the further change

$$z_1 \rightarrow z_1 + A(z), \quad A(z) = A_\alpha z^\alpha, \quad (4.6)$$

under which $c(x)$ in (4.5) changes by

$$\begin{aligned} c(x) &\rightarrow c(x) + 2aA(x) + b\bar{A}(x), \\ \bar{c}(y) &\rightarrow \bar{c}(x) + bA(x) + 2a\bar{A}(x). \end{aligned}$$

If $\gamma = |a/b| \neq \frac{1}{2}$, then the determinant $b^2 - 4a^2 \neq 0$, and $A(z)$ can be chosen

uniquely to make $c(x) \rightarrow 0$. If $\gamma = \frac{1}{2}$, we take $b = 1$ and $a = \frac{1}{2}$. Then (4.6) results in

$$c(x) \rightarrow c(x) + 2 \operatorname{Re} A(x),$$

which may be used to make the $c(x)$ in (4.5) purely imaginary. The $x^\alpha x^\beta$ terms introduced by (4.6) can then be removed by a transformation of the form (4.4). This gives (4.2) and the form (4.3) for the quadratic term Q .

We must investigate the third order terms in the parabolic case. Since $b = 1$, the change $z^\alpha \rightarrow z^\alpha - b^\alpha z_n$, followed by one of the type (4.4), makes $f^\alpha \equiv h^\alpha = O(3)$ in (4.1). We put

$$h^\alpha = h_0^\alpha + h_{\beta\gamma}^\alpha x^\beta x^\gamma + c_{\beta\gamma\rho}^\alpha x^\beta x^\gamma x^\rho + O(4),$$

$$h_0^\alpha = c^\alpha z_1^3 + e^\alpha z_1^2 \bar{z}_1 + \bar{e}^\alpha z_1 \bar{z}_1^2 + \bar{c}^\alpha \bar{z}_1^3,$$

$$h_\beta^\alpha = c_\beta^\alpha z_1^2 + e_\beta^\alpha z_1 \bar{z}_1 + \bar{c}_\beta^\alpha \bar{z}_1^2, e_\beta^\alpha \text{ real},$$

$$h_{\beta\gamma}^\alpha = c_{\beta\gamma}^\alpha z_1 + \bar{c}_{\beta\gamma}^\alpha \bar{z}_1, c_{\beta\gamma\rho}^\alpha \text{ real}.$$

The transformation

$$z^\alpha \rightarrow z^\alpha + 2i\{c^\alpha z_1^3 + c_\beta^\alpha z_1^2 z^\beta + c_{\beta\gamma}^\alpha z_1 z^\beta z^\gamma + \frac{1}{2}c_{\beta\gamma\rho}^\alpha z^\beta z^\gamma z^\rho\} \quad (4.7)$$

reduces h^α to the form

$$h^\alpha = (c^\alpha z_1 + \bar{c}^\alpha \bar{z}_1 + c_\beta^\alpha x^\beta) z_1 \bar{z}_1 + O(4). \quad (4.8)$$

The substitution

$$z^\alpha \rightarrow z^\alpha + 2i\{c^\alpha z_1 + \frac{1}{2}c_\beta^\alpha z^\beta\} z_n, \quad (4.9)$$

followed by another one of type (4.7) (to remove any newly introduced third order terms already removed by (4.7)) results in

$$\begin{aligned} z_n &= Q + H, & H &= O(3) \\ y^\alpha &= h^\alpha, & h^\alpha &= O(4). \end{aligned} \quad (4.10)$$

Next we consider the third order terms in h ,

$$\begin{aligned} H &= H_0 + H_\alpha x^\alpha + H_{\alpha\beta} x^\alpha x^\beta + K_{\alpha\beta\gamma} x^\alpha x^\beta x^\gamma + O(4), \\ H_0 &= K_0 z_1^3 + K_1 z_1^2 \bar{z}_1 + K_2 z_1 \bar{z}_1^2 + K_3 \bar{z}_1^3, \\ H_\alpha &= K_{\alpha 0} z_1^2 + K_{\alpha 1} z_1 \bar{z}_1 + K_{\alpha 2} \bar{z}_1^2, \\ H_{\alpha\beta} &= K_{\alpha\beta 0} z_1 + K_{\alpha\beta 1} \bar{z}_1. \end{aligned} \quad (4.11)$$

We shall simplify this by means of a transformation of the form

$$\begin{aligned}
 z_1 &\rightarrow z_1 + A(z_1, z^\alpha, z_n), & A &= A_2 + A_0 z_n, \\
 z_n &\rightarrow z_n + B(z_1, z^\alpha, z_n), & B &= B_3 + B_1 z_n, \\
 A_2 &= A_{20} z_1^2 + A_{2\alpha} z_1 z^\alpha + A_{2\alpha\beta} z^\alpha z^\beta, & A_0 &= \text{const.}, \\
 B_3 &= B_{30} z_1^3 + B_{3\alpha} z_1^2 z^\alpha + B_{3\alpha\beta} z_1 z^\alpha z^\beta + B_{3\alpha\beta\gamma} z^\alpha z^\beta z^\gamma, \\
 B_1 &= B_{10} z_1 + B_{1\alpha} z^\alpha.
 \end{aligned} \tag{4.12}$$

This will not alter any of the previous normalizations. Note that

$$Q(z_1 + A, x) = Q(z_1, x) + (z_1' + \bar{z}_1)(A + \bar{A}) + Q(A, x).$$

Therefore, when we substitute (4.12) into (4.10), we get

$$H \rightarrow H + (z_1 + \bar{z}_1)(A + \bar{A}) - B + i(A - \bar{A})c(x) + \frac{1}{2}(A + \bar{A})^2, \tag{4.13}$$

in which we must make the substitution (4.10). We shall simplify the terms of H in order of increasing degree in x^α . This allows us to ignore the term $i(A - \bar{A})c(x)$, and hence $Q(A, x)$, since $(A + \bar{A})^2$ is of fourth order.

In simplifying H_0 we ignore terms in x^α and $z^\alpha = x^\alpha + O(4)$, so that

$$H_0 \rightarrow H_0 + (z_1 + \bar{z}_1)(A_2 + \bar{A}_2 + (A_0 + \bar{A}_0)Q) - B_3 - B_1 Q,$$

with $A_2 \equiv A_{20} z_1^2$, $B_3 = B_{30} z_1^3$, $B_1 \equiv B_{10} z_1$, $Q \equiv \frac{1}{2}(z_1 + \bar{z}_1)^2$. Comparison of coefficients shows that

$$\begin{aligned}
 K_0 &\rightarrow K_0 + A_{20} + \frac{1}{2}(A_0 + \bar{A}_0) - B_{30} - \frac{1}{2}B_{10}, \\
 K_1 &\rightarrow K_1 + A_{20} + \frac{3}{2}(A_0 + \bar{A}_0) - B_{10}, \\
 K_2 &\rightarrow K_2 + \bar{A}_{20} + \frac{3}{2}(A_0 + \bar{A}_0) - \frac{1}{2}B_{10}, \\
 K_3 &\rightarrow K_3 + \bar{A}_{20} + \frac{1}{2}(A_0 + \bar{A}_0).
 \end{aligned}$$

By proper choice of A_{20} and B_{30} we can realize $K_0 = K_3 = 0$, after which $A_{20} = -\text{Re } A_0$, $B_{30} = -\frac{1}{2}B_{10}$. Then $K_1 - K_2 \rightarrow K_1 - K_2 - \frac{1}{2}B_{10}$, so that we can make $K_1 = K_2$, and restrict to $B_{10} = 0$. This leaves the change $K_1 \rightarrow K_1 + 2 \text{Re } A_0$, by which we make $K_1 = -i\eta$, purely imaginary.

To simplify $H_\alpha x^\alpha$ in (4.11), we set $A_{20} = A_0 = B_{30} = B_{10} = 0$ in (4.12) and work mod $x^\alpha x^\beta$, $z^\alpha z^\beta$. With $A_0 = A_{20} = 0$, $i(A - \bar{A})c(x) \equiv 0$, mod $x^\alpha x^\beta$, so

$$H_\alpha x^\alpha \rightarrow H_\alpha x^\alpha + (z_1 + \bar{z}_1)(A + \bar{A}) - B_3 - B_1 Q,$$

with $A \equiv A_{2\alpha} z_1 z^\alpha$, $B_3 \equiv B_{3\alpha} z_1^2 z^\alpha$, $B_1 \equiv B_{1\alpha} z_1^\alpha$, and $Q \equiv \frac{1}{2}(z_1 + \bar{z}_1)^2$. Comparison of coefficients gives

$$K_{\alpha 0} \rightarrow K_{\alpha 0} + A_{2\alpha} - B_{3\alpha} - \frac{1}{2}B_{1\alpha},$$

$$K_{\alpha 1} \rightarrow K_{\alpha 1} + A_{2\alpha} + \bar{A}_{2\alpha} - B_{1\alpha},$$

$$K_{\alpha 2} \rightarrow K_{\alpha 2} + \bar{A}_{2\alpha} - \frac{1}{2}B_{1\alpha}.$$

So we normalize to $K_{\alpha 0} = K_{\alpha 2} = 0$ and restrict to $A_{2\alpha} = B_{3\alpha} + \frac{1}{2}B_{1\alpha} = \frac{1}{2}\bar{B}_{1\alpha}$. It follows that $K_{\alpha 1} \rightarrow K_{\alpha 1} + \frac{1}{2}(\bar{B}_{1\alpha} - B_{1\alpha})$, so that we can make $K_{\alpha 1} = \eta_\alpha = \bar{\eta}_\alpha$, real.

Now we further restrict to $A_{2\alpha} = B_{3\alpha} = B_{1\alpha} = 0$ in (4.12) and work mod $x^\alpha x^\beta x^\gamma$, $z^\alpha z^\beta z^\gamma$. Again $i(A - \bar{A})c(x)$ can be ignored in (4.13). We have

$$H_{\alpha\beta} x^\alpha x^\beta \rightarrow H_{\alpha\beta} x^\alpha x^\beta + (z_1 + \bar{z}_1)(A + \bar{A}) - B_3,$$

where $A \equiv A_{2\alpha\beta} z^\alpha x^\beta$, $B_3 \equiv B_{3\alpha\beta} z_1 z^\alpha z^\beta$. This results in the change

$$K_{\alpha\beta 0} \rightarrow K_{\alpha\beta 0} + A_{2\alpha\beta} - B_{3\alpha\beta},$$

$$K_{\alpha\beta 1} \rightarrow K_{\alpha\beta 1} + \bar{A}_{2\alpha\beta}.$$

It's clear that we can make $K_{\alpha\beta 0} = K_{\alpha\beta 1} = 0$. Finally, we remove the term $K_{\alpha\beta\gamma} x^\alpha x^\beta x^\gamma$ by a transformation

$$z_n \rightarrow z_n + B_{3\alpha\beta\gamma} z^\alpha z^\beta z^\gamma.$$

This achieves the form (4.3). If $\eta \neq 0$, a dilation $(z_1, z^\alpha, z_n) \rightarrow (\lambda z_1, \lambda z^\alpha, \lambda^2 z_n)$ with λ real results in $\eta \rightarrow \lambda\eta$, so we can make $\eta = 1$.

At a parabolic point (1.8) and (4.3) give

$$B = (i/2)^{n-2} (z_1 + \bar{z}_1 + ic(x) + i\eta(\bar{z}_1^2 + 2z_1\bar{z}_1) + \eta_\beta x^\beta \bar{z}_1) + O(3). \quad (4.14)$$

It follows that $dB \wedge d\bar{B} = 4^{1-n} ic(dx) \wedge dx_1 + O(1)$, so that $c(x) \neq 0$ if the transversality condition holds. We make a linear change in the coordinates (x^2, \dots, x^{n-1}) so that $c(x) = x^2$, then N has the local equations

$$\begin{aligned} x_1 &= O(3), \\ x^2 &= -\eta y_1^2 + \sum_{\beta=3}^{n-1} \eta_\beta x^\beta y_1 + O(3). \end{aligned} \quad (4.15)$$

The conditions $Xr = X\bar{r} = Xr^\alpha = 0$, which determine X give

$$X = \frac{\partial}{\partial z_1} + (Q + H)_{z_1} \frac{\partial}{\partial z_n} + O(3). \quad (4.16)$$

Also,

$$\begin{aligned} XB &= B_{z_1} + O(3) = 1 + \eta y_1 + O(2), \\ \bar{X}B &= B_{\bar{z}_1} + O(3) = 1 + \sum_{\beta=3}^{n-1} \eta_\beta x^\beta + O(2). \end{aligned} \quad (4.17)$$

The condition (1.11) gives $a(0) + \bar{a}(0) = 0$, so we may take $a = a' + i$, a' real, $a'(0) = 0$. Then

$$YB = a'(2 + \eta y_1 + \eta_\beta x^\beta) + i(\eta y_1 - \eta_\beta x^\beta) + O(2),$$

so that $a' = O(2)$. Thus, in coordinates $(y_1, x^2, \dots, x^{n-1})$

$$Y = \partial/\partial y_1 + O(2). \quad (4.18)$$

From (4.17) and (1.9)

$$\Delta = 2\eta y_1 - 2 \sum_{\beta=3}^{n-1} \eta_\beta x^\beta + O(2); \quad (4.19)$$

so that $Y[\Delta] = 2\eta + O(1)$. It follows that Y is tangent to $N_p : \Delta = 0$ if and only if $\eta = 0$. If $\eta = 1$, then $Y[\Delta] > 0$ implies that Y points toward N_h .

5. A formula for the Euler number

To derive our formula we shall make use of the Poincaré–Hopf theorem characterizing the Euler number $\chi(M)$ as the sum of the indices of the zeros of a vector field tangent to M . This does not require M to be orientable and is applicable to compact manifolds with boundary, provided the vector field points outward along the boundary. For M^n immersed in the complex n -manifold \tilde{M} with normal bundle F , $\chi(F)$ denotes the sum of the indices of the zeros of a suitable section of F . The index at an isolated zero $m \in M$ is well defined since $T_m \tilde{M} = T_m M \oplus F_m$ as oriented vector spaces locally. A reversal of the local orientation of M near m results in a reversal of that of F as well as of TM .

In this section we prove the following, which does not require M to be orientable.

THEOREM (5.1). *Suppose that the compact n -dimensional manifold M is immersed in the complex n -dimensional manifold \tilde{M} with at most nondegenerate, one-dimensional complex tangents as in section 1 and Proposition (2.1c). Then*

$$\chi(M) = \varepsilon_n \chi(F) + e - h + p, \quad \varepsilon_n = (-1)^{(n-1)n/2}, \quad (5.1)$$

where $e = \chi(N_e)$, $h = (-1)^n \chi(N_h)$, and p is the parabolic index.

If M is also orientable and embedded in \mathbb{C}^n , then a theorem of Whitney (see [4] or [9]) asserts that $\chi(F) = 0$. Theorem (0.1) follows immediately from this. As mentioned after Proposition (2.1) the assumptions of Theorem (5.1) are generic if $n \leq 5$. The remainder of this section is devoted to the proof of Theorem (5.1).

We choose some convenient hermitian metric on \tilde{M} and denote by $\pi_m : T_m \tilde{M} \rightarrow F_m$, the orthogonal projection onto F_m along $T_m \equiv T_m M$. Then $\pi_m \circ J_m$ gives a linear mapping from T_m to F_m , which will be a linear isomorphism if m is a totally real point of M . If v is a vector field tangent to M , then πJv is a section of F . The idea of the proof is to relate the index sum of πJv to that of v for a suitable choice of v .

About any particular m in M we choose holomorphic coordinates $z = x + iy$ for \tilde{M} centered at m . The orientation of \tilde{M} is given by the local form

$$\tilde{\Omega} = \prod_{\alpha=1}^n \left(\frac{i}{2} dz^\alpha \wedge d\bar{z}^\alpha \right) = \varepsilon_n dx^1 \wedge \cdots \wedge dx^n \wedge dy^1 \wedge \cdots \wedge dy^n, \quad (5.2)$$

and the operator J is identified with $(x, y) \rightarrow (-y, x)$. Suppose m is a totally real point of M . Then the coordinates may be chosen so that T_m is the x -space and F_m is the y -space, which by (5.2) have the orientations

$$\Omega_T = dx^1 \wedge \cdots \wedge dx^n, \quad \Omega_F = \varepsilon_n dy^1 \wedge \cdots \wedge dy^n.$$

Since π is smoothly deformable to $(x, y) \rightarrow (0, y)$ and $\pi \circ J$ to $(x, 0) \rightarrow (0, x)$, we have

$$(\pi \circ J)^* \Omega_F = c \Omega_T, \quad \text{sgn } c = \varepsilon_n.$$

It follows that the effect of πJ on the index of a vector field v with isolated zero at

m is

$$\text{ind}_{F,m}(\pi Jv) = \varepsilon_n \text{ind}_{M,m}(v),$$

so that

$$\sum_{m \notin N} \text{ind}_{F,m}(\pi Jv) = \varepsilon_n \chi(M). \quad (5.3)$$

This proves (5.1) if M is totally real.

In the general case we start with a smooth vector field v_0 tangent to N with the following properties. It is to have only finitely many zeros m_j , $1 \leq j \leq l$, which are non-degenerate and lie in $N_e \cup N_h$, and is to be transverse to N_p and point toward N_h along N_p . Furthermore, the line field k along N_p spanned by v_0 is to satisfy $k_m = l_m$ for only finitely many $m \in N_p$, and at such m this intersection is transverse in the space P (see (1.15)). We find disjoint neighborhoods U_j of m_j in $N - N_p$ and smooth sections v_j of H , compactly supported in U_j , with $v_j(m_j) \neq 0$. Then we smoothly extend $v_0 + \sum v_j$ to a vector field v on M having a finite number of non-degenerate zeros. By construction v does not vanish on N ; however πJv will have a zero at each m_j and at each m in N_p where $v(m) \in l_m \subset H_m$, as well as at each zero of v . There is much freedom in the choice of such a v , which we shall specify more precisely later.

Let m_j be one of the zeros of v_0 , and choose coordinates as in (4.1), (4.2), so that (z_1, x^α) are coordinates on M . We may assume that the hermitian metric on \tilde{M} has been chosen so that F_m coincides with the (y^α, z_n) -space for all m near m_j . The local orientations are given by

$$\Omega_T = \frac{i}{2} dz_1 \wedge d\bar{z}_1 \wedge dx^2 \wedge \cdots \wedge dx^{n-1}, \quad (5.4)$$

$$\Omega_F = \varepsilon_{n-2} dy^2 \wedge \cdots \wedge dy^{n-1} \wedge \frac{i}{2} dz_n \wedge d\bar{z}_n. \quad (5.5)$$

We set $G(z_1, x^\alpha) = (z_1, x^\alpha + if^\alpha, F)$, so that $G_{x_1}, G_{y_1}, G_{x^\alpha}$ span $T(M)$. In the local coordinates (z_1, x^α) on M we have

$$v = v_1 \partial / \partial z_1 + \bar{v}_1 \partial / \partial \bar{z}_1 + v^\alpha \partial / \partial x^\alpha, \quad (5.6)$$

so that as a vector in \mathbb{C}^n

$$v \equiv v[G] = (v_1, v^\alpha + iv[f^\alpha], v[F]),$$

where $v[\cdot]$ denotes directional derivative. It follows that

$$Jv \equiv iv[G] = (iv_1, -v[f^\alpha] + iv^\alpha, iv[F]),$$

so that

$$\begin{aligned} \pi Jv &\equiv iv[G] - c'G_{x_1} - c''G_{y_1} - c^\alpha G_{x^\alpha} \\ &= iv[G] - cG_{z_1} - \bar{c}G_{\bar{z}_1} - c^\alpha G_{x^\alpha} = (0, 0 + i*, *). \end{aligned}$$

Here $G_{z_1} = (1, if_{z_1}^\alpha, F_{z_1})$, $G_{\bar{z}_1} = (0, if_{\bar{z}_1}^\alpha, F_{\bar{z}_1})$, and $G_{x^\alpha} = (0, \delta_\alpha^\beta + if_{x^\alpha}^\beta, F_{x^\alpha})$, so that $c = iv_1$ and $c^\alpha = -v[f^\alpha]$. Hence, as a map from (z_1, x^α) -space to (y^α, z_n) -space, πJv has the form

$$\begin{aligned} y^\alpha &= v^\alpha - iv_1 f_{z_1}^\alpha + i\bar{v}_1 f_{\bar{z}_1}^\alpha + v[f^\beta]f_{x^\beta}^\alpha, \\ z_n &= iv[F] - iv_1 F_{z_1} + i\bar{v}_1 F_{\bar{z}_1} + v[f^\beta]F_{x^\beta}. \end{aligned} \quad (5.7)$$

If we substitute (5.7) into (5.5), we get (5.4) multiplied by the Jacobian factor

$$\varepsilon_{n-2} \frac{\partial(y^\alpha, z_n, \bar{z}_n)}{\partial(z_1, \bar{z}_1, x^\beta)}, \quad (5.8)$$

the sign of which gives the index of πJv at m_j . If we take into account (4.2), (5.7) becomes

$$\begin{aligned} y^\alpha &= v^\alpha - iv_1 b^\alpha \bar{z}_1 + i\bar{v}_1 b^\alpha z_1 + O(2), \\ z_n &= 2i\bar{v}_1 q_{\bar{z}_1} + O(2). \end{aligned}$$

We may assume that the H -component v_j added to v_0 is such that $v_1 \equiv 1$ near 0. Also, we assume that the extension of v from N to M is made so that the coefficients of v are locally independent of z_1 . Then at the origin (5.8) has the value

$$4(b^2 - 4a^2) \det(\partial v_0^\alpha / \partial x^\beta)(0). \quad (5.9)$$

The sign of the determinant is the index of v_0 at m_j , and $b^2 - 4a^2$ is positive if m_j is elliptic and negative if m_j is hyperbolic. Hence,

$$\varepsilon_{n-2} \text{ind}_{F, m_j}(\pi Jv) = \delta \text{ind}_{N, m_j}(v_0), \quad (5.10)$$

where $\delta = +1$ if m_i is elliptic or $\delta = -1$ if m_i is hyperbolic. If we sum (5.10) over the m_i in N_e , the right hand side is $\chi(N_e)$. To get $\chi(N_h)$ we must use $-v_0$ which multiplies the determinant in (5.9) by $(-1)^{n-2}$. Thus we get

$$\varepsilon_{n-2} \sum_j \text{ind}_{F, m_j}(\pi J v) = \chi(N_e) - (-1)^n \chi(N_h), \quad (5.11)$$

which accounts for the term $e - h$ in (5.1).

Finally, we consider a zero of $\pi J v$ at a point m in N_p which arises when $v(m)$, which spans the line k_m , lies in l_m . We first elaborate further on the construction of v along N_p . It is initially defined so that $k(N_p)$ intersects $l(N_p)$ transversely at m . Then it will be extended to N . We take coordinates as in (4.1), (4.3) with $c(x) = x^2$, so that N is given by (4.15). (x^3, \dots, x^{n-1}) gives coordinates on N_p , and $(y_1, x^3, \dots, x^{n-1})$ coordinates on N . In (5.6) we take $v_1 = v^1 + i$, $v^1 = \bar{v}^1$, so that

$$v = \partial/\partial y_1 + \sum_{j=1}^{n-1} v^j \partial/\partial x^j, \quad v^j(0) = 0. \quad (5.12)$$

The condition that v be tangent to N gives, via (4.15) and (5.12),

$$v^1 = O(2), \quad v^2 = -2\eta y_1 + \sum_{\beta=3}^{n-1} \eta_\beta x^\beta + O(2). \quad (5.13)$$

Thus we start with

$$v^\alpha = v^\alpha(x^3, \dots, x^{n-1}), \quad v^\alpha(0) = 0, \det \frac{\partial v^\alpha}{\partial x^\beta}(0) \neq 0, \quad 3 \leq \alpha, \beta \leq n-1, \quad (5.14)$$

and determine v^1 and v^2 by (5.13). We then extend this vector v locally from N_p to N by keeping (5.14) independent of y_1 , and from N to M by keeping (5.14) independent of x_1 and x^2 . Again we assume that F_m is the (y^α, z_n) -space for m near 0. Note that we may take $\eta = 1$, since $l_0 = k_0$ is transverse to N_p .

The parabolic index as defined in section 1 is computed relative to a coordinate system $(x_\star^\alpha, y_\star)$ with $y_\star = 0$ on N_p . Therefore we set (4.19)

$$y_\star = \frac{1}{2}\Delta = y_1 - \sum_{\beta=3}^{n-1} \eta_\beta x^\beta + O(2), \quad x_\star^\alpha = x^\alpha.$$

The chain rule in (1.14) gives

$$w_*^\alpha = w^\alpha (\partial y_*/\partial y_1 + w^\beta \partial y_*/\partial x^\beta)^{-1} = w^\alpha (1 - \eta_\beta w^\beta + O(1))^{-1}.$$

Since $w^\alpha = O(1)$ for both Y (4.18) and v (5.12), and $\partial/\partial x_*^\beta = \partial/\partial x^\beta$ for functions defined along N_p , we have $\partial w_*^\alpha/\partial x_*^\beta(0) = \partial w^\alpha/\partial x^\beta(0)$. Thus the parabolic intersection index at $m = 0$ is given (see (1.15)) by the sign of

$$\Omega_P\left(\frac{\partial l}{\partial x}, \frac{\partial k}{\partial x}\right)(0) = \det \begin{bmatrix} \delta_{\alpha\beta} & 0 \\ \delta_{\alpha\beta} & \partial v^\alpha/\partial x^\beta(0) \end{bmatrix}.$$

Hence,

$$\text{ind}_{P,m}(l, k) = \text{sgn} \det (\partial v^\alpha/\partial x^\beta(0))_{3 \leq \alpha, \beta \leq n-1}. \quad (5.15)$$

For the index of πJv at m we again compute the determinant (5.8). We substitute (4.3) into (5.7) and ignore second order terms. By (5.12) and (5.13) we get

$$\begin{aligned} y^2 &\equiv v^2 \equiv -2y_1 + \sum_{\beta=3}^{n-1} \eta_\beta x^\beta, & y^\alpha &\equiv v^\alpha(x^3, \dots, x^{n-1}), & 3 \leq \alpha \leq n-1, \\ z_n &\equiv 2i\bar{v}_1 Q_{z_1} \equiv 4x_1 + 2ix^2. \end{aligned}$$

Thus,

$$\frac{\partial(y^2, y^\alpha, z_n, \bar{z}_n)}{\partial(z_1, \bar{z}_1, x^2, x^\beta)}(0) = 16 \det [\partial v^\alpha/\partial x^\beta(0)]_{3 \leq \alpha, \beta \leq n-1}.$$

Comparison with (5.15) gives

$$\text{ind}_{F,m}(\pi Jv) = \varepsilon_{n-2} \text{ind}_{P,m}(l, k),$$

so that

$$\sum_{N_p} \text{ind}_{F,m}(\pi Jv) = \varepsilon_{n-2} p. \quad (5.16)$$

Combining (5.3), (5.11), and (5.16) gives (5.1), since $\varepsilon_n \varepsilon_{n-2} = -1$.

REFERENCES

- [1] E. BISHOP, *Differentiable manifolds in complex Euclidean space*, Duke Math. Jour. 32 (1965) 1–22.
- [2] S. S. CHERN, *Characteristic classes of hermitian manifolds*, Ann. Math. 47 (1946) 85–121.
- [3] S. S. CHERN and E. H. SPANIER, *A theorem on orientable surfaces in four-dimensional space*, Comment. Math. Helv. 25 (1951) 1–5.
- [4] M. W. HIRSCH, *Differential Topology*, Springer-Verlag, New York, 1976.
- [5] L. R. HUNT and R. O. WELLS, *The envelope of holomorphy of a two-manifold in \mathbb{C}^2* , Rice Univ. Studies 56 (1970) 51–62.
- [6] C. KENIG and S. WEBSTER, *The local hull of holomorphy of a surface in the space of two complex variables*, Invent. Math. 67 (1982) 1–21.
- [7] —, *On the hull of holomorphy of an n -manifold in \mathbb{C}^n* , Ann. Sc. Norm. Sup. Pisa vol. 11, no. 2 (1984) 261–280.
- [8] H. F. LAI, *Characteristic classes of real manifolds immersed in complex manifolds*, Trans. AMS 172 (1972) 1–33.
- [9] J. W. MILNOR and J. D. STASHEFF, *Characteristic classes*, Annals of Math. Studies, Princeton Univ. Press, 1974.
- [10] J. K. MOSER and S. M. WEBSTER, *Normal forms for real surfaces in \mathbb{C}^2 near complex tangents and hyperbolic surface transformations*, Acta Math 150 (1983) 255–296.
- [11] R. O. WELLS, *Compact real submanifolds of a complex manifold with nondegenerate holomorphic tangent bundles*, Math. Ann. 179 (1969) 123–129.

University of Minnesota
School of Mathematics
127 Vincent Hall
206 Church Street SE.
Minneapolis Minnesota 55455
USA

Received February 7, 1984