

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 60 (1985)  
  
**Artikel:** A non-immersion theorem for hyperbolic manifolds.  
**Autor:** Xavier, Frederico  
**DOI:** <https://doi.org/10.5169/seals-46313>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 11.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## A non-immersion theorem for hyperbolic manifolds

FREDERICO XAVIER

Let  $M^n$  be a complete hyperbolic  $n$ -manifold, so that  $M^n = H^n/\Gamma$  where  $\Gamma$  is a discrete group of isometries of hyperbolic space  $H^n$ . The manifold  $M^n$  is said to be *elementary* if the limit set of  $\Gamma$  in the sphere at infinity has at most two points. It is not difficult to see that in the case of non-elementary  $M^n$  the group  $\Gamma$  has infinite limit set. The purpose of this paper is to establish the following

**THEOREM.** *A non-elementary complete hyperbolic manifold  $M^n$  cannot be isometrically immersed in  $\mathbb{R}^{2n-1}$ .*

To put this result in the proper perspective let us recall that a well-known result of Hilbert [4] states that  $H^2$  cannot be isometrically immersed in  $\mathbb{R}^3$ . It has been conjectured for a long time that hyperbolic  $n$ -space  $H^n$  cannot be isometrically immersed in  $\mathbb{R}^{2n-1}$ ,  $n \geq 2$ . In this regard our theorem represents a partial verification of this conjecture.

The local analysis in Hilbert's theorem carries over to the higher dimensional case but it is not clear how to use this information at the global level. One source of difficulty lies in the fact that the sine-Gordon system, a highly non-linear system of partial differential equations associated to the problem ([1], [5]) is not yet fully understood. In our proof we abandon this system and concentrate on the metric dissimilarities between  $H^n$  and  $\mathbb{R}^n$ . However, the crucial relation (2) below – that leads in the case of surfaces to Tchebychef nets and, ultimately, to a contradiction – is not used in its full force. A better understanding of its significance might lead to sharper results. On the other hand, it seems that the existence of Tchebychef nets in higher dimensions is a less compelling evidence of a contradiction. In two dimensions they yield coverings of  $H^2$  by disjoint “squares” (quadrilaterals all of whose sides are asymptotic curves of equal lengths). Thinking about the effect that the isoperimetric inequality will have on the shape of the squares we eventually become convinced that they will have unbounded distortion and that singularities will arise. In the case  $n = 3$ , say, the “cubes” which are far

away will also have some degenerating angles but, at least in principle, their labels can change in a complicated manner and it is not geometrically clear how this will force appearance of singularities.

In the first version of this paper our theorem was proved in the case  $n = 3$  only, where we used existence of incompressible surfaces to guarantee that the group  $\Gamma$  was “big” enough. We are indebted to Dennis Sullivan for pointing out to us that, by an argument of Klein, non-elementary groups contain non-trivial free subgroups, regardless of dimension. This enabled us to extend our proof to all dimensions.

*Proof of the theorem.* Let us start by establishing the fact that if  $M^n = H^n/\Gamma$  is non-elementary then  $\Gamma$  contains non-trivial free subgroups. A hyperbolic (parabolic) isometry  $g \in \Gamma$  has a fundamental domain in the Poincaré model which is the complement of two disjoint balls about the fixed points (two externally tangent balls about the fixed point). For a power  $g^n$  the balls shrink to the fixed point ( $s$ ). Since the group is non-elementary we can find two isometries with disjoint sets of fixed points. Taking powers we can therefore assume that  $\Gamma$  contains isometries  $g_1, g_2$  having fundamental regions  $X_1, X_2$  satisfying  $X_1 \cup X_2 = H^n$ ,  $X_1 \cap X_2 \neq \emptyset$ . It follows that the image of a point  $p \in X_1 \cap X_2$  by a non-trivial word in  $g_1$  and  $g_2$  lies outside  $X_1 \cap X_2$ . In particular, there are no relations and the group generated by  $g_1$  and  $g_2$  is free.

Suppose now that  $j: M^n \rightarrow \mathbb{R}^{2n-1}$  is an isometric immersion. According to the previous paragraph we can pass to a covering and suppose that  $\pi_1(M^n)$  is free on two generators. Let  $p: H^n \rightarrow M^n$  be the canonical projection. As a consequence of Cartan’s theorem on exteriorly orthogonal forms it is possible to diagonalize simultaneously all second fundamental forms of the immersion  $p \circ j$  by orthonormal “principal” vectors  $e_1, \dots, e_n$  ([3]). Moreover, they can be extended to a global frame on  $H^n$ . Furthermore, the functions

$$a_i = \left( 1 + \sum_{\lambda=n+1}^{2n-1} b_{ii}^\lambda b_{ii}^\lambda \right)^{-1/2} \quad (\text{see [3], eq. 10}), \quad (1)$$

where the  $b_{ii}^\lambda$  are the components of the  $(n-1)$  second fundamental forms associated to an orthonormal basis of the normal space of the immersion, satisfy

$$\sum_{i=1}^n a_i^2 = 1, \quad [a_i e_i, a_j e_j] = 0 \quad (2)$$

(see bottom of page 161 and top of page 166 of [3]).

Following the flow lines of the (complete) vector fields  $a_i e_i$  we get a parametrization  $J: \mathbb{R}^n \rightarrow H^n$  whose coordinate vectors are the vectors  $a_i e_i$ . The pull back of the hyperbolic metric by  $J$  is  $g = \sum_{i=1}^n a_i^2 dx_i^2$ . We then have the following situation

$$(\mathbb{R}^n, g) \xrightarrow{J} H^n \xrightarrow{p} M^n \xrightarrow{i} \mathbb{R}^{2n-1}.$$

Let  $L$  be a compact neighborhood of two loops generating  $\pi_1(M^n)$  and let  $C$  be a connected component of  $J^{-1}(p^{-1}(L))$ . For a fixed  $q \in C$  above the base point, let  $C^s$  be the set of points in  $C$  that can be joined to  $q$  by a curve having  $g$ -length not greater than  $s$  and lying entirely in  $C$ . Let also  $B(r)$  be the euclidean ball centered at  $q$  of radius  $r$ . It follows from (1) that  $a_i \geq (1 + |b|^2)^{-1/2}$ , where  $b$  is the vector-valued second fundamental form. In particular, since  $C$  covers the compact set  $L$ , the coefficients  $a_i$  are bounded from below on  $C$  by a constant  $\delta > 0$ . Let now  $\alpha: [a, b] \rightarrow C \cap B(r)$  be a path starting at  $q$ . From

$$\int_a^b \left( \sum_{i=1}^n a_i^2 \alpha_i'^2 \right)^{1/2} dt \geq \delta \int_a^b \left( \sum_{i=1}^n \alpha_i'^2 \right)^{1/2} dt$$

we get  $C \cap B(r) \supset C^{\delta r}$ . Denoting by  $\text{Vol}(A)$  the  $g$ -volume of a set  $A \subset \mathbb{R}^n$  we have

$$\text{Vol } C^{\delta r} \leq \text{Vol } C \cap B(r) \leq \text{Vol } B(r) = \int_{B(r)} \left( \prod_{i=1}^n a_i \right) dx_1 \cdots dx_n = O(r^n),$$

by (2). This shows  $\text{Vol } C^s = O(s^n)$ . We claim that  $\text{Vol } C^s$  must actually grow exponentially. Let  $\mu$  denote the maximum length of the loops associated to the set  $L$ . The end point of the lift through  $q$  of a word of length at most  $s$  belongs to  $C^{\mu s}$ . Hence, for small enough  $\varepsilon > 0$ ,  $C^{\mu s + \varepsilon}$  contains at least as many disjoint balls of radius  $\varepsilon$  as there are words of length  $\leq s$ . Since  $\pi_1(M^n)$  is free on two generators the number of words grows exponentially with the length. Hence  $\text{Vol } C^s$  must grow exponentially with  $s$  and we have established a contradiction.

## REFERENCES

- [1] AMINOV, J. A., *Isometric immersions of domains of  $n$ -dimensional Lobachevsky space in  $(2n-1)$ -dimensional euclidean space*, Math. USSR Sbornik, Vol. 39 (1981), 3, 359–386.
- [2] CARTAN, E., *Sur les variétés de courbure constante d'un espace euclidien ou non-euclidien*, Bull. Soc. Math. France, 47 (1919), 125–160 and 48 (1920), 132–208.

- [3] MOORE, J. D., *Isometric immersions of space forms in space forms*, Pacific J. Math. 40 (1972), 157–166.
- [4] STOKER, J. J., *Differential Geometry*, Wiley-Interscience, New York 1969.
- [5] TENENBLAT, K. and TERNG, C. L., *Blacklund's theorem for  $n$ -dimensional submanifolds of  $R^{2n-1}$* , Annals of Math., 111 (1980), 477–490.

*Departamento de Matemática*  
*Universidade Federal de Pernambuco*

Received: August 7, 1984.