

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 60 (1985)

Artikel: Parallelizability of Finite H-Spaces.
Autor: Cappell, Sylvain / Weinberger, Sh.
DOI: <https://doi.org/10.5169/seals-46335>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Parallelizability of Finite H-Spaces

SLYVAIN CAPPELL and SHMUEL WEINBERGER

This note is a contribution to the study of how closely topological spaces with a multiplication (*H*-spaces) geometrically resemble Lie groups.

THEOREM. *Let X be a finite H -space and suppose that $X_{(2)}$ has a factor which at the prime 2 is S^7 , or a Lie group and that*

- 1) X is simply-connected, or
- 2) $\pi_1 X$ is a odd p -group, or
- 3) $\pi_1 X$ is infinite and has no 2-torsion.

Then X has the homotopy type of a closed parallelizable smooth manifold.

For a history and for our other related results see [2]. For instance, without the assumption on $X_{(2)}$ it is shown there that X is homotopy equivalent to a closed manifold. All known examples of finite *H*-spaces in fact are products of (just) S^7 , RP^7 and Lie groups at the prime 2, so that our theorem for the given fundamental groups includes all the known examples. Here, we just prove the theorem in case $\pi_1 X = 0$; the non-simply connected cases are readily dealt with using similar ideas together with methods of handling fundamental group difficulties developed in [2].

Notice that S^7 is the two-fold cover of RP^7 , and every Lie group is the two-fold cover of a closed manifold [2, Lemma 2.1]. Thus $X_{(2)}$ is the two-fold cover of the localization at 2 of a finite complex Y . There is a map $Y \rightarrow X_{(0)}$ so that the composite $X_{(2)} \rightarrow Y_{(2)} \rightarrow X_{(0)}$ is just localization. Let Z be the homotopy pullback:

$$\begin{array}{ccc} Z & \longrightarrow & X[\frac{1}{2}] \\ \downarrow & & \downarrow \\ Y_{(2)} & \longrightarrow & X_{(0)} \end{array}$$

Observation 1. Z is a finite Poincaré complex with 2-fold cover homotopy equivalent to X .

Observation 2. There is a map $Z \rightarrow BO$ lifting the Spivak normal fibration such that $X \rightarrow Z \rightarrow BO$ is nullhomotopic.

For the proofs of related statements see [3]. Now, consider a degree-one normal map $f: M \rightarrow Z$ associated to the lifting given by (2), and more importantly, the surgery obstruction of the two-fold cover $\tilde{f}: \tilde{M} \rightarrow X$. Note that \tilde{f} is covered by trivial bundle data. The only nonzero elements in $L_n(0)$ in the image of the transfer occur for $n \equiv 0 \pmod{4}$ and are detected by signature. Now $\text{sign}(\tilde{M}) = 0$ by the Hirzebruch signature formula and the stable parallelizability of M and $\text{sign}(X) = 0$ by an easy application of the Milnor–Moore theorem on Hopf algebras (e.g. X is rationally a product of odd spheres) so that this obstruction vanishes as well. Thus, we can complete surgery on \tilde{f} producing a stably parallelizable manifold N homotopy equivalent to X . Except in low dimensions where parallelizability is automatic, the obstructions to N being parallelizable are in fact just the Euler characteristic and, if the dimension is $\equiv 1 \pmod{2}$, the $Z/2Z$ semicharacteristic, (see e.g. [1]), but these vanish for N since they obviously do for X . Q.E.D.

REFERENCES

- [1] G. BREDON and A. KOSINSKI, *Vector fields on π -manifolds*, Ann. of Math. 84 (1966), 85–90.
- [2] S. CAPPELL and S. WEINBERGER, *Which H-spaces are manifolds? I*, (preprint).
- [3] S. CAPPELL and S. WEINBERGER, *Homology propagation of group actions*, (preprint).
- [4] S. WEINBERGER, *Homologically trivial group actions, I: Simply connected manifolds*. Amer. J. of Math., (to appear).

*Courant Institute of the Mathematical Sciences
The University of Chicago*

Received January 16, 1984