

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 62 (1987)

Artikel: Length of curves under conformal mappings.
Autor: Fernández, J.L. / Hamilton, D.H.
DOI: <https://doi.org/10.5169/seals-47343>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Length of curves under conformal mappings

JOSÉ L. FERNÁNDEZ and DAVID H. HAMILTON

1. Introduction

It is well known that for any homeomorphism f of the unit disk \mathbb{D} onto a domain Ω , where f is ACL and $\nabla f \in L^2(\mathbb{D})$, $f^{-1}(\Omega \cap L)$ has finite length for almost all rectifiable curves L . Suppose now that f is analytic, and let $\lambda(E)$ denote the Hausdorff linear measure of a set E . Hayman and Wu [8] proved that for any line L

$$\lambda(f^{-1}[\Omega \cap L]) < A, \quad (1)$$

for some absolute constant A . This was generalized by Garnett, Gehring and Jones [7] who gave conditions on a rectifiable Jordan curve in order that (1) holds for all Ω as above. It is necessary that L satisfy a regularity condition introduced by Ahlfors, i.e. there is a constant c_1 :

$$\lambda[L \cap \{|\xi - w| < r\}] \leq c_1 r \quad (2)$$

for all $\xi \in L$ and $r > 0$. Garnett, Gehring and Jones conjectured that (1) could fail for a regular quasicircle, i.e. L satisfies (2) together with

$$|z_1 - z_2| > c_2 \min_i \text{dia}(\gamma_i) \quad (3)$$

for any $z_1, z_2 \in L$ where γ_1, γ_2 are the two components of $L \setminus \{z_1, z_2\}$. In fact we show that the example suggested in [7] cannot work. A curve L is called quasismooth (or chord-arc) if there is a constant $M > 0$ such that for any $z_1, z_2 \in L$ we have

$$\min_{i=1,2} \lambda(\gamma_i) \leq M |z_1 - z_2|, \quad (4)$$

see Jerison and Kenig [9], Pommerenke [12]. Actually (2) and (3) are equivalent to (4). We prove:

THEOREM 1. *For any quasismooth curve L and any simply connected domain Ω with Riemann mapping f*

$$\lambda[f^{-1}(\Omega \cap L)] \leq A < \infty,$$

where A depends only on the chord arc constant M .

We conjecture that (2) is a necessary and sufficient condition on L in order that (1) hold for all conformal maps.

Next we consider the case of the universal covering mapping f of a multiply connected planar domain. Flinn [5] had obtained the following theorem: suppose that Ω is a hyperbolic planar domain and one component of $\mathbb{C} \setminus L$ is contained in Ω . Then if l is one component of $f^{-1}(\Omega \cap L)$ we have $\lambda(l) < \infty$. On the other hand if $\Omega = \mathbb{D} \setminus E$ where $E \subset (0, 1)$ is a closed set of zero logarithmic capacity then Belna, Cohn, Piranian and Stephenson [3] proved that there are circles L which do not satisfy (1).

Suppose that G is the Fuchsian group of Möbius transformations $T: \mathbb{D} \rightarrow \mathbb{D}$ which represents the cover group for Ω . The Dirichlet fundamental region \mathcal{D} for G is

$$\mathcal{D} = \{z \in \mathbb{D} : |T'(z)| < 1, \forall T \in G \setminus \{I\}\}$$

We say that \mathcal{D} is of finite length type if

$$\sum_G \lambda(\partial T\mathcal{D}) < \infty$$

THEOREM 2. *For any hyperbolic planar domain Ω of finite length type with universal covering map $f: \mathbb{D} \rightarrow \Omega$ and any quasismooth curve L :*

$$\lambda[f^{-1}(\Omega \cap L)] < \infty$$

From this we prove:

COROLLARY 1. *Suppose that Ω is a finitely connected hyperbolic planar domain with no point boundary components, and let f be the universal covering*

map. Then for any quasismooth curve L ,

$$\lambda[f^{-1}(\Omega \cap L)] < \infty.$$

The argument of the proof of Corollary 1 also shows how to construct infinitely connected domains for which the theorem holds.

However we do have:

COROLLARY 2. *Suppose that Ω is a Denjoy domain, i.e. $\partial\Omega \subset \mathbb{R}$. Then $\lambda[f^{-1}(\Omega \cap L)] < A$ for all quasismooth curves L if and only if Ω has finite length type.*

The fact that \mathcal{D} has finite length type says something about the “size” of the group G . The usual way of measuring that size is through the exponent of convergence $\delta(G)$

$$\delta(G) = \inf \left\{ \delta > 0: \sum_{T \in G} (1 - |T(0)|)^\delta < \infty \right\}$$

(see, e.g. [14]).

We have

COROLLARY 3. *Suppose that Ω is a planar domain and G the Fuchsian group uniformizing Ω then if $\delta(G) < \frac{1}{2}$ then for all quasismooth curves L*

$$\lambda(f^{-1}(\Omega \cap L)) < \infty.$$

The condition on $\delta(G)$ is sharp because for $\mathbb{D} \setminus \{0\}$ we have $\delta = \frac{1}{2}$ while $\lambda(f^{-1}(\mathbb{D} \setminus \{0\}) \cap \mathbb{R}) = \infty$. But on the other hand the condition is not necessary because there are finitely connected domains with no point boundary components for which $\delta > \frac{1}{2}$, e.g. take $\Omega_\varepsilon = \{z \in \mathbb{C} : |z| < \varepsilon, |z - 1| < \varepsilon, |z| > 1/\varepsilon\}$ with ε small enough (actually $\delta(\Omega_\varepsilon) \uparrow 1$ as $\varepsilon \rightarrow 0$).

2. Preliminary results

We shall be dealing with domains G which are regular for the Dirichlet problem. By dw_G^z we denote the unique probability measure such that if g is continuous on ∂G then the Perron solution u to the Dirichlet problem in G with

boundary values g is given by

$$u(z) = \int_{\partial G} g dw_G^z.$$

The harmonic measure of a Borel subset E of ∂G at a point $z \in G$ with respect to G is then

$$\omega(z, E, G) = \int_E dw_G^z.$$

Also the disk $\{|z - a| < r\}$ is denoted by $\Delta(a, r)$.

We make frequent use of the following results which are simple consequences of the Carleman–Milloux inequality, see [1], and Hall's lemma respectively, see [6].

LEMMA 1. *There is a positive function $c(\delta)$, $\delta > 0$, such that if the closure of a domain Ω contains continuum E which meets $\partial\Omega$ then for any $z \in \Omega \setminus E$ satisfying*

$$\text{dist}(z, E) \leq c(\delta) \text{dist}(z, \partial\Omega \setminus E)$$

we have

$$\omega(z, E, \Omega \setminus E) \geq 1 - \delta.$$

Let us denote the upper half plane by H . Also if $0 < a < b < 1$ and $\theta \in (0, \pi/2)$ then

$$S(a, b, \theta) = \{z \in H : |z| \in (a, b), \arg z \in (\theta, \pi - \theta)\}.$$

LEMMA 2. *Given $0 < a < b$, and $\theta \in (0, \pi/2)$ there exists $R > 0$ and $\eta > 0$ such that for any $r \geq R$ and for any continuum $E \subset H$ joining $|z| = 1$ to $|z| = r$*

$$\omega(z, (\Delta(0, r) \cap H) \setminus E) > \eta$$

for each z in the sector $S(a, b, \theta)$.

Also we shall be using quasiconformal mappings. We need

LEMMA 3. *Given $0 < a < b < 1$ and $\theta \in (0, \pi/2)$ there exists $R > 0$ and a positive function $\eta(k)$ such that for any $r \geq R$:*

If E is a continuum joining $|z|=1$ to $|z|=r$ in H , then for any k -quasiconformal mapping $\Phi: \mathbb{C} \rightarrow \mathbb{C}$ we have

$$\omega(\Phi(z), \Phi(E), \Phi(\Delta(0, r) \cap H \setminus E)) \geq \eta(k)$$

for each $z \in S(a, b, \theta)$.

The lemma follows from Lemma 2 and the distortion theorem of Mori, see [2]. Let Ω be the component of $\Delta(0, r) \cap H \setminus E$ containing $z \in S(a, b, \theta)$ and suppose f and g are the Riemann mappings from the unit disk \mathbb{D} onto (respectively) Ω and $\Phi(\Omega)$ with $f(0) = z$, $g(0) = \Phi(z)$. If $z \in S(a, b, \theta)$ then because of Lemma 2 $\omega(z, E, \Omega) > \eta$. On the other hand $2\pi \omega(z, E, \Omega)$ is the length of the subarc I_1 which is the closure of $\{e^{i\theta} : \lim_{r \rightarrow 1} f(re^{i\theta}) \in E\}$. One should note that as E is a continuum in \bar{H} , by a theorem of Beurling (see Collingwood and Lohwater [4]), “ $f^{-1}(E)$ ” is an arc of $\partial\mathbb{D}$ with a set of capacity zero removed. Similarly $2\pi \omega(\Phi(z), \Phi(\Omega))$ is the length of a subarc I_2 . But $g^{-1} \circ \Phi \circ f = \psi$ is a quasiconformal mapping of \mathbb{D} onto itself which fixes 0. Thus as $\psi(I_1) = I_2$ we see by Mori’s theorem that:

$$\lambda(I_1) \geq c \{\lambda(I_2)\}^\delta$$

where $c, \delta > 0$ depend only on k , which concludes the proof of the lemma.

The following is derived from estimates of Jerison and Kenig [9] and Kaufman and Wu [10, p. 269, 273].

LEMMA 4. *Suppose that U is a domain whose boundary is a quasismooth curve with constant M . If $z_0 \in U$, $\xi_0 \in \partial U$ satisfy (for some $r > 0$)*

$$\text{dist}(z_0, \partial U) \geq ar$$

and

$$|z_0 - \xi_0| \leq br$$

for some $a, b > 0$, then for any set $F \subset \partial U$ satisfying $F \subset \Delta(\xi_0, r)$ and $\lambda(F) \geq r/2$ we have

$$\omega(z_0, F, U) \geq \eta$$

where $\eta > 0$ depends only on a, b and M .

This is most easily proved by using Lemma 1 of [10] which provides us with a point $z_1 \in U$ satisfying

$$a_1^{-1}r \leq \text{dist}(z_1, \partial U) \leq a_1r$$

$$b_1^{-1}r \leq |\xi_0 - z_1| \leq b_1r$$

and

$$\omega(z_1, F, U) \geq \eta_1 > 0$$

where $a_1, b_1, \eta_1 > 0$ depend only on a, b and M . Now U is an (ε, ∞) domain (see [11]) and so there exists a rectifiable arc $\gamma \subset U$ joining z_0 to z_1 and satisfying

$$\lambda(\gamma) \leq a_2r$$

and

$$\text{dist}(\gamma, \partial U) \geq a_2^{-1}r$$

where a_2 depends only on a, b and K . Consequently Harnack's inequality is applied and we see that it is impossible that $\omega(z_0, F, U)$ may become arbitrarily small.

The connection between estimating harmonic measures and $\lambda(f^{-1}(\Omega \cap L))$ is derived from the notion of a Carleson measure (see [6]). Now a positive measure μ on the unit disk may be defined to be a Carleson measure if

$$\int_{\mathbb{D}} |T'(z)| d\mu < c \tag{5}$$

for any Möbius transformation $T: \mathbb{D} \rightarrow \mathbb{D}$. Clearly then, by considering $f \circ T$, any L satisfying (1) will have the property that arc length measure on $f^{-1}(\Omega \cap L)$ is a Carleson measure. This was observed in [7] and gives the extra conclusion that we have a Carleson measure.

LEMMA 5. *Suppose that L is a Jordan curve satisfying Ahlfors' regularity condition (2). Then to obtain $\lambda(f^{-1}[\Omega \cap L]) < c$ for all simply connected Ω the following are sufficient:*

There is a $\alpha > 0$, $\varepsilon > 0$ and $\beta < 1$ such that for any sequence $w_j \in L \cap \Omega$ with

$$|w_j - w_k| \geq \alpha \text{dist}(w_j, \partial \Omega), \quad j \neq k, \tag{6}$$

we have

$$\omega(w_j, K_j, \Omega \setminus K_j) \leq \beta \quad (7)$$

where

$$K_j = \bigcup_{k \neq j} \bar{\Delta}(w_k, \alpha \varepsilon \operatorname{dist}(w_k, \partial \Omega)) \quad (8)$$

3. Proof of Theorem 1

We let M denote the chord arc constant of L , see (4). Now we fix $\alpha = \frac{1}{3}$ and determine ε and β so that (7) of Lemma 5 is verified for any sequence $\{w_k\} \subset L \cap \Omega$ satisfying (6).

Fix j and let $z = w_j$. Also we define $d = \operatorname{dist}(z, \partial \Omega)$ and $J = L \cap \Omega$. Denote by J_1 the component of J which contains z and by J_0 the component of $J_1 \cap \Delta(z, \alpha d)$ containing z . Consider now the closed (in Ω) set $K = \bigcup_{w \in J \setminus J_0} \bar{\Delta}(w, \alpha \varepsilon \operatorname{dist}(w, \partial \Omega))$. Clearly $K \supset K_j \cup (J - J_0)$ and in particular by the maximum principle

$$\omega(z, K, \Omega \setminus K) \geq \omega(z, K_j, \Omega \setminus K_j) \quad (9)$$

So we have only to show that if we choose ε appropriately (depending only on L) we obtain $\beta = \beta(\varepsilon, M) < 1$ so that

$$\omega(z, K, \Omega \setminus K) \leq \beta. \quad (10)$$

Recall the function $c(\delta)$ of Lemma 1; then if $\varepsilon \leq c(\delta)$ we have

$$\omega(z, J \setminus J_0, \Omega \setminus (J \setminus J_0)) \geq (1 - \delta) \omega(z, K, \Omega \setminus K). \quad (11)$$

To see this we write

$$\omega(z, J \setminus J_0, \Omega \setminus (J \setminus J_0)) = \int_{\partial K} \omega(\zeta, J \setminus J_0, \Omega \setminus (J \setminus J_0)) dw_{\Omega \setminus K}^z(\zeta).$$

But by Lemma 1, if $\zeta \in K$ then $\omega(\zeta, J \setminus J_0, \Omega \setminus (J \setminus J_0)) \geq (1 - \delta)$, and (11) follows. In fact, if $w \in J \setminus J_0$ and $|\zeta - w| \leq \alpha \varepsilon \operatorname{dist}(w, \partial \Omega)$ then if E is the closure of the

component of $J \setminus J_0$ containing w we have

$$\begin{aligned} \text{dist}(\zeta, E) &\leq \alpha \varepsilon \text{dist}(w, \partial \Omega) \\ &\leq \alpha \varepsilon (1 - \alpha \varepsilon)^{-1} \text{dist}(\zeta, \partial \Omega) \\ &\leq c(\delta) \text{dist}(\zeta, \partial \Omega). \end{aligned}$$

The next step is to estimate $\omega(z, J \setminus J_0, \Omega \setminus (J \setminus J_0))$. Let U_1, U_2 be the complementary domains of L . Suppose that Ω_0 is the component of $\Omega \setminus (J \setminus J_0)$ containing z , and $\Omega_i = \Omega_0 \cap U_i$, $i = 1, 2$. Also we define $J_{i,j}$ to be the components of $(J \setminus J_1) \cap \partial \Omega_i$. Note that $J_{i,j}$ belongs to only one of the boundaries $\partial \Omega_i$.

The disk $\Delta(z, d\alpha/2M)$ contains no point of $J \setminus J_0$. We set $r = d\alpha/2M$. Since L is chord arc we have subarcs I_j of $\partial \Delta(z, r)$ such that for some $\rho, \tau > 0$ (depending only on M)

$$I_i \subset \Omega_i \tag{12}$$

$$\lambda(I_i) = \frac{\rho}{2} \pi r \tag{13}$$

$$\text{dist}(I_i, L) > \tau r. \tag{14}$$

Therefore

$$\begin{aligned} \omega(z, J \setminus J_0, \Omega \setminus (J \setminus J_0)) &= \int_{\partial \Delta(z, r)} \omega(\zeta, J \setminus J_0, \Omega \setminus (J \setminus J_0)) d\omega_{\Delta(z, r)}^z(\zeta) \\ &\leq (1 - \rho) + \frac{\rho}{2} + \frac{\rho}{2} \min_{i=1, 2} \max_{I_i} \omega(\zeta, J \setminus J_0, \Omega \setminus (J \setminus J_0)) \end{aligned} \tag{15}$$

But for $\zeta \in I_i$

$$\omega(\zeta, J \setminus J_0, \Omega \setminus (J \setminus J_0)) \leq \omega(\zeta, J, \Omega_i) \tag{16}$$

and we see from (15) that it is enough to show that

$$\min_i \max_{I_i} \omega(\zeta, J, \Omega_i) \leq \beta. \tag{17}$$

We consider now two cases. Let $B > 1$ be a constant to be determined later; B will depend only on M .

In this first case we suppose there exists $\zeta_0 \in (L \setminus J_1) \cap \partial\Omega_0$ so that

$$|\zeta_0 - z| \leq Bd. \quad (18)$$

In this case we let (following [10]) $S_i = L \setminus (\bigcup_j J_{i,j})$. Clearly $L \setminus J_1 = S_1 \cup S_2$. From the maximum principle we obtain for $\zeta \in I_i$

$$1 - \omega(\zeta, J, \Omega_i) = \omega(\zeta, \partial\Omega_i \setminus J, \Omega_i) \geq \omega(\zeta, S_i, U_i). \quad (19)$$

Now we use Lemma 4 for the chord arc domain U_i . Since $\zeta_0 \in L \setminus J_1$ we have that

$$\max_i \lambda(\Delta(\zeta_0, r) \cap S_i) \geq \frac{r}{2}. \quad (20)$$

Consequently by (18), (20) and Lemma 4 we obtain that

$$\max_{i=1,2} \min_{\zeta \in I_i} \omega(\zeta, S_i, U_i) \geq \eta > 0. \quad (21)$$

where η depends only on B and the chord arc constant M and so only on M . Then from (21), (19), (17), (15) and (11) we get

$$\omega(z, K, \Omega \setminus K) \leq \frac{\beta_1}{1 - \delta} \quad (22)$$

where $\beta_1 < 1$ depends only on M .

This leaves the case where for each $\zeta \in (L \setminus J_1) \cap \partial\Omega_0$ we have $|\zeta - z| > Bd$. Let $w \in \partial\Omega \cap \partial\Omega_0$, $|w - z| = d$. Notice that $w \notin L$ for $w \notin L \setminus J_1$ by our assumption (and $w \notin J_1$ as $w \notin \Omega$). We can join w to a point on $\partial\Delta(z, Bd)$ with a continuum

$$F \subset \partial\Omega \cap \partial\Omega_0 \cap (\bar{\Delta}(z, Bd) \setminus \Delta(z, d)),$$

because if not there would exist $w_1 \in \partial\Omega \cap \partial\Omega_0$, $|w_1 - z| < Bd$ and $w_1 \in L \setminus J_1$ contradicting our assumption in the second case. Since $F \subset \partial\Omega_0$ we have $F \cap (L \setminus J_1) = \emptyset$ and $F \cap J_1 = \emptyset$ so $F \cap L = \emptyset$.

Use a quasiconformal mapping Φ from \mathbb{C} to \mathbb{C} mapping \mathbb{R} onto L , $\Phi(0) = z$, $\Phi(\infty) = \infty$. Also we may assume $|\Phi(1) - z| = r$. Since F does not meet L , without loss of generality $F \subset U_1$ and Φ maps H onto U_1 . Now from the uniform bounds for quasiconformal mapping, (13) and (14) we obtain constants a, b, θ depending

only on the chord arc constant M so that

$$\Phi^{-1}(I_1) \subset S(a, b, \theta). \quad (23)$$

Now, if $\zeta \in I_1$

$$\omega(\zeta, \partial\Omega_1 \setminus J, \Omega_1) \geq \omega(\zeta, F, U_1). \quad (24)$$

But if $E = \Phi^{-1}(F)$ then E is a continuum running from $|z| = c_1$ to $|z| = c_2$, where c_1 depends only on k and c_2 depends on M and B , and $c_2 \rightarrow \infty$ as $B \rightarrow \infty$. Consequently from Lemma 3, (23) and (24) show that if $B \geq B_0(M)$

$$\omega(\zeta, J, \Omega_1) \leq \beta(M) < 1$$

for each $\zeta \in I_1$, and so, as in the first case, we obtain

$$\omega(z, K, \Omega \setminus K) \leq \frac{\beta_2}{1 - \delta} \quad (25)$$

where $\beta_2 < 1$ depends only on M .

Therefore we choose $\beta = \max(\beta_1, \beta_2)$ and $\delta < 1 - \beta$ and with $\varepsilon = c(\delta)$ see that the proof of the theorem is complete.

4. Proof of Theorem 2

We need the following (see [7]):

LEMMA 6. *Let U be a simply connected domain with rectifiable boundary, and f a conformal mapping of U onto Ω . Then for any quasismooth curve L*

$$\lambda(f^{-1}(\Omega \cap L)) < c_1 \lambda(\partial U).$$

Let g be the Riemann mapping from \mathbb{D} to U . Thus by Theorem 1 arc length $d\mu$ on $g^{-1} \circ f^{-1}(\Omega \cap L)$ is a Carleson measure and hence as $g' \in H^1$

$$\lambda(f^{-1}(\Omega \cap L)) = \int_{\mathbb{D}} |g'| d\mu \leq c_2 \int_{\partial\mathbb{D}} |g'| d\theta = c_2 \lambda(\partial U)$$

which proves the lemma.

Now let Ω be a hyperbolic planar domain and $f: \mathbb{D} \rightarrow \Omega$ be the universal covering map. Suppose that G is the Fuchsian group of Möbius transformations $T: \mathbb{D} \rightarrow \mathbb{D}$ which represents the cover group for Ω . The Dirichlet fundamental region \mathcal{D} for G is

$$\{z \in \mathbb{D} : |T'(z)| < 1, \forall T \in G \setminus \{I\}\}.$$

Now \mathcal{D} is a convex set in the hyperbolic metric with rectifiable boundary. Thus by Lemma 6

$$\lambda\{f^{-1}(\Omega \cap L) \cap \overline{T(\mathcal{D})}\} \leq c_3 \lambda(\partial T\mathcal{D}) \quad (26)$$

for any quasismooth curve L and $T \in G$. This immediately proves theorem 2.

Corollary 1 follows from

LEMMA 7. *Suppose that G is the Fuchsian group of a finitely connected planar domain with no point boundary components. Let \mathcal{D} be the Dirichlet region for G . Then*

$$\sum_G \lambda(T\partial\mathcal{D}) < \infty.$$

The boundary $\partial\mathcal{D}$ consists of a finite number of disjoint nonconcentric circles orthogonal to the unit disk. Let us denote \mathcal{D} as \mathcal{D}_1 . The region \mathcal{D}_2 is obtained from \mathcal{D} by reflecting \mathcal{D} through each of the orthogonal circles, and adding \mathcal{D}_1 . At the n^{th} stage we obtain \mathcal{D}_n with boundaries exactly n reflections of the original circles. Thus $\sum_G \lambda(T\mathcal{D}) = \sum_{n=1}^{\infty} \lambda(\partial\mathcal{D}_n)$. We need

LEMMA 8. *There is a constant $\beta < 1$ such that*

$$\lambda(\partial\mathcal{D}_n \cap \mathbb{D}) < \pi\beta^n$$

Let $E_n = \partial\mathcal{D}_n \cap \mathbb{D}$, F_n be a circle of E_{n-1} and G_n the part of E_n separated from the rest of E_n by F_n . By conformal invariance there is $\beta < 1$ such that $\text{dia}(G_n) \leq \beta \text{dia}(F_n)$. However as G_n consists of orthogonal semicircles $\lambda(G_n) \leq \pi \text{dia}(G_n)$. Summing over the components of E_{n-1} gives

$$\lambda(E_n) \leq \beta\lambda(E_{n-1})$$

Thus we prove Lemmas 7, 8 and complete the proof of Corollary 1.

The necessary part of Corollary 2 is derived from using the real line as our curve L . In this symmetric situation $f^{-1}(\mathbb{R})$ is $\bigcup_G T(\partial\mathcal{D})$.

Corollary 3 follows immediately from

LEMMA 9. *With the notations above if Ω is a hyperbolic planar domain then*

$$\sum_{T \in G} \lambda(\partial T\mathcal{D}) \leq 2\pi \left(\inf_{T \neq I} |T(0)| \right)^{-1} \sum_{T \in G} (1 - |T(0)|^2)^{1/2}$$

where C is a universal constant.

Proof. To see this we will associate to each side of \mathcal{D} and the $T(\mathcal{D})$'s an element $R \in G$ in a 1-1 fashion and in such a way that if $z \in s$ then $\rho(z, R(0)) \leq \rho(z, 0)$, where ρ denotes hyperbolic distance in \mathbb{D} . This is enough because then s is contained in a euclidean disk of radius $|R(0)|^{-1} (1 - |R(0)|^2)^{1/2}$ and so

$$\lambda(s) \leq \pi |R(0)|^{-1} \cdot (1 - |R(0)|^2)^{1/2}.$$

So consider the side s . It separates two contiguous images of \mathcal{D} , say $A(\mathcal{D})$, $B(\mathcal{D})$ with $A, B \in G$.

The transformations $\{T_i\}$ in G which pairwise identify the sides of \mathcal{D} generate G and in fact since Ω is planar G is freely generated by the $\{T_i\}$.

Now $A = B \circ T_0$ for some generator T_0 so that if $B = T_n \circ \cdots \circ T_1$ is a reduced word then the word length of A is $n - 1$ or $n + 1$ according to $T_1 = T_0^{-1}$ or $T_1 \neq T_0^{-1}$. Changing the roles of A and B we may assume that the latter case occurs and to s we associate $A = T_n \circ \cdots \circ T_1 \circ T_0$. Notice that $A = T_n \circ \cdots \circ T_1 \circ T_0$ determines s by being the side separating $T_n \circ \cdots \circ T_1(\mathcal{D})$ from $A(\mathcal{D})$. Finally $s \subset \partial A(\mathcal{D})$ and so for each $z \in s$ $\rho(z, A(0)) \leq \rho(z, 0)$.

REFERENCES

- [1] AHLFORS, L. V., *Conformal Invariants*, McGraw Hill, New York, (1973).
- [2] —, *Lectures on quasiconformal mapping*, Van Nostrand, Princeton, (1966).
- [3] BELNA, C., COHN, W., PIRANIAN, G. and STEPHENSON, K., *Level sets of special Blaschke products*, Michigan Math. J. 29 (1982), 79–81.
- [4] COLLINGWOOD, E. F. and LOHWATER, A. J., *Theory of Cluster Sets*, Cambridge University Press, (1966).
- [5] FLINN, B. B., *Hyperbolic Convexity and Level Sets of Analytic Functions*, Indiana Math. J. 32(6) (1983), 831–841.
- [6] GARNETT, J. B., *Bounded Analytic Functions*, Academic Press, New York, (1981).
- [7] GARNETT, J. F., GEHRING, F. W. and JONES, P. W., *Conformal invariant length sums*, Indiana Math. J. 32(6) (1983), 809–829.

- [8] HAYMAN, W. K. and WU, J. M., *Level sets of univalent functions*, Comment Mat. Helv. 56 (1981), 366–403.
- [9] JERISON, D. and KENIG, C., *Boundary behaviour of harmonic functions in non-tangentially accessible domains*, Advances in Math. 46 (1982), 80–147.
- [10] KAUFMAN, R. and WU, J. M., *Distortion of the boundary under conformal mapping*, Mich. Math. J. 29 (1982), 267–280.
- [11] MARTIO, O. and SARVAS, J., *Injectivity theorems in the plane and space*, Ann. Acad. Sci. Fenn. Ser. AI. Math. 4 (1979), 383–401.
- [12] POMMERENKE, C., *Boundary behaviour of conformal mapping*, in *Aspects of Contemporary Complex Analysis*, edited by Brannan, D. and Clunie, J., Academic Press, New York, 1980.
- [13] SULLIVAN, D., *The density at infinity of a discrete group of hyperbolic motions*, Publ. IHES 50 (1979), 419–450.

Dept. of Mathematics
University of Maryland
College Park, Maryland 20742
USA

Received June 18 1986