

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 62 (1987)

Artikel: SK1 of finite group rings: V.
Autor: Oliver, Robert
DOI: <https://doi.org/10.5169/seals-47357>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

SK₁ of finite group rings: V

ROBERT OLIVER

We continue here the study of

$$SK_1(\mathbb{Z}G) = \text{Ker } [K_1(\mathbb{Z}G) \rightarrow K_1(\mathbb{Q}G)]$$

for finite G : the group shown by Wall [26] to be precisely the torsion subgroup of $\text{Wh}(G)$. In earlier papers in this series, $SK_1(\mathbb{Z}G)$ has been studied via the extension

$$0 \rightarrow Cl_1(\mathbb{Z}G) \rightarrow SK_1(\mathbb{Z}G) \rightarrow \sum_{p \mid |G|} SK_1(\hat{\mathbb{Z}}_p G) \rightarrow 0; \quad (0.1)$$

where $Cl_1(\mathbb{Z}G) \subseteq SK_1(\mathbb{Z}G)$ is the subgroup of elements described via K_2 in localization sequences.

This paper contains the last step in deriving a combinatorial algorithm for describing the odd torsion in $SK_1(\mathbb{Z}G)$. By [17, Theorem 4.8], $SK_1(\mathbb{Z}G)[\frac{1}{2}]$ splits naturally as a sum

$$SK_1(\mathbb{Z}G)[\frac{1}{2}] \cong Cl_1(\mathbb{Z}G)[\frac{1}{2}] \oplus \sum_{p > 2} SK_1(\hat{\mathbb{Z}}_p G).$$

The groups $SK_1(\hat{\mathbb{Z}}_p G)$ (also for $p = 2$) are described by [15, Theorem 3] and [16, Theorem 2], in terms of $H_2(Z_i)$ for certain subgroups $Z_i \subseteq G$. On the other hand, in [17], the problem of describing $Cl_1(\mathbb{Z}G)_{(p)}$ for any odd prime p and any finite G is reduced to the case where G is a p -group (see [17, Theorem 4.8], and the discussion at the end of Section 3 below).

The following theorem is the central result of this paper, and gives a relatively simple way of describing $Cl_1(\mathbb{Z}G)$ when G is a p -group (and p odd). Note that if G is any group, and G acts on $\mathbb{Z}G$ by conjugation, then for any set $S \subseteq G$ of conjugacy class representatives,

$$H_1(G; \mathbb{Z}G) \cong \sum_{h \in S} H_1(Z_G(h)) \otimes \mathbb{Z}(h).$$

(If $X \subseteq G$ is any conjugacy class, and $h \in X$, then $\mathbb{Z}(X) \cong \text{Ind}_{\mathbb{Z}_G(h)}^G(\mathbb{Z})$ as $\mathbb{Z}G$ -modules.) Thus, $H_1(G; \mathbb{Z}G)$ is generated by elements $g \otimes h$ for commuting $g, h \in G$.

THEOREM 3.6. *Fix an odd prime p and a p -group G . Write $\mathbb{Q}G = \prod_{i=1}^k B_i$, where each B_i is simple with center F_i and irreducible representation V_i . For each i , let $(\mu_{F_i})_p$ be the group of p -th power roots of unity. Define*

$$\psi_G : H_1(G; \mathbb{Z}G) \rightarrow \prod_{i=1}^k (\mu_{F_i})_p,$$

where G acts on $\mathbb{Z}G$ by conjugation, by setting

$$\psi_G(g \otimes h) = [\det_{F_i}(g, V_i^h)]_i \quad (g, h \in G, gh = hg, V_i^h = \{x \in V_i : hx = x\}).$$

Then $Cl_1(\mathbb{Z}G) \cong \text{Coker } (\psi_G)$.

Examples of computations of $Cl_1(\mathbb{Z}G)$ using Theorem 3.6 for non-abelian G are given in Section 4. For abelian G , the isomorphism $SK_1(\mathbb{Z}G) \cong \text{Coker } (\psi_G)$ is proven in [1, Theorem 1.8], and some examples of calculations of $SK_1(\mathbb{Z}G)$ using that are given in Section 5 of the same paper.

Theorem 3.6 (and the other theorems referred to above) are stated, for simplicity, as describing the components of $SK_1(\mathbb{Z}G)$ as abstract groups only. But the proofs also contain enough information so that one can take a specific element in $SK_1(\mathbb{Z}G)$ (e.g., a specific element in $\text{Coker } (\psi_G)$ as described above), and represent it by a matrix. The opposite problem, taking a specific matrix over $\mathbb{Z}G$ and deciding how it sits in $SK_1(\mathbb{Z}G)$ (if it does) is harder in general; the study in [20] of the Whitehead transfer homomorphism for oriented S^1 -fiber bundles gives one example where this can be done.

In general, for any finite group G , $Cl_1(\mathbb{Z}G)$ is described by localization exact sequences

$$K_2^{\text{top}}(\hat{\mathbb{Z}}_p G) \xrightarrow{\varphi} C_p(\mathbb{Q}G) \xrightarrow{\partial} Cl_1(\mathbb{Z}G)_{(p)} \rightarrow 0$$

for each prime p ; where for any maximal order $\mathfrak{M} \subseteq \mathbb{Q}G$:

$$\begin{aligned} C_p(\mathbb{Q}G) &\cong \varprojlim_n \text{Coker } [K_2(\mathfrak{M}) \rightarrow K_2(\mathfrak{M}/p^n \mathfrak{M})] \cong \varprojlim_n Cl_1(\mathfrak{M}; p^n \mathfrak{M}) \\ &\cong \text{Coker } \left[K_2\left(\mathfrak{M}\left[\frac{1}{p}\right]\right) \rightarrow K_2^{\text{top}}(\hat{\mathbb{Q}}_p G) \right]_{(p)}. \end{aligned}$$

The $C_p(\mathbb{Q}G)$ are described by the work of Bak and Rehmann on the congruence subgroup problem [3]. The remaining problem is then to find a set of generators for $K_2^{\text{top}}(\hat{\mathbb{Z}}_p G)$, or at least for its image in $C_p(\mathbb{Q}G)$. In the case of an odd prime p and a p -group G , the formula

$$Cl_1(\mathbb{Z}G) \cong \text{Coker} \left[\psi_G : H_1(G; \mathbb{Z}G) \rightarrow \prod_{i=1}^k (\mu_{F_i})_p \right]$$

can be explained by noting that norm residue symbols define an isomorphism of $C_p(\mathbb{Q}G)$ with $\prod (\mu_{F_i})_p$, and that $H_1(G; \mathbb{Z}G) (\cong H_1(G; \hat{\mathbb{Z}}_p G))$ and $K_2(\hat{\mathbb{Z}}_p G)$ both are closely related to the cyclic homology group $HC_1(\hat{\mathbb{Z}}_p G)$ (see [21]).

The key new result here about generators for $K_2(\hat{\mathbb{Z}}_p G)$ is:

THEOREM 1.4. *Let p be any prime, and fix a p -group G and an element $z \in Z(G)$. Then*

$$\begin{aligned} \text{Ker} [K_2^{\text{top}}(\hat{\mathbb{Z}}_p G) \rightarrow K_2^{\text{top}}(\hat{\mathbb{Z}}_p[G/z])] \\ = \langle \{g, 1 + \lambda(1 - z)^i h\} : \lambda \in \hat{\mathbb{Z}}_p, i \geq 1, g, h \in G, gh = hg \rangle. \end{aligned}$$

Since $\text{Coker} [K_2^{\text{top}}(\hat{\mathbb{Z}}_p G) \rightarrow K_2^{\text{top}}(\hat{\mathbb{Z}}_p[G/z])]$ is also known in the above situation (see Proposition 2.1 below), it should in principle now be possible to inductively construct a set of generators for $K_2^{\text{top}}(\hat{\mathbb{Z}}_p G)$. Unfortunately, it's not always easy to explicitly lift elements from $K_2^{\text{top}}(\hat{\mathbb{Z}}_p[G/z])$ to $K_2^{\text{top}}(\hat{\mathbb{Z}}_p G)$, even where they are known to lift. But such an inductive procedure does work to give generators for $K_2^{\text{top}}(\hat{\mathbb{Z}}_p G)^+$ when p is odd, and this suffices when computing $Cl_1(\mathbb{Z}G)$.

Another consequence of Theorem 1.4 involves a comparison of $Cl_1(RG)$ – when G is any finite group and R the ring of integers is some number field $K \subseteq \mathbb{C}$ – with the “complex Artin cokernel”

$$A_{\mathbb{C}}(G) = \text{Coker} \left[\sum \{R_{\mathbb{C}}(H) : H \subseteq G \text{ cyclic}\} \xrightarrow{\Sigma \text{ Ind}} R_{\mathbb{C}}(G) \right].$$

Natural epimorphisms $I_{RG} : A_{\mathbb{C}}(G) \twoheadrightarrow Cl_1(RG)$ are constructed, for such R and G , via localization sequences. Theorem 1.4 can then be applied to show that for any G , I_{RG} is an isomorphism for R large enough. Thus, $A_{\mathbb{C}}(G)$ represents the “largest possible” $Cl_1(RG)$ when G is fixed and R varies. This is the second unexpected appearance of Artin cokernels when studying $K_n(RG)$: it was shown in [18] that $D(\mathbb{Z}G)^+ \cong A_{\mathbb{Q}}(G)$ when G is a p -group and p any odd regular prime.

The obvious remaining question is: what about 2-power torsion in $SK_1(\mathbb{Z}G)$? Unlike the case of odd torsion, this cannot be completely reduced to studying $Cl_1(\mathbb{Z}G)$ for 2-groups G , but the results in [17] show that the main problem is with 2-groups. If G is a p -group (for any p) and $[G, G]$ is central and cyclic, then we can show that $K_2^{\text{top}}(\hat{\mathbb{Z}}_p G)$ is generated by $\{-1, -1\}$ and symbols $\{g, u\}$ for $g \in G$ and $u \in (\hat{\mathbb{Z}}_p[Z_G(g)])^*$; and when $p = 2$ this suffices to get a description of $Cl_1(\mathbb{Z}G)$. But there are 2-groups G for which $K_2^{\text{top}}(\mathbb{Z}_2 G)$ is not generated by such symbols, and there may not be any simple algorithm for describing $Cl_1(\mathbb{Z}G)$ in general. The best conjecture we have been able to make so far gives upper and lower bounds for $Cl_1(\mathbb{Z}G)$, bounds which differ by exponent two. The question of whether the inclusion $Cl_1(\mathbb{Z}G)_{(2)} \subseteq SK_1(\mathbb{Z}G)_{(2)}$ ever fails to split is also still open.

The paper is organized as follows. Section 1 and 2 deal with the problems of finding generators for $\text{Ker}(K_2(\hat{\mathbb{Z}}_p \alpha))$, and of detecting $\text{Coker}(K_2(\hat{\mathbb{Z}}_p \alpha))$, respectively, when α is a surjection of p -groups whose kernel is central and cyclic. This is applied in Section 3 to prove that $Cl_1(\mathbb{Z}G) \cong \text{Coker}(\psi_G)$ when G is an odd p -group; and ways of using that to compute the odd torsion in $Cl_1(\mathbb{Z}G)$ for arbitrary finite G are discussed. Examples are given in Section 4 to illustrate how Theorem 3.6 works in practice for computing $Cl_1(\mathbb{Z}G)$. Finally, in Section 5, the relationship between $Cl_1(RG)$ and the complex Artin cokernel is studied, and the isomorphism $Cl_1(RG) \cong A_{\mathbb{C}}(G)$ proven for large R .

As for notation, C_n always denotes a (multiplicative) cyclic group of order n , and ζ_n a primitive n -th root of unity. If F is any field, then μ_F denotes the group of roots of unity in F , and $(\mu_F)_p$ the group of p -th power roots of unity.

If R is a $\hat{\mathbb{Q}}_p$ -algebra or a $\hat{\mathbb{Z}}_p$ -order (e.g., $R = \hat{\mathbb{Q}}_p G$ or $\hat{\mathbb{Z}}_p G$), then $K_2(R)$ always denotes the topological K_2 . The precise definition of these groups, and their occurrence in localization sequences, is described in [20]: in Theorem 2.1 and the preceding discussion (see also [3]). Here we just note that if R is a $\hat{\mathbb{Z}}_p$ -order, then

$$K_2^{\text{top}}(R) \cong \varprojlim_n K_2(R/p^n R).$$

Section 1

If R is a ring, and $I \subseteq R$ is a 2-sided ideal, we define here

$$K_2(R, I) = \text{Ker}[K_2(R) \rightarrow K_2(R/I)].$$

A braid diagram analogous to that in [12, Remark 6.6] shows that for any ideals

$\bar{I} \subseteq I \subseteq R$, there is an exact sequence

$$0 \rightarrow K_2(R, \bar{I}) \rightarrow K_2(R, I) \rightarrow K_2(R/\bar{I}, I/\bar{I}) \xrightarrow{\partial} K_1(R, \bar{I}) \rightarrow K_1(R, I) \rightarrow \dots$$

The main result of this section is to describe a set of generators for $K_2(\hat{\mathbb{Z}}_p G, (1 - z))$; when p is any prime, G is any p -group, and $z \in Z(G)$. Three lemmas will first be needed.

LEMMA 1.1. *Fix a prime p , and a finite ring R of p -power order. Let $J \subseteq R$ be the Jacobson radical, and let $\{\alpha_1, \dots, \alpha_k\} \subseteq J$ be any set of elements such that $\{p, \alpha_1, \dots, \alpha_k\}$ generates J (as an ideal). Then for any ideal $I \subseteq J$ of R such that $IJ = JI = 0$, and such that $I \subseteq \langle \alpha_1, \dots, \alpha_k \rangle_R$ if $p = 2$, $K_2(R, I)$ is generated by symbols of the form*

$$\{1 - \alpha_i, 1 - x\} : 1 \leq i \leq k, \quad x \in I. \quad (1)$$

Proof. We use the notation and relations for pointed bracket symbols in [25, Proposition 96–97]. By [17, Proposition 2.3], $K_2(R, I)$ is generated by symbols of the form

$$\{1 - \alpha, 1 + x\} = \langle \alpha, 1 + x \rangle = \langle \alpha, x \rangle$$

for $\alpha \in J$ and $x \in I$ ($\alpha x = x\alpha = 0$). Write $\alpha = pr_0 + \alpha_1 r_1 + \dots + \alpha_k r_k$; so that

$$\begin{aligned} \langle \alpha, x \rangle &= \langle pr_0, x \rangle + \sum_{i=1}^k \langle \alpha_i r_i, x \rangle = \langle p, r_0 x \rangle + \sum_{i=1}^k \langle \alpha_i, r_i x \rangle \\ &= \sum_{i=1}^k \{1 - \alpha_i, 1 + r_i x\} + \langle p, r_0 x \rangle + p \langle -1, r_0 x \rangle \\ &= \sum_{i=1}^k \{1 - \alpha_i, 1 + r_i x\} + \langle p, r_0 x \rangle + \left\langle -p + \binom{p}{2} r_0 x, r_0 x \right\rangle \quad (x^2 = 0) \\ &= \sum_{i=1}^k \{1 - \alpha_i, 1 + r_i x\} + \left\langle \binom{p}{2} r_0 x, r_0 x \right\rangle. \quad (px \in JI = 0). \end{aligned}$$

If p is odd, then $\binom{p}{2} x = 0$, and we are done. If $p = 2$ and $I \subseteq \langle \alpha_1, \dots, \alpha_k \rangle_R$, then the same procedure shows that for any $x \in I$, $\langle x, x \rangle$ is a sum of symbols of the form in (1). \square

The following technical relation between symbols will be needed in the calculations.

LEMMA 1.2. *Let R be any ring. Fix $a, u \in R^*$ and $n \geq 2$ such that*

$$[a^n, u] = 1 = [a^i u a^{-i}, a^j u a^{-j}]$$

for any i, j . Then

$$\begin{aligned} \{a, u(aua^{-1})(a^2ua^{-2}) \cdots (a^{n-1}ua^{1-n})\} \\ = \{a^n, u\} + (n-1)\{u, u\} + \sum_{i=1}^{n-1} \{a^i u a^{-i}, u\}. \end{aligned}$$

Proof. In $St(R)$, set $x = h_{12}(u)$, $y = h_{13}(a)$, and

$$T = (yxy^{-1})(y^2xy^{-2}) \cdots (y^{n-1}xy^{1-n}).$$

Then

$$\begin{aligned} \{a, u(aua^{-1}) \cdots (a^{n-1}ua^{1-n})\} &= [y, xT] \\ &= (yxy^{-1})(yTy^{-1})T^{-1}x^{-1} = T(y^nxy^{-n})T^{-1}x^{-1} = [T, y^nxy^{-n}][y^n, x] \\ &= (\text{diag}(aua^{-1} \cdot a^2ua^{-2} \cdots a^{n-1}ua^{1-n}, u^{1-n}) * \text{diag}(u, u^{-1})) + \{a^n, u\} \\ &= \sum_{i=1}^{n-1} \{a^i u a^{-i}, u\} + \{u^{1-n}, u^{-1}\} + \{a^n, u\}. \end{aligned}$$

Here, for commuting matrices $M, N \in E(R)$, $M^*N \in K_2(R)$ denotes the commutator $[\tilde{M}, \tilde{N}]$ of liftings to $\tilde{M}, \tilde{N} \in St(R)$. \square

The third lemma will be needed when constructing filtrations of group rings by ideals. By a *p-ring* is meant the ring of integers in any finite extension of \mathbb{Q}_p .

LEMMA 1.3. *Fix a prime p , a p -group G , and some $z \in Z(G)$. Let $p^n = |z|$. Then, for any p -ring A , there are isomorphisms*

$$f_k : A/p^n[G/z] \xrightarrow{\cong} \frac{(1-z)^k AG}{(1-z)^{k+1} AG} \quad (k \geq 1)$$

and

$$f'_k: A/p[G/z] \xrightarrow{\cong} \frac{(1-z)^k A/p[G]}{(1-z)^{k+1} A/p[G]}; \quad (1 \leq k \leq p^n - 1)$$

both induced by sending ξ to $(1-z)^k \xi$ for $\xi \in AG$.

Proof. Note first that for any $\xi \in AG$, and any $k \geq 1$,

$$(1-z)^k p^n \xi \equiv (1-z)^k (1+z+z^2+\cdots+z^{p^n-1}) \xi = 0 \pmod{(1-z)^{k+1} AG}. \quad (1)$$

Thus, $(1-z)^k AG / (1-z)^{k+1} AG$ has exponent at most p^n for $k \geq 1$; and is in particular finite. So the map

$$(1-z)^k: (1-z)AG \xrightarrow{\cong} (1-z)^{k+1}AG$$

is an isomorphism: it is clearly onto, and the groups are free A -modules of the same rank.

Thus, for $\xi \in AG$ and $k \geq 1$, if $(1-z)^k \xi = (1-z)^{k+1} \eta$ for some $\eta \in AG$, then $(1-z)(\xi - (1-z)\eta) = 0$, and so

$$\xi \in (1-z)\eta + (1+z+\cdots+z^{p^n-1})AG \subseteq (1-z)AG + p^nAG.$$

Together with (1), this shows that $(1-z)^k \xi \in (1-z)^{k+1}AG$ if and only if $\xi \in p^nAG + (1-z)AG$. So f_k is well defined and an isomorphism.

If $1 \leq k \leq p^n - 1$, and $\xi' \in A/p[G]$ is such that $(1-z)^k \xi' \in (1-z)^{k+1}A/p[G]$ then

$$(1+z+\cdots+z^{p^n-1})\xi' = (1-z)^{p^n-1}\xi' \in (1-z)^{p^n}A/p[G] = 0;$$

and so $\xi' \in (1-z)A/p[G]$. The converse is clear, and so f'_k is a well defined isomorphism. \square

The main result of this section can now be shown:

THEOREM 1.4. *Fix a prime p , an unramified p -ring A , a p -group G , and an element $z \in Z(G)$. Then*

$$K_2(AG, (1-z)AG) = \text{Ker } [K_2(AG) \rightarrow K_2(A[G/z])]$$

is a finite group, and is generated by symbols of the form

$$\{g, 1 - \lambda(1 - z)^i h\} : g, h \in G, [g, h] = 1, \lambda \in A, i \geq 1.$$

Proof. Let $H_0 = \langle z \rangle$, and fix a series of subgroups

$$H_0 \subseteq H_1 \subseteq \cdots \subseteq H_n = G$$

such that for each $i = 1, \dots, n$,

$$H_i \triangleleft G \quad \text{and} \quad [H_i : H_{i-1}] = p.$$

For each i , fix $z_i \in H_i \setminus H_{i-1}$. Note that in G/H_{i-1} , z_i is central of order p .

Let $p^m = |z|$. Define

$$S = \{(k; r, i_0, \dots, i_k) : 0 \leq k \leq n, i_0 \geq 1, 0 \leq r \leq m-1, 0 \leq i_1, \dots, i_k \leq p-1\}.$$

For each $\sigma = (k; r, i_0, \dots, i_k) \in S$, set $k(\sigma) = k$, and

$$X(\sigma) = p^r(1 - z)^{i_0}(1 - z_1)^{i_1} \cdots (1 - z_k)^{i_k} \in AG.$$

Define ideals $I'(\sigma) \subseteq I(\sigma) \subseteq AG$ by setting

$$I'(\sigma) = \langle (1 - z)^{i_0+1}, p^{r+1}(1 - z)^{i_0}; p^r(1 - z)^{i_0}(1 - z_1)^{i_1} \cdots (1 - z_j)^{i_j+1} : 1 \leq j \leq k \rangle$$

and

$$I(\sigma) = I'(\sigma) + \langle X(\sigma) \rangle.$$

The idea now is to use S as a bookkeeping system for filtering the ideal $(1 - z)AG$ into “pieces” small enough so that the theorem can be proven starting with Lemma 1.1. The following diagram gives a visual overview of this

filtration in the case where $p = 3$, $m = 2$, and $n = 2$ (i.e., $|z| = 9$ and $|G| = 81$):

		$k(\sigma) = 0$	$k(\sigma) = 1$	$k(\sigma) = 2$	
$(1 - z)AG$					
				(2; 0, 1, 0, 0)	
			(1; 0, 1, 0)	(2; 0, 1, 0, 1)	
				(2; 0, 1, 0, 2)	
				(2; 0, 1, 1, 0)	
			(1; 0, 1, 1)	(2; 0, 1, 1, 1)	
				(2; 0, 1, 1, 2)	
				(2; 0, 1, 2, 0)	
			(1; 0, 1, 2)	(2; 0, 1, 2, 1)	
				(2; 0, 1, 2, 2)	
				(2; 1, 1, 0, 0)	
			(1; 1, 1, 0)	(2; 1, 1, 0, 1)	
				(2; 1, 1, 0, 2)	(1)
				(2; 1, 1, 1, 0)	
			(1; 1, 1, 1)	(2; 1, 1, 1, 1)	
				(2; 1, 1, 1, 2)	
				(2; 1, 1, 2, 0)	
			(1; 1, 1, 2)	(2; 1, 1, 2, 1)	
				(2; 1, 1, 2, 2)	
$(1 - z)^2AG$					
				(2; 0, 2, 0, 0)	
			(1; 0, 2, 0)	(2; 0, 2, 0, 1)	
				(2; 0, 2, 0, 2)	
				(2; 0, 2, 1, 0)	
			(1; 0, 2, 1)	(2; 0, 2, 1, 1)	
				(2; 0, 2, 1, 2)	

The horizontal lines represent ideals in AG , ordered sequentially with the largest at the top. Each box represents some element $\sigma \in S$; the horizontal line at the top of the box represents $I(\sigma)$, while the line at the bottom represents $I'(\sigma)$. That the $I(\sigma)$ and $I'(\sigma)$ actually do correspond with this picture will be shown in Step 2A below.

Step 1. We now show that for any $\sigma \in S$, there is an isomorphism

$$f_\sigma : A/p[G/H_{k(\sigma)}] \xrightarrow{\cong} I(\sigma)/I'(\sigma) \quad (2)$$

defined by setting $f_\sigma([\xi]) = [X(\sigma) \cdot \xi]$ for $\xi \in AG$. This will be proven by induction on $k = k(G)$. If $k = 0$, so $\sigma = (0; r, i)$ for some $i \geq 1$ and $0 \leq r \leq m - 1$, then

$$(1 - z)^i AG / (1 - z)^{i+1} AG \cong A/p^m[G/Z] = A/p^m[G/H_0]$$

by Lemma 1.3; and so

$$I(\sigma)/I'(\sigma) = \frac{p^r(1 - z)^i AG + (1 - z)^{i+1} AG}{p^{r+1}(1 - z)^i AG + (1 - z)^{i+1} AG} \cong \frac{p^r A/p^m[G/H_0]}{p^{r+1} A/p^m[G/H_0]} \cong A/p[G/H_0].$$

Now assume $k \geq 1$, and write $\sigma = (k; r, i_0, \dots, i_k)$. Set

$$\hat{\sigma} = (k - 1; r, i_0, \dots, i_{k-1}) \in S.$$

By induction, we can assume that $I(\hat{\sigma})/I'(\hat{\sigma}) \cong A/p[G/H_{k-1}]$. By definition

$$I'(\sigma) = I'(\hat{\sigma}) + X(\hat{\sigma})(1 - z_k)^{i_k+1} AG,$$

$$I(\sigma) = I'(\hat{\sigma}) + X(\hat{\sigma})(1 - z_k)^{i_k} AG,$$

$$I(\hat{\sigma}) = I'(\hat{\sigma}) + X(\hat{\sigma})AG.$$

Thus, $I(\hat{\sigma}) \supseteq I(\sigma) \supseteq I'(\sigma) \supseteq I'(\hat{\sigma})$; and by Lemma 1.3:

$$I(\sigma)/I'(\sigma) \cong \frac{(1 - z_k)^{i_k} A/p[G/H_{k-1}]}{(1 - z_k)^{i_k+1} A/p[G/H_{k-1}]} \cong A/p[G/H_k].$$

(Recall that $H_k = \langle H_{k-1}, z_k \rangle$, and that $0 \leq i_k \leq p - 1$.)

Step 2. We next show that for any $\sigma \in S$,

$$K_2(AG/I'(\sigma), I(\sigma)/I'(\sigma)) = \langle \{g, 1 - X(\sigma)\lambda h\} : [g, h] \in H_{k(\sigma)}, \lambda \in A \rangle. \quad (3)$$

This will be proven by downwards induction on $k = k(\sigma)$.

Note first that AG is a local ring with Jacobson radical

$$J(AG) = \langle p, 1 - g : g \in G \rangle. \quad (4)$$

If $\sigma \in S$ and $k(\sigma) = n$, then $H_n = G$, and so $I(\sigma)/I'(\sigma) \cong A/p$ by Step 1. In particular,

$$(I(\sigma)/I'(\sigma)) \cdot J(AG/I'(\sigma)) = 0 = J(AG/I'(\sigma)) \cdot (I(\sigma)/I'(\sigma)).$$

So (3) follows in this case from Lemma 1.1 (applied using $\{1 - g : g \in G\}$ for the α_i 's).

Now fix some $\sigma = (k; r, i_0, \dots, i_k) \in S$, where $k < n$. For each $0 \leq i \leq p-1$, set

$$\sigma_i = (k+1; r, i_0, \dots, i_k, i) \in S.$$

Assume inductively that (3) holds for the σ_i .

Step 2A. We now show that the $I(\sigma_i) \supseteq I'(\sigma_i)$ and $I(\sigma) \supseteq I'(\sigma)$ have the relations implied by diagram (1) above. By definition, $I(\sigma_0) = I(\sigma)$ ($X(\sigma_0) = X(\sigma)$). For any $0 \leq i \leq p-2$,

$$I'(\sigma_i) = I'(\sigma) + X(\sigma)(1 - z_{k+1})^{i+1}AG = I(\sigma_{i+1}). \quad (5)$$

Furthermore,

$$I'(\sigma_{p-1}) = I'(\sigma) + X(\sigma)(1 - z_{k+1})^pAG = I'(\sigma)$$

by (2): since $X(\sigma)(1 - z_{k+1})^p = f((1 - z_{k+1})^p)$ and

$$(1 - z_{k+1})^p = (1 - z_{k+1}^p) = 0 \in A/p[G/H_k]. \quad (z_{k+1}^p \in H_k).$$

We thus have a filtration

$$I(\sigma) = I(\sigma_0) \supseteq I(\sigma_1) \supseteq \dots \supseteq I(\sigma_{p-1}) \supseteq I'(\sigma_{p-1}) = I'(\sigma); \quad (6)$$

and $I(\sigma_i) = I'(\sigma_{i-1})$ for $1 \leq i \leq p-1$.

Step 2B. For shortness in notation, we now write $K_1(I)$, $K_2(I)$ for $K_1(R, I)$, $K_2(R, I)$: R is always a quotient ring of AG . We are assuming that (3) holds for

the σ_i ; i.e., that

$$K_2(I(\sigma_i)/I'(\sigma_i)) = \langle \{g, 1 - X(\sigma_i)\lambda h\} : [g, h] \in H_{k+1}, \lambda \in A \rangle \quad (7)$$

for each $0 \leq i \leq p - 1$. Let $\{\lambda_1, \dots, \lambda_s\}$ be a \mathbb{Z}_p -basis for A . Let $h_1, \dots, h_t \in G$ be conjugacy class representatives $(\bmod H_{k+1})$ for those elements such that $[g_l, h_l] \in z_{k+1}H_k$ for some $g_l \in G$; fix also such g_l . Then (7) takes the form

$$K_2(I(\sigma_i)/I'(\sigma_i)) = M_i + \langle \{g_l, 1 - X(\sigma_i)\lambda_j h_l\} : 1 \leq j \leq s, 1 \leq l \leq t \rangle; \quad (8)$$

where

$$M_i = \langle \{g, 1 - \lambda X(\sigma_i)h\} : [g, h] \in H_k, \lambda \in A \rangle. \quad (9)$$

Step 2C. Now assume that $i < p - 1$; and consider the relative exact sequence

$$K_2(I(\sigma_i)/I'(\sigma_{i+1})) \rightarrow K_2(I(\sigma_i)/I'(\sigma_i)) \xrightarrow{\partial} K_1(I(\sigma_{i+1})/I'(\sigma_{i+1}))$$

(recall that $I'(\sigma_i) = I(\sigma_{i+1})$). By (2) (and [24, Corollary 2.6]):

$$K_1(I(\sigma_{i+1})/I'(\sigma_{i+1})) \cong H_0(G; A/p[G/H_{k+1}]), \quad (10)$$

where G acts by conjugation. Furthermore, for $1 \leq j \leq s$, $1 \leq l \leq t$,

$$\begin{aligned} \partial(\{g_l, 1 - X(\sigma_i)\lambda_j h_l\}) &= [g_l, 1 - X(\sigma_i)\lambda_j h_l] = 1 - X(\sigma_i)\lambda_j(g_l h_l g_l^{-1} - h_l) \\ &= 1 + X(\sigma_i)(1 - z_{k+1})\lambda_j h_l = 1 + X(\sigma_{i+1})\lambda_j h_l \pmod{I'(\sigma_{i+1})} \end{aligned}$$

(recall that $[g_l, h_l] \in z_{k+1}H_k$). By (10), these elements are all independent in $K_1(I(\sigma_{i+1})/I'(\sigma_{i+1}))$. So by (8) and (9),

$$\begin{aligned} \text{Im } [K_2(I(\sigma_i)/I'(\sigma_i)) \rightarrow K_2(I(\sigma_i)/I'(\sigma_i))] \\ &= \text{Im } [K_2(I(\sigma_i)/I'(\sigma_{i+1})) \rightarrow K_2(I(\sigma_i)/I'(\sigma_i))] \\ &= M_i = \langle \{g, 1 - \lambda X(\sigma_i)h\} : [g, h] \in H_k, \lambda \in A \rangle: \quad (11) \end{aligned}$$

all elements in M_i lift (using (2)) to $K_2(I(\sigma_i)/I'(\sigma)) \subseteq K_2(I(\sigma)/I'(\sigma))$.

Step 2D. By (8) and (11) (and (6)),

$$K_2(I(\sigma)/I'(\sigma)) = M + \langle \{g, 1 - \lambda X(\sigma)(1 - z_{k+1})^{p-1}h\} : [g, h] \in z_{k+1}H_k, \lambda \in A \rangle \quad (12)$$

where

$$\begin{aligned} M &= \langle \{g, 1 - \lambda X(\sigma)(1 - z_{k+1})^i h\} : 0 \leq i \leq p-1, [g, h] \in H_k, \lambda \in A \rangle \\ &= \langle \{g, 1 - \lambda X(\sigma)h\} : [g, h] \in H_k, \lambda \in A \rangle. \end{aligned}$$

(Note that $X(\sigma)^2 = 0$ in $I(\sigma)/I'(\sigma)$.) We want to show that $K_2(I(\sigma)/I'(\sigma)) = M$. Fix $\lambda \in A$ and $g, h \in G$ such that $[g, h] \in z_{k+1}H_k$, and set $u = 1 - X(\sigma)\lambda h$. Then

$$1 - X(\sigma)\lambda(1 - z_{k+1})^{p-1}h = \prod_{i=0}^{p-1} (1 - X(\sigma)\lambda z_{k+1}^i h) = \prod_{i=0}^{p-1} g^i u g^{-i} \in AG/I'(\sigma)$$

by (2) ($I(\sigma)/I'(\sigma) \cong A/p[G/H_k]$). So by Lemma 1.2,

$$\begin{aligned} \{g, 1 - x(\sigma)\lambda(1 - z_{k+1})^{p-1}h\} &= \{g, u \cdot gug^{-1} \cdots g^{p-1}ug^{1-p}\} \\ &= \{g^p, u\} + (p-1)\{u, u\} + \sum_{j=1}^{p-1} \{g^j ug^{-j}, u\}. \end{aligned}$$

By definition, $\{g^p, u\} \in M$. For any $0 \leq j \leq p-1$:

$$\begin{aligned} \{g^j ug^{-j}, u\} &= \{1 - X(\sigma)\lambda z_{k+1}^j h, 1 - X(\sigma)\lambda h\} \\ &= \left\{ 1 - (1 - z), 1 - X(\sigma) \frac{X(\sigma)}{1-z} \lambda^2 z_{k+1}^j h^2 \right\} \in M \end{aligned}$$

(see [17, Lemma 2.2] for the last step). So from (12) we now get that $K_2(I(\sigma)/I'(\sigma)) = M$; and this finishes the proof of (3).

Step 3. Now fix some $i \geq 1$. For any $0 \leq r \leq m-1$, (3) applied to $\sigma = (0; r, i)$ says that

$$\begin{aligned} K_2(p^r(1-z)^i AG / \langle p^{r+1}(1-z)^i, (1-z)^{i+1} \rangle) \\ = \langle \{g, 1 - \lambda p^r(1-z)^i h\} : [g, h] \in \langle z \rangle, \lambda \in A \rangle. \end{aligned}$$

For any such g , h , and λ , note that (in AG)

$$[g, 1 - \lambda p^r(1 - z)^i h] \equiv 0; 1 - \lambda p^r(1 - z)^i h \equiv (1 - \lambda(1 - z)^i h)^{p^r} \pmod{(1 - z)^{i+1}AG}.$$

It follows that

$$K_2((1 - z)^i AG / (1 - z)^{i+1} AG) = \langle \{g, 1 - \lambda(1 - z)^i h\} : [g, h] \in \langle z \rangle, \lambda \in A \rangle. \quad (13)$$

Step 4. The rest of the proof is analogous to Step 2B and 2C. Let $\lambda_1, \dots, \lambda_s$ be a \mathbb{Z}_p -basis for A , and let $h_1, \dots, h_t \in G$ be conjugacy class representatives for G/z . For $1 \leq l \leq t$, choose $g_l \in G$ so that $[g_l, h_l] = z^{q_l}$, and $1 \leq q_l \leq p^m = |z|$ is minimal. Then by (13),

$$K_2((1 - z)^i AG / (1 - z)^{i+1} AG) = N_i + \langle \{g_l, 1 - \lambda_j(1 - z)^i h_l\} : 1 \leq l \leq t, 1 \leq j \leq s \rangle, \quad (14)$$

where

$$N_i = \langle \{g, 1 - \lambda(1 - z)^i h\} : [g, h] = 1, \lambda \in A \rangle.$$

Consider the exact sequence

$$K_2\left(\frac{(1 - z)^i AG}{(1 - z)^{i+2} AG}\right) \rightarrow K_2\left(\frac{(1 - z)^i AG}{(1 - z)^{i+1} AG}\right) \xrightarrow{\partial} K_1\left(\frac{(1 - z)^{i+1} AG}{(1 - z)^{i+2} AG}\right). \quad (15)$$

For any j, l :

$$\partial(\{g_l, 1 - \lambda_j(1 - z)^i h_l\}) = [g_l, 1 - \lambda_j(1 - z)^i h_l] = 1 + q_l \lambda_j(1 - z)^{i+1} h_l.$$

By Lemma 1.3, these elements are independent in

$$K_1((1 - z)^{i+1} AG / (1 - z)^{i+2} AG) \cong H_0(G; A/p^m[G])$$

and have order p^m/q_l (q_l is a power of p). Furthermore, for each j and l , $[g_l^{p^m/q_l}, h_l] = 1$, and so

$$p^m/q_l \cdot \{g_l, 1 - \lambda_j(1 - z)^i h_l\} \in N_i.$$

So by (14), and the exactness of (15),

$$\text{Im} \left[K_2 \left(\frac{(1-z)^i AG}{(1-z)^{i+2} AG} \right) \rightarrow K_2 \left(\frac{(1-z)^i AG}{(1-z)^{i+1} AG} \right) \right] = \text{Ker} (\partial) = N_i.$$

Every element of N_i lifts to $K_2((1-z)^i AG) \subseteq K_2((1-z)AG)$. Thus, for any $i \geq 1$,

$$K_2((1-z)^i AG) = K_2((1-z)^{i+1} AG) + \langle \{g, 1 - \lambda(1-z)^i h\} : gh = hg, \lambda \in A \rangle. \quad (16)$$

By induction, for any $N > 1$,

$$K_2((1-z)AG) = K_2((1-z)^N AG) + \langle \{g, 1 - \lambda(1-z)^i h\} : gh = hg, \lambda \in A, 1 \leq i < N \rangle. \quad (17)$$

Let $p^k = \exp(G)$, and recall that $|z| = p^m$. Then $p(1-z) \mid (1-z)^{p^m}$, and so

$$1 + (1-z)^{(k+1)p^m} AG \subseteq 1 + p^{k+1}(1-z)AG \subseteq \{(1 + (1-z)\xi)^{p^k} : \xi \in AG\}.$$

Thus, for any commuting $h, g \in G$, any $\lambda \in A$, and any $i \geq (k+1)p^m$:

$$\{g, 1 - \lambda(1-z)^i h\} = \{g, (1 - (1-z)\xi)^{p^k}\} = \{g^{p^k}, 1 - (1-z)\xi\} = 0. \quad (\text{some } \xi \in AG).$$

By (16), for any $N > (k+1)p^m$, $K_2((1-z)^{(k+1)p^m} AG) = K_2((1-z)^N AG)$; and so

$$K_2((1-z)^{(k+1)p^m}) = \varprojlim_N K_2((1-z)^{(k+1)p^m} AG / (1-z)^N AG) = 0 \quad (18)$$

Equation (17) now takes the form

$$K_2((1-z)AG) = \langle \{g, 1 - \lambda(1-z)^i h\} : gh = hg, \lambda \in A, 1 \leq i < (k+1)p^m \rangle.$$

Furthermore, it suffices to take λ belonging to some $\hat{\mathbb{Z}}_p$ -basis for A . This shows that $K_2((1-z)AG)$ is generated by a finite set of elements of finite order, and is hence finite. \square

With some more work, one can in fact show that $K_2(AG, (1-z)AG)$ is

generated by symbols $\{g, 1 - \lambda(1 - z)h\}$, where $gh = hg$ in G and λ lies in any fixed $\hat{\mathbb{Z}}_p$ -basis for A .

One easy consequence of Theorem 1.4 is:

THEOREM 1.5. *For any prime p , any unramified p -ring A , and any p -group G , $K_2(AG)$ is finite.*

Proof. Fix some $1 \neq z \in Z(G)$. Then $K_2(AG, (1 - z)AG)$ is finite by Theorem 1.4. We may assume inductively that $K_2(A[G/z])$ is finite; and so $K_2(AG)$ is also finite. \square

In fact, using the results in [17], this can be extended to arbitrary finite G . Whether it is true for arbitrary \mathbb{Z}_p -orders, we do not know.

Section 2

Theorem 1.4 gives a set of generators for $\text{Ker}(K_2(A\alpha))$, when $\alpha: \tilde{G} \twoheadrightarrow G$ is a central extension of p -groups with cyclic kernel. In this section, we study $\text{Coker}(K_2(A\alpha))$ when $\text{Ker}(\alpha) \subseteq Z(\tilde{G})$. This problem was studied in [19]: $\text{Coker}(K_2(A\alpha))$ is described there for an arbitrary surjection α , but only up to a mysterious contribution by $H_3(G)$. What we show here is that the $H_3(G)$ contribution vanishes when α is a central extension.

PROPOSITION 2.1. *Let p be any prime, let A be an unramified p -ring, and let $\alpha: \tilde{G} \twoheadrightarrow G$ be any central extension of p -groups (i.e., $\text{Ker}(\alpha) \subseteq Z(\tilde{G})$). Then there is an exact sequence*

$$\begin{aligned} 0 \rightarrow \text{Coker}(H_2(\alpha)) &\xrightarrow{T_\alpha} \text{Coker}[K_2(A\alpha): K_2(A\tilde{G}) \rightarrow K_2(AG)] \\ &\xrightarrow{\Gamma_2^*(\alpha)} H_1(G; AG)/\langle g \otimes \lambda h : [\alpha^{-1}g, \alpha^{-1}h] = 1 \rangle. \end{aligned}$$

Here, T_α is included by the usual inclusion $H_2(G) \rightarrow K_2(AG)/\{-1, G\}$, and $\Gamma_2^*(\alpha)$ is induced by the homomorphism

$$\Gamma_2^*(G): K_2(AG) \rightarrow H_1(G; AG)/\langle g \otimes \lambda g^n : g \in G, \lambda \in A, n \in \mathbb{Z} \rangle$$

of [19, Theorem 3.6]. In particular, for any $g \in G$, $H = Z_G(g)$, and any

$u \in (AH)^*$,

$$\Gamma_2^*(\alpha)(\{g, u\}) = g \otimes \Gamma_H(u) \in H_1(G; AG)/\langle g \otimes \lambda h : [\alpha^{-1}g, \alpha^{-1}h] = 1 \rangle.$$

Proof. Define the group \hat{G} and the order \mathfrak{A} to be the pullbacks:

$$\begin{array}{ccc} \hat{G} & \xrightarrow{r_1} & \tilde{G} \\ \downarrow r_2 & & \downarrow \alpha \\ \tilde{G} & \xrightarrow{\alpha} & G \end{array} \quad \begin{array}{ccc} \mathfrak{A} & \xrightarrow{\hat{r}_1} & A\tilde{G} \\ \downarrow \hat{r}_2 & & \downarrow A\alpha \\ A\tilde{G} & \xrightarrow{A\alpha} & AG. \end{array}$$

Set

$$I_1 = \text{Ker } [A\hat{G} \xrightarrow{Ar_1} A\tilde{G}], \quad I_2 = \text{Ker } [A\hat{G} \xrightarrow{Ar_2} A\tilde{G}], \quad I = \text{Ker } [A\tilde{G} \xrightarrow{A\alpha} AG].$$

Then $\mathfrak{A} \cong A\hat{G}/(I_1 \cap I_2)$; and so by Lemma 2.4 in [16],

$$\mathfrak{A} \cong A\hat{G}/I_1 I_2.$$

Step 1. By [26, Theorem 4.1],

$$\text{tors}(K_1(A\hat{G})) \cong \mu_A \times \hat{G}^{ab} \times SK_1(A\hat{G}); \quad (1)$$

where μ_A denotes the group of roots of unity in A . We first claim that

$$\hat{G}^{ab} \rightarrow K_1(A\hat{G}/I_1 I_2) \cong K_1(\mathfrak{A}) \quad (2)$$

is injective. To see this, let $I(A\hat{G})$ denote the augmentation ideal of $A\hat{G}$. Then $I(A\hat{G})^2 \supseteq I_1 I_2$, and by [19, Proposition 2.2]:

$$A\hat{G}/I(A\hat{G})^2 \cong A \times (A \otimes \hat{G}^{ab}).$$

The isomorphism identifies $g \in \hat{G}^{ab}$ with $(1, 1 \otimes g)$, and so $\hat{G}^{ab} \subseteq K_1(A\hat{G}/I(A\hat{G})^2)$.

Now set $K = \text{Ker } (\alpha) \cong \text{Ker } (r_1)$, and consider the following diagram:

$$\begin{array}{ccccccc} H_2(\hat{G}) & \longrightarrow & H_2(\tilde{G}) & \xrightarrow{\delta'_{r_1}} & K & \longrightarrow & \hat{G}^{ab} \xrightarrow{H_1(r_1)} G^{ab} \longrightarrow 0 \\ \downarrow & & \downarrow H_2(\alpha) & & \downarrow Id & & \downarrow H_1(r_2) \\ H_2(\tilde{G}) & \xrightarrow{H_2(\alpha)} & H_2(G) & \xrightarrow{\delta^{\alpha}} & K & \longrightarrow & \tilde{G}^{ab} \longrightarrow G^{ab} \longrightarrow 0. \end{array}$$

The rows are the five-term exact sequences for the extensions $r_1: \hat{G} \rightarrow \tilde{G}$ and

$\alpha: \tilde{G} \twoheadrightarrow G$ (see [8, Corollary VI. 8.2]). It follows that

$$\text{Ker } [H_1(r_1 \times r_2): \hat{G}^{ab} \rightarrow \tilde{G}^{ab} \times \tilde{G}^{ab}] \cong \text{Coker } (H_2(\alpha)). \quad (3)$$

Furthermore, $\delta^{r_1} = \delta^\alpha \circ H_2(\alpha) = 0$, so $\text{Ker } (r_1) \cap [\hat{G}, \hat{G}] = 1$, and

$$SK_1(Ar_1): SK_1(A\hat{G}) \rightarrow SK_1(A\tilde{G}) \quad (4)$$

is injective by [15, Proposition 7].

Step 2. Now define

$$\Gamma_{AG}: K_1(A\hat{G}) \rightarrow H_0(\hat{G}; A\hat{G}); \quad \Gamma_{A\tilde{G}}: K_1(A\tilde{G}) \rightarrow H_0(\tilde{G}; A\tilde{G})$$

as in [20, Theorem 2.7], and recall that they are isomorphisms modulo torsion. By Theorem 1.1 in [19],

$$\Gamma_{A\hat{G}}(1 + I_1 I_2) = \text{Im } [I_1 I_2 \rightarrow H_0(\hat{G}; A\hat{G})]. \quad (5)$$

So $\Gamma_{A\hat{G}}$ induces a homomorphism

$$\Gamma_{\mathfrak{A}}: K_1(\mathfrak{A}) \rightarrow H_0(\hat{G}; \mathfrak{A}).$$

Consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mu_A \times \hat{G}^{ab} \times SK_1(A\hat{G}) & \longrightarrow & K_1(\mathfrak{A}) & \xrightarrow{\Gamma_{\mathfrak{A}}} & H_0(\hat{G}; \mathfrak{A}) \\ & & \downarrow f & & \downarrow K_1(r_1 \times r_2) & & \downarrow H_0(r_1 \times r_2) \\ 0 & \longrightarrow & [\mu_A \times \tilde{G}^{ab} \times SK_1(A\tilde{G})]^2 & \longrightarrow & [K_1(A\tilde{G})]^2 & \xrightarrow{\Gamma_{A\tilde{G}}} & [H_0(\tilde{G}; A\tilde{G})]^2 \end{array} \quad (6)$$

The bottom row is exact since $\text{Ker } (\Gamma_{A\tilde{G}}) = \text{tors } (K_1(A\tilde{G}))$. The top row is exact at $K_1(\mathfrak{A})$ since by (5), $\text{Ker } (\Gamma_{A\hat{G}}) \rightarrow \text{Ker } (\Gamma_{\mathfrak{A}})$ is onto. By (3) and (4), $\hat{G}^{ab} \cong \text{Ker } (f) \cong \text{Coker } (H_2(\alpha))$, and this injects into $K_1(\mathfrak{A})$ by (2). So the top row in (6) is exact, and there is an exact sequence

$$0 \rightarrow \text{Coker } (H_2(\alpha)) \rightarrow \text{Ker } (K_1(r_1 \times r_2)) \rightarrow \text{Ker } (H_0(r_1 \times r_2)). \quad (7)$$

By the Mayer–Victoris sequence for a pullback square,

$$\text{Ker } (K_1(r_1 \times r_2)) \cong \text{Coker } [K_2(A\alpha): K_2(A\tilde{G}) \rightarrow K_2(A\hat{G})]. \quad (8)$$

Step 3. The extension $0 \rightarrow I \rightarrow \mathfrak{A} \xrightarrow{r_1} AG \rightarrow 0$ is \hat{G} -equivariantly split by the diagonal map. Thus,

$$\text{Ker } [\hat{r}_{1*} : H_0(\hat{G}; \mathfrak{A}) \rightarrow H_0(\tilde{G}; A\tilde{G})] \cong H_0(\tilde{G}; I);$$

and so

$$\begin{aligned} \text{Ker } (H_0(r_1 \times r_2)) &\cong \text{Ker } [H_0(\tilde{G}; I) \rightarrow H_0(\tilde{G}; A\tilde{G})] \\ &\cong \text{Coker } [H_1(\tilde{G}; A\tilde{G}) \rightarrow H_1(\tilde{G}; AG)] \\ &\cong H_1(G; AG) / \langle g \otimes \lambda h : \lambda \in A, [\alpha^{-1}g, \alpha^{-1}h] = 1 \rangle. \end{aligned} \quad (9)$$

Upon substituting (8) and (9) into (7), we get the exact sequence

$$0 \rightarrow \text{Coker } (H_2(\alpha)) \xrightarrow{T_\alpha} \text{Coker } (K_2(A\alpha)) \xrightarrow{\Gamma_2^*(\alpha)} H_1(G; AG) / \langle g \otimes \lambda h : [\alpha^{-1}g, \alpha^{-1}h] = 1 \rangle.$$

That $\Gamma_2^*(\alpha)$ is the reduction of the map $\Gamma_2^*(G)$ of [19] follows since the constructions are identical. By diagram chasing, T_α is seen to be the reduction of the standard inclusion $H_2(G) \rightarrow K_2(AG) / \{-1, G\}$. \square

In fact, in the above situation, $\text{Im } (\Gamma_2^*(\alpha))$ can be described precisely with the help of Theorem 3.6 in [19].

Proposition 2.1 will be applied directly in Section 3, when describing $Cl_1(\mathbb{Z}G)$ for odd p -groups G . But we first note one consequence of particular interest. The next theorem is useful when constructing maps

$$\Gamma_2 : K_2(AG) \rightarrow H_1(G; AG) / \langle g \otimes \lambda g \rangle$$

for non-abelian p -groups G (compare with [21]).

THEOREM 2.2. *Let $\alpha : \tilde{G} \rightarrow G$ be any surjection of p -groups such that $\text{Ker } (\alpha) \cap [\tilde{G}, \tilde{G}] = 1$. Then for any unramified p -ring A , the map*

$$K_2(A\alpha) : K_2(A\tilde{G}) \rightarrow K_2(AG)$$

is onto, and its kernel is generated by elements of the form $\{g, 1 + (1 - z)^i h\}$ for $z \in \text{Ker } (\alpha)$, $i \geq 1$, and commuting $g, h \in G$.

Proof. Note first that

$$[\text{Ker}(\alpha), \tilde{G}] \subseteq \text{Ker}(\alpha) \cap [\tilde{G}, \tilde{G}] = 1;$$

so that $\text{Ker}(\alpha) \subseteq Z(\tilde{G})$. The exact sequence

$$H_2(\tilde{G}) \xrightarrow{H_2(\alpha)} H_2(G) \xrightarrow{\delta^\alpha} \text{Ker}(\alpha) \rightarrow \tilde{G}^{ab} \rightarrow G^{ab} \rightarrow 0$$

(see [8, Corollary VI. 8.2]) shows that $H_2(\alpha)$ is onto. By hypothesis,

$$H_1(G; AG)/\langle g \otimes \lambda h : [\alpha^{-1}g, \alpha^{-1}h] = 1 \rangle = 0:$$

commuting elements in G lift to commuting elements in \tilde{G} . So $K_2(A\alpha)$ is onto by Proposition 2.1.

Now write α as a composite

$$\alpha : \tilde{G} = G_0 \xrightarrow{\alpha_1} G_1 \xrightarrow{\alpha_2} G_2 \rightarrow \dots \xrightarrow{\alpha_n} G_n = G;$$

and so that $\text{Ker}(\alpha_j)$ is cyclic for all j . By Theorem 1.4,

$$\text{Ker}(K_2(A\alpha_j)) = \langle \{g, 1 + (1 - z)^i h\} : z \in \text{Ker}(\alpha_j), i \geq 1, g, h \in G_{j-1}, gh = hg \rangle$$

for each j . But all such symbols lift to $K_2(A\tilde{G})$; and so $\text{Ker}(K_2(A\alpha))$ is generated as described. \square

Section 3

We can now derive algorithms for computing the groups $Cl_1(\mathbb{Z}G)[\frac{1}{2}]$ and $SK_1(\mathbb{Z}G)[\frac{1}{2}]$ for finite G . The key extra tool when working with odd torsion is the standard involution on $K_n(\mathbb{Z}G)$ and $K_n(\hat{\mathbb{Z}}_p G)$; for example, this is what was used in [17] to construct natural splittings

$$SK_1(\mathbb{Z}G)[\frac{1}{2}] \cong Cl_1(\mathbb{Z}G)[\frac{1}{2}] \oplus \sum_{2 < p \mid |G|} SK_1(\hat{\mathbb{Z}}_p G).$$

Recall that for any group G and any commutative ring R , an antiinvolution $x \rightarrow \bar{x}$ on RG is defined by setting

$$\overline{\sum r_i g_i} = \sum r_i g_i^{-1} \quad (r_i \in R, g_i \in G).$$

This extends to an antiinvolution on $GL(RG)$ – defined by setting $\overline{(a_{ij})} = (\bar{a}_{ji})$ – and hence an involution on $K_1(RG)$. Similarly, an antiinvolution on $St(RG)$ is induced by setting $\overline{x_{ij}(a)} = x_{ji}(\bar{a})$ ($a \in RG$); and this restricts to an involution on $K_2(RG)$.

LEMMA 3.1. *For any group ring RG as above, and any commuting units, $u, v \in (RG)^*$, $\overline{\{u, v\}} = \{\bar{v}, \bar{u}\}$. In particular, for any $g \in G$, and $u \in (RG)^*$ such that $gu = ug$, $\overline{\{g, u\}} = \{g, \bar{u}\}$.*

Proof. Recall that $\{u, v\} = [X, Y]$, where $X, Y \in St(RG)$ are arbitrary liftings of $\text{diag}(u, u^{-1}, 1)$ and $\text{diag}(v, 1, v^{-1})$. Then

$$\overline{\{u, v\}} = \overline{[X, Y]} = \bar{Y}^{-1}\bar{X}^{-1}\bar{Y}\bar{X} = \{\bar{v}^{-1}, \bar{u}^{-1}\} = \{\bar{v}, \bar{u}\}.$$

The last statement follows since $\bar{g} = g^{-1}$. \square

The importance of the involution for simplifying the computation of $Cl_1(\mathbb{Z}G)$ follows from:

LEMMA 3.2. *For any odd prime p and any p -group G , the involution on $K_2(\hat{\mathbb{Q}}_p[G])_{(p)}$ is the identity.*

Proof. By [22, Section 2 and 3], for any p -group G and any irreducible $\mathbb{Q}G$ -module V , there are subgroups $K \triangleleft H \subseteq G$ and a faithful $\mathbb{Q}[H/K]$ -representation W such that $V = \text{Ind}_H^G(W)$, $\text{End}_{\mathbb{Q}H}(W) \cong \text{End}_{\mathbb{Q}G}(V)$, and H/K is cyclic. Let $A \subseteq \mathbb{Q}H$ and $B \subseteq \mathbb{Q}G$ denote the corresponding simple summands. Then the induction map restricts to a Morita equivalence from A to B , and hence induces an isomorphism of $K_2(\hat{\mathbb{Q}}_p \otimes_{\mathbb{Q}} A)$ to $K_2(\hat{\mathbb{Q}}_p \otimes_{\mathbb{Q}} B)$. Thus, if

$$S = \{(H, K) : K \triangleleft H \subseteq G, H/K \text{ cyclic}\},$$

then the map

$$\sum \text{Ind}_{H/K}^G : \sum_{(H, K) \in S} K_2(\hat{\mathbb{Q}}_p[H/K]) \rightarrow \hat{\mathbb{Q}}_p[G] \quad (1)$$

is onto. Here, $\text{Ind}_{H/K}^G$ is the composite

$$\text{Ind}_{H/K}^G : K_2(\hat{\mathbb{Q}}_p[H/K]) \xrightarrow{\text{incl}} K_2(\hat{\mathbb{Q}}_p H) \xrightarrow{\text{Ind}_H^G} K_2(\hat{\mathbb{Q}}_p G);$$

where the first map is induced by the inclusion of $\hat{\mathbb{Q}}_p[H/K]$ as a direct summand of $\hat{\mathbb{Q}}_pH$.

The $\text{Ind}_{K/H}^G$ commute with the involution, and so by (1) it suffices to prove the lemma when G is cyclic. If $G \cong C_{p^n}$, write $\hat{\mathbb{Q}}_pG \cong \prod_{i=0}^n F_i$, where $F_i \cong \hat{\mathbb{Q}}_p[\zeta_{p^i}]$ (a field). For each i , the involution inverts elements in μ_{F_i} . So from the isomorphism $K_2(F_i) \cong \mu_{F_i}$ and its naturality with respect to automorphisms of F_i , we get that $\{\bar{u}, \bar{v}\} = -\{u, v\} = \{v, u\}$ for $u, v \in F_i^*$. But $\{\bar{n}, \bar{v}\} = \{\bar{v}, \bar{u}\}$ by Lemma 3.1, and so the involution on $K_2(F_i)$, and hence on $K_2(\hat{\mathbb{Q}}_pG)$, is trivial. \square

In fact, Lemma 3.2 also holds for 2-groups, and for arbitrary finite G if $K_2(\tilde{\mathbb{Q}}_pG)_{(p)}$ is replaced by $C_p(\mathbb{Q}G)$ (see the definition in the introduction).

The main problem when describing $Cl_1(\mathbb{Z}G)$ for a p -group G is computing the image of $K_2(\hat{\mathbb{Z}}_pG)$ in $K_2(\hat{\mathbb{Q}}_pG)$. Lemma 3.2 shows that when p is odd, it is enough to concentrate attention on $K_2(\hat{\mathbb{Z}}_pG)^+$; and (recall the formula $\{\bar{g}, \bar{u}\} = \{g, \bar{u}\}$) on $K_1(\hat{\mathbb{Z}}_pG)^+$.

PROPOSITION 3.3. *For any odd prime p , any unramified p -ring A , and any p -group G , Γ_{AG} restricts to an isomorphism*

$$\Gamma_{AG}^+: K_1(AG)^+ \rightarrow H_0(G; AG)^+.$$

Proof. By [20, Theorem 2.7], there is an exact sequence

$$0 \rightarrow G^{ab} \times SK_1(AG) \rightarrow K_1(AG) \xrightarrow{\Gamma_{AG}} H_0(G; AG) \xrightarrow{\omega} G^{ab} \rightarrow 0 \quad (1)$$

where $\omega(\sum \lambda_i g_i) = \prod g_i^{\text{Tr}(\lambda_i)}$. These maps all commute with the involution; and $(G^{ab})^+ = 0$ by definition. That $SK_1(AG)^+ = 0$ follows from the definition of the isomorphism

$$\Theta_{AG}: SK_1(AG) \rightarrow H_2(G)/H_2^{ab}(G)$$

in [15, diagram on p. 215]. So (1) restricts to an isomorphism

$$\Gamma_{AG}^+: K_1(AG)^+ \rightarrow H_0(G; AG)^+. \quad \square$$

If A is an unramified p -ring, and G is an *abelian* p -group, we can now define for any $\lambda \in A$ and $g \in G$ a unit $u(\lambda g) \in (AG)^{**} \cong K_1(AG)^+$ to be the unique element such that $\Gamma_G^+(u(\lambda g)) = \frac{1}{2}\lambda(g + g^{-1})$. If G is an arbitrary p -group and $g \in G$, we let $u(\lambda g) \in (AG)^*$ be the image of $u(\lambda g) \in (AH)^*$, when $H = \langle g \rangle$.

The results of Sections 1 and 2 can now be used to describe $K_2(\hat{\mathbb{Z}}_p G)^+$:

PROPOSITION 3.4. *For any odd prime p , any unramified p -ring A , and any p -group G ,*

$$K_2(AG)^+ = \langle \{g, u(\lambda h)\} : \lambda \in A, g, h \in G, [g, h] = 1 \rangle.$$

Proof. For any G , define an involution on $H_1(G; AG)$ by setting $\overline{g \otimes \lambda h} = g \otimes \lambda h^{-1}$. Define

$$\Delta_G^+ : H_1(G; AG)^+ \rightarrow K_2(AG)^+$$

by setting $\Delta_G^+(g \otimes \frac{1}{2}\lambda(h + h^{-1})) = \{g, u(\lambda h)\}$ for any $\lambda \in A$ and commuting $g, h \in G$.

Fix some G , choose $z \in Z(G)$ of order p , set $H = G/z$, and let $\alpha : G \twoheadrightarrow H$ be the projection. Assume inductively that Δ_H^+ is surjective, and consider the following diagram:

$$\begin{array}{ccccccc} 0 \rightarrow \text{Ker } (H_1(A\alpha))^+ \rightarrow H_1(G; AG)^+ \xrightarrow{H_1(A\alpha)} H_1(H; AH)^+ \rightarrow \text{Coker } (H_1(A\alpha))^+ \rightarrow 0 \\ \downarrow f_1 \qquad \qquad \downarrow \Delta_G^+ \qquad \qquad \downarrow \Delta_H^+ \qquad \qquad \Gamma_2^+ \uparrow \downarrow f_2 \\ 0 \rightarrow \text{Ker } (K_2(A\alpha))^+ \rightarrow K_2(AG)^+ \xrightarrow{K_2(A\alpha)} K_2(AH)^+ \rightarrow \text{Coker } (K_2(A\alpha))^+ \rightarrow 0 \end{array} \quad (1)$$

Here, f_1 and f_2 are induced by Δ_G^+ and Δ_H^+ , and Γ_2^+ is the restriction of the homomorphism of Proposition 2.1. For any $\lambda \in A$ and commuting $g, h \in G$,

$$\Gamma_2^+ \circ f_2(g \otimes \frac{1}{2}\lambda(h + h^{-1})) = \Gamma_2^+(\{g, u(h)\}) = g \otimes \Gamma_{AG}(u(h)) = g \otimes \frac{1}{2}\lambda(h + h^{-1});$$

and so f_2 is injective. By Theorem 1.4,

$$\begin{aligned} \text{Ker } (K_2(A\alpha))^+ \\ = \langle \{g, (1 - \lambda(1 - z)^i h)(1 - \lambda(1 - z^{-1})^i h^{-1})\} : \lambda \in A, i \geq 1, gh = hg \rangle; \end{aligned}$$

and so by Proposition 3.3 (applied to the $K_1(A[Z_G(g)])^+$):

$$\text{Ker } (K_2(A\alpha))^+ \subseteq \langle \{g, u(\lambda h)\} : \lambda \in A, [g, h] = 1 \rangle = \text{Im } (\Delta_G^+).$$

By diagram chasing in (1), Δ_G^+ is now seen to be onto. \square

It seems quite likely that the homomorphisms Δ_G^+ defined above actually induce isomorphisms

$$HC_1(AG)^+ \cong [H_1(G; AG)/\langle g \otimes \lambda g \rangle]^+ \cong K_2(AG)^+.$$

This is the case at least for abelian p -groups [21, Theorem 3.9].

It remains only to find a description of the image of any $\{g, u(h)\}$ in $K_2(\hat{\mathbb{Q}}_p G)$, when $p > 2$ and G is a p -group. Recall that $K_2(\hat{\mathbb{Q}}_p G)$ is described in terms of norm residue symbol isomorphisms

$$(\cdot, \cdot)_F : K_2(F) \xrightarrow{\cong} \mu_F$$

defined for any finite extension F of $\hat{\mathbb{Q}}_p$ [12, Theorem A.14].

LEMMA 3.5. *Fix an odd prime p and a p -group G ; and let $u(g) \in (\hat{\mathbb{Z}}_p G)^*$ for $g \in G$ be defined as above. Write*

$$\hat{\mathbb{Q}}_p G = \prod_{i=1}^k B_i; \quad B_i = M_{r_i}(F_i),$$

where for each i , $F_i \cong \hat{\mathbb{Q}}_p \zeta_{p^m}$ (a field) for some $m \geq 0$ (see [22]). Let

$$\lambda_G : K_2(\hat{\mathbb{Q}}_p G) \rightarrow \prod_{i=1}^k (\mu_{F_i})_p$$

be the product of the norm residue symbol homomorphisms

$$\lambda_G^i : K_2(B_i) \cong K_2(M_{r_i}(F_i)) \cong K_2(F_i) \xrightarrow{(\cdot, \cdot)} (\mu_{F_i})_p.$$

For each i , let V_i be the irreducible B_i -representation. Then, for any commuting $g, h \in G$,

$$\lambda_G(\{g, u(h)\}) = [\det_{F_i}(g, V_i^h)]_{i=1}^k. \quad (V_i^h = \{x \in V_i : hx = x\}).$$

Proof. Fix some i , set $B = B_i$, $V = V_i$, $F = F_i$, $r = r_i$; and let

$$\alpha : \hat{\mathbb{Q}}_p G \twoheadrightarrow B \cong \text{End}_F(V) \cong M_r(F)$$

be the projection. Let m be such that $F \cong \hat{\mathbb{Q}}_p \zeta_{p^m}$. Set $p^m = \exp(G)$, and let

$$f : B \cong M_r(F) \rightarrow M_r(\hat{\mathbb{Q}}_p \zeta_{p^m})$$

be an inclusion. Note that taking norm residue symbols commutes (p is odd) with inclusions of cyclotomic fields: this follows, for example, from the formulas in [2].

Fix commuting $g, h \in G$. Then $\langle g, h \rangle$ is an abelian group of exponent dividing p^n ; and so $f\alpha(g)$ and $f\alpha(h)$ are conjugate (simultaneously) to diagonal matrices:

$$f\alpha(g) \sim \text{diag}(u_1, \dots, u_r), f\alpha(h) \sim \text{diag}(v_1, \dots, v_r) \quad (u_l, v_l \in \langle \zeta_{p^m} \rangle).$$

with

$$u(h) = \sum_j \lambda_j h^j; \quad (\lambda_j \in \hat{\mathbb{Z}}_p)$$

so that

$$K_2(f\alpha)(\{g, u(h)\}) = \prod_{l=1}^r \left\{ u_l, \sum_j \lambda_j v_l^j \right\}.$$

By the formulas of Artin and Hasse [2],

$$\lambda_G^i(\{g, u(h)\}) = \prod_{l=1}^r \left(u_l, \sum_j \lambda_j v_l^j \right)_F = \prod_{l=1}^r u_l^{N_l};$$

where

$$N_l = \frac{1}{p^n} \text{Tr} \left(\log \left(\sum_j \lambda_j v_l^j \right) \right). \quad (\text{Tr} : \hat{\mathbb{Q}}_p \zeta_{p^m} \rightarrow \hat{\mathbb{Q}}_p)$$

Recall that $\Gamma_G(u(h)) = \frac{1}{2}(h + h^{-1})$, where $\Gamma_G = (1 - (1/p)\Phi) \circ \log$, and $\Phi(\sum \lambda_i g_i) = \sum \lambda_i g_i^p$. Thus,

$$\begin{aligned} \log(u(h)) &= \left(1 - \frac{1}{p} \Phi \right)^{-1} \left(\frac{1}{2}(h + h^{-1}) \right) \\ &= \frac{p}{p-1} + \frac{1}{2} \left[(h + h^{-1} - 2) + \frac{1}{p} (h^p + h^{-p} - 2) + \dots \right]. \end{aligned}$$

Hence, for $1 \leq l \leq r$,

$$\begin{aligned} N_l &= \frac{1}{p^n} \text{Tr} \left(\frac{p}{p-1} + \frac{1}{2} \left[(v_l + v_l^{-1} - 2) + \frac{1}{p} (v_l^p + v_l^{-p} - 2) + \dots \right] \right) \\ &= \begin{cases} 1 & \text{if } v_l = 1 \\ 0 & \text{if } v_l \neq 1. \end{cases} \quad (v_l \in \langle \zeta_{p^n} \rangle) \end{aligned}$$

It follows that

$$\lambda_G^i(\{g, u(h)\}) = \prod_{v_l=1} u_l = \det_F(g, V^h). \quad \square$$

The main result can now be shown.

THEOREM 3.6. *Let p be an odd prime, and let G be a p -group. Write $\mathbb{Q}G = \prod_{i=1}^k B_i$, where each B_i is a matrix algebra over a field F_i with irreducible representation V_i . Define*

$$\psi_G : H_1(G; \mathbb{Z}G) \rightarrow \prod_{i=1}^k (\mu_{F_i})_p$$

by setting, for any commuting $g, h \in G$,

$$\psi_G(g \otimes h) = [\det_{F_i}(g, V_i^h)]_{i=1}^k.$$

Then $Cl_1(\mathbb{Z}G) \cong \text{Coker } (\psi_G)$ and

$$SK_1(\mathbb{Z}G) \cong \text{Coker } (\psi_G) \oplus (H_2(G)/H_2^{ab}(G)).$$

More precisely, there is a commutative square

$$\begin{array}{ccc} K_2(\hat{\mathbb{Q}}_p G)_{(p)} & \xrightarrow{\lambda_G} & \prod_{i=1}^k (\mu_{F_i})_p \\ \partial \downarrow & & \downarrow \text{proj} \\ Cl_1(\mathbb{Z}G) & \xrightarrow{\Delta_G} & \text{Coker } (\psi_G); \end{array}$$

where λ_G is induced by the norm residue symbol, and ∂ is the boundary map in the localization sequence.

Proof. By [20, Theorem 2.1 and 2.2], there is an exact sequence

$$K_2(\hat{\mathbb{Z}}_p G) \xrightarrow{\varphi_G} \text{Coker} \left[K_2 \left(\mathbb{Z} \left[\frac{1}{p} \right] [G] \right) \rightarrow K_2(\hat{\mathbb{Q}}_p G) \right] \xrightarrow{\partial} Cl_1(\mathbb{Z} G) \rightarrow 0$$

and an isomorphism

$$\text{Coker} \left[K_2 \left(\mathbb{Z} \left[\frac{1}{p} \right] [G] \right) \rightarrow K_2(\hat{\mathbb{Q}}_p G) \right] \cong K_2(\hat{\mathbb{Q}}_p G)_{(p)} \xrightarrow{\lambda_G} \prod_{i=1}^k (\mu_{F_i})_p.$$

(note that $\mathbb{Z}[1/p][G]$ is a maximal order). Consider the diagram

$$\begin{array}{ccccccc} H_1(G; \mathbb{Z} G)^+ & \xrightarrow{\psi_G^+} & \prod_{i=1}^k (\mu_{F_i})_p & \xrightarrow{\text{proj}} & \text{Coker}(\psi_G) & \longrightarrow 0 \\ \downarrow \Delta_G^+ & (1) & \uparrow \cong \lambda_G & (2) & \uparrow \Lambda_G & & \\ K_2(\hat{\mathbb{Z}}_p G)^+ & \xrightarrow{\varphi_G^+} & K_2(\hat{\mathbb{Q}}_p G)_{(p)} & \xrightarrow{\partial} & Cl_1(\mathbb{Z} G) & \longrightarrow 0. \end{array}$$

By Lemma 3.2, $\text{Im}(\varphi_G^+) = \text{Im}(\varphi_G)$; and $\text{Im}(\psi_G^+) = \text{Im}(\psi_G)$ since $\psi_G(g \otimes h) = \psi_G(g \otimes h^{-1})$ by definition. So the rows above are exact. The map Δ_G^+ , defined by setting $\Delta_G^+(g \otimes h) = \{g, u(h)\}$, is onto by Proposition 3.4, and (1) commutes by Lemma 3.5. So there is a unique isomorphism

$$\Lambda_G : Cl_1(\mathbb{Z} G) \xrightarrow{\cong} \text{Coker}(\psi_G)$$

which makes (2) commute.

The exact sequence

$$0 \rightarrow Cl_1(\mathbb{Z} G) \rightarrow SK_1(\mathbb{Z} G) \rightarrow SK_1(\hat{\mathbb{Z}}_p G) \rightarrow 0$$

is naturally split by [17, Theorem 4.8], and

$$SK_1(\hat{\mathbb{Z}}_p G) \cong H_2(G)/H_2^{ab}(G) \cong H_2(G)/\langle g \wedge h : g, h \in G, gh = hg \rangle$$

by [15, Theorem 3]. So

$$SK_1(\mathbb{Z} G) \cong \text{Coker}(\psi_G) \oplus (H_2(G)/H_2^{ab}(G)). \quad \square$$

In [17, Theorem 4.8], the computation of $Cl_1(\mathbb{Z} G)_{(p)}$ for odd p and arbitrary

finite G was reduced to the case of a p -group. More precisely, if C_1, \dots, C_k are conjugacy class representatives for cyclic subgroups in G of order prime to p , and $N_i = N_G(C_i)$, $Z_i = Z_G(C_i)$, and $\mathfrak{P}(Z_i)$ is the set of p -subgroups, then

$$Cl_1(\mathbb{Z}G)_{(p)} \cong \sum_{i=1}^k H_0\left(N_i/Z_i; \varinjlim_{H \in \mathfrak{P}(Z_i)} Cl_1(\mathbb{Z}H)\right). \quad (3.7)$$

Here, the limits are taken with respect to inclusion and conjugation among subgroups.

This direct sum decomposition is somewhat awkward, and hence a more direct description of $Cl_1(\mathbb{Z}G)_{(p)}$ seems also desirable. In fact, one can define homomorphisms

$$\psi_G : H_1(G; \mathbb{Z}G) \rightarrow \prod_{i=1}^k (\mu_{F_i}), \quad \left(\mathbb{Q}G \cong \prod_{i=1}^k B_i, F_i = Z(B_i) \right)$$

for arbitrary finite G , such that $Cl_1(\mathbb{Z}G)[\frac{1}{2}] \cong \text{Coker } (\psi_G)[\frac{1}{2}]$. But alone the definition of ψ_G become quite complicated as soon as we start working with non- p -groups; and the most efficient way of describing $Cl_1(\mathbb{Z}G)[\frac{1}{2}]$ for concrete G does seem to be by means of (3.7) above, together with Theorem 3.6. Some techniques for calculating with the help of (3.7) are presented in [17, Section 5].

Section 4

Theorem 3.6 reduces the calculation of $Cl_1(\mathbb{Z}G)$, for an odd order p -group G , to a straightforward combinatorial algorithm. We now give some examples to illustrate how this works in practice. Examples of calculations for abelian G are presented in [1]; and for non-abelian G of order p^3 , $Cl_1(\mathbb{Z}G)$ is calculated in [19, Theorem 7.5] using a weaker form of the theorem. So here we take some non-abelian groups of order p^4 to give a sample of some of the techniques which can be used. Throughout this section, p denotes a fixed odd prime.

Note first that for any p -group G and any commuting $g, h \in G$,

$$\psi_G(g \otimes g) = 0; \quad \psi_G(g \otimes h^n) = \psi_G(g \otimes h) \quad (\text{if } p \nmid n);$$

and

$$\psi_G(aga^{-1} \otimes aha^{-1}) = \psi_G(g \otimes h) \quad (\text{any } a \in G).$$

Thus, when describing $\text{Im}(\psi_G)$, it suffices to consider $\psi_G(g \otimes h)$ as h runs through a set of \mathbb{Q} -conjugacy class representatives in G , and g a set of generators for $Z_G(h)/h$.

An irreducible representation V of G will be described by listing eigenvalues for the actions of various group elements on V – or, when necessary, by describing the irreducible components of $V|H$ for some appropriate $H \subseteq G$.

Finally, note that when $|G| = p^4$, then $SK_1(\hat{\mathbb{Z}}_p G) = 0$ by [15, Proposition 23]. So $SK_1(\mathbb{Z}G) = Cl_1(\mathbb{Z}G)$ in this case.

PROPOSITION 4.1. *Assume $G \cong H \times C_p$, where H is non-abelian, $|H| = p^3$, and $\exp(H) = p$. Then*

$$SK_1(\mathbb{Z}G) \cong (\mathbb{Z}/p)^{(p^2+3p-6)/2}.$$

Proof. Fix generators $a, b \in H$ and $c \in C_p$; and set $z = [a, b]$. Then $Z(G) = \langle z, c \rangle$, and for any $g \in G \setminus Z(G)$, $Z_G(g) = \langle Z(G), g \rangle$. Set $\zeta = \zeta_p$, and note that

$$\mathbb{Q}[G] \cong \mathbb{Q} \times \prod^{p^2+p+1} \mathbb{Q}[\zeta] \times \prod^p M_p(\mathbb{Q}[\zeta]).$$

The following table describes $\psi = \psi_G$. Here, $(H^{ab})^*$ denotes the set of irreducible complex characters of H^{ab} , and $(*)$ for eigenvalues means that all powers of ζ occur.

Representation	$U \cong \mathbb{Q}\zeta$	$V_m \cong \mathbb{Q}\zeta$	$W_\chi \cong \mathbb{Q}\zeta$	$X_m \cong (\mathbb{Q}\zeta)^p$
Indexed by	—	$0 \leq m < p$	$\chi \in (H^{ab})^*$	$0 \leq m < p$
$E'\text{val } (a, b, c, z)$	$(\zeta, 1, 1, 1)$	$(\zeta^m, \zeta, 1, 1)$	$(\chi(a), \chi(b), \zeta, 1)$	$(*, *, \zeta^m, \zeta)$
$\psi(a \otimes cz^{-i})$	ζ	ζ^m	1	1
$\psi(b \otimes cz^{-i})$	1	ζ	1	1
$\psi(a \otimes (1 - c))$	1	1	$\chi(a)$	1
$\psi(b \otimes (1 - c))$	1	1	$\chi(b)$	1
$\psi(c \otimes 1)$	1	1	ζ	1
$\psi(G \otimes (1 - z))$	1	1	1	1
$\psi(z \otimes gc^{-i})$	1	1	1	ζ
$\psi(c \otimes gc^{-i})$ ($g \in H \setminus \langle z \rangle$)	1	1	ζ (if $\chi(g) = \zeta^i$) 1 (if $\chi(g) \neq \zeta^i$)	ζ^m
$\psi(c \otimes \xi)$	1	1	1	ζ^m

Here, in the last line, $\xi = a(1 + b + \cdots + b^{p-1}) - b(c + c^2 + \cdots + c^{p-1})$. By inspection,

$$SK_1(\mathbb{Z}G) \cong \text{Coker } (\psi) \cong (\mathbb{Z}/p)^{p-1} \oplus (\mathbb{Z}/p[C_p \times C_p]/I) \oplus (\mathbb{Z}/p)^{p-2}, \quad (1)$$

where $I \subseteq \mathbb{Z}/p[C_p \times C_p]$ is the ideal generated by elements $(\sum_{g \in K} g)$ for subgroups $K \subseteq C_p \times C_p$ of order p .

Write $C_p \times C_p = \langle g \rangle \times \langle h \rangle$, and let $J = \langle 1 - g, 1 - h \rangle \subseteq \mathbb{Z}/p[C_p \times C_p]$ denote the Jacobson radical. Then

$$I = \langle (1 - g)^{p-1}, (1 - h)^{p-1}; (1 - g^i h)^{p-1} : 1 \leq i \leq p - 1 \rangle.$$

Furthermore, for any $1 \leq i \leq p - 1$:

$$(1 - g^i h) = 1 - [1 - (1 - g)]^i [1 - (1 - h)] \equiv i(1 - g) + (1 - h) \pmod{J^2}$$

and so

$$\begin{aligned} (1 - g^i h)^{p-1} &\equiv \sum_{k=0}^{p-1} \binom{p-1}{k} i^k (1 - g)^k (1 - h)^{p-1-k} \\ &= \sum_{k=0}^{p-1} (-i)^k (1 - g)^k (1 - h)^{p-1-k} \pmod{J^p}. \end{aligned}$$

The determinant of $\{(-i)^k\}_{i,k=1}^{p-2}$ is invertible over \mathbb{Z}/p (a van der Monde determinant), and so

$$I + J^p = \langle (1 - g)^k (1 - h)^{p-1-k} : 0 \leq k \leq p - 1 \rangle = J^{p-1}.$$

But $J^{2p-1} = 0$, and hence this implies that $I = J^{p-1}$. So as a group,

$$\mathbb{Z}/p[C_p \times C_p]/I \cong (\mathbb{Z}/p)^{1/2p(p-1)}$$

$$\text{with basis } \{(1 - g)^i (1 - h)^j : i, j \geq 0, i + j < p - 1\}.$$

The result now follows from (1). \square

In the above example, the fact that $[G, G]$ was central helped to keep the description of ψ_G simple. The next example illustrates additional complexities which can arise when this is no longer the case. First, a lemma is needed.

LEMMA 4.2. *Let G be cyclic of order p^n ($n \geq 1$) with generator $g \in G$.*

Then, for any

$$0 \neq \alpha = \sum_{i=0}^{p^n-1} a_i g^i \in \mathbb{Z}/p[G] \quad (a_i \in \mathbb{Z}/p);$$

$\mathbb{Z}/p[G]/(\alpha) \cong (\mathbb{Z}/p)^k$ (as groups), where

$$k = \min \left\{ m \geq 0 : \sum_{i=0}^{p^n-1} \binom{i}{m} a_i \neq 0 \text{ in } \mathbb{Z}/p \right\}.$$

Proof. By direct calculation,

$$\begin{aligned} \alpha &= \sum_{i=0}^{p^n-1} a_i g^i = \sum_{i=0}^{p^n-1} a_i (1 + (g-1))^i = \sum_{i=0}^{p^n-1} a_i \sum_{m=0}^i \binom{i}{m} (g-1)^m \\ &= \sum_{m=0}^{p^n-1} \left(\sum_{i=0}^{p^n-1} \binom{i}{m} a_i \right) (g-1)^m. \end{aligned} \tag{1}$$

Recall that $\mathbb{Z}/p[G]$ is a local ring with maximal ideal generated by $(g-1)$. So if k is defined as above, then $\alpha = (g-1)^k u$ for some unit u in $\mathbb{Z}/p[G]$, and

$$rk[\mathbb{Z}/p[G]/(\alpha)] = rk[\mathbb{Z}/p[G]/(g-1)^k] = k. \quad \square$$

PROPOSITION 4.3. Set $H = \langle a \rangle \times \langle b \rangle \times \langle c \rangle = C_p^3$, $K = \langle x \rangle \cong C_p$, and let G be any extension of the form

$$1 \rightarrow H \rightarrow G \rightarrow K \rightarrow 1$$

such that

$$xax^{-1} = ab, \quad xbx^{-1} = bc, \quad xcx^{-1} = c.$$

Then

$$SK_1(\mathbb{Z}G) \cong Cl_1(\mathbb{Z}G) \cong (\mathbb{Z}/p)^{3(p-1)/2}.$$

Proof. The action of x on $\mathbb{Q}H$ fixes $\mathbb{Q}[H/\langle b, c \rangle]$, and permutes the other $p^2 + p$ summands freely. Thus,

$$\mathbb{Q}G \cong \mathbb{Q}[G^{ab}] \times \prod_{i=1}^{p+1} M_p(\mathbb{Q}[\zeta]);$$

where $\zeta = \zeta_p$. The following table presents ψ_G , where the nonabelian representations are described by their restrictions to H :

Representation	$U \cong \mathbb{Q}\zeta$	$V_m \cong \mathbb{Q}\zeta$	$W \cong (\mathbb{Q}\zeta)^p$	$X_m \cong (\mathbb{Q}\zeta)^p$
Indexed by	—	$0 \leq m < p$	—	$0 \leq m < p$
E'val $(a, b, c; x)$	$(1, 1, 1; \zeta)$	$(\zeta, 1, 1; \zeta^m)$	$(\zeta^r, \zeta, 1)$ $(1 \leq r \leq p)$	$(\zeta^{m+1/2r(r-1)}, \zeta^r, \zeta)$ $(1 \leq r \leq p)$
$\psi(a \otimes c)$	1	ζ	1	1
$\psi(x \otimes c)$	ζ	ζ^m	1	1
$\psi(a \otimes (1 - c))$	1	1	1	ζ^T
$\psi(x \otimes (1 - c))$	1	1	1	$\zeta^u (x^p = c^u)$
$\psi(a \otimes (b - c))$	1	1	1	ζ^m
$\psi(c \otimes b)$	1	1	1	ζ
$\psi(b \otimes ac^{-i})$	1	1	ζ	$\zeta^{R(i-m)}$
$\psi(c \otimes ac^{-i})$	1	1	1	$\zeta^{S(i-m)}$
$\psi(c \otimes gx) (g \in H)$	1	1	1	$\begin{cases} \zeta & ((gx)^p = 1) \\ 1 & ((gx)^p \neq 1). \end{cases}$

Here, $T = \sum_{r=1}^p \frac{1}{2}r(r-1)$;

$$R(i) = \sum \{r: 1 \leq r \leq p, \frac{1}{2}r(r-1) \equiv i \pmod{p}\};$$

$$S(i) = \#\{r: 1 \leq r \leq p, \frac{1}{2}r(r-1) \equiv i \pmod{p}\}.$$

Note that solutions to $\frac{1}{2}r(r-1) \equiv i$ come in pairs $\{r, p+1-r\}$ (unless $r = (p+1)/2$). This shows that for all i ,

$$R(i) = \frac{p+1}{2} S(i) \equiv \frac{1}{2} S(i) \pmod{p}.$$

Identify $\prod_{X_m} \langle \zeta \rangle$ with $\mathbb{Z}/p[C_p]$, by identifying X_m with g^m for some generator g of C_p . Then

$$SK_1(\mathbb{Z}G) \cong \text{Coker } (\psi_G)$$

$$\cong (\mathbb{Z}/p)^{p-1} \oplus \left(\mathbb{Z}/p[C_p] / \left\langle \sum_m g^m, \sum_m mg^m, \sum_m S(i-m)g^m \pmod{p} \right\rangle \right)$$

$$\cong (\mathbb{Z}/p)^{p-1} \oplus \mathbb{Z}/p[C_p]/I,$$

where I is the ideal generated by

$$\alpha = \sum_m S(m)g^{-m} = \sum_{k=1}^p g^{-1/2k(k-1)}.$$

By Lemma 4.2, we will be done upon showing that

$$\sum_{k=1}^p \binom{\frac{1}{2}k(k-1)}{n} \begin{cases} \equiv 0 \pmod{p} & \text{for } 0 \leq n < \frac{p-1}{2} \\ \not\equiv 0 \pmod{p} & \text{for } n = \frac{p-1}{2} \end{cases} \quad (1)$$

But the sum is a polynomial in k (over \mathbb{Z}/p) of degree exactly $2n$; and (1) follows since

$$p-1 = \min \left\{ m \geq 0 : \sum_{k=1}^p k^m \not\equiv 0 \pmod{p} \right\}. \quad \square$$

The groups covered above turn out to be the most difficult cases for computing $SK_1(\mathbb{Z}G)$ when $|G| = p^4$. In fact, all other groups of order p^4 are covered by the following proposition (this can easily be checked directly, but also follows from the classification in [9, section III.12]).

PROPOSITION 4.4. *Assume that G is non-abelian of order p^4 , and that there is a subgroup $H \triangleleft G$ such that $H \cong C_{p^3}$ or $H \cong C_{p^2} \times C_p$. Then*

$$\begin{aligned} SK_1(\mathbb{Z}G) &= Cl_1(\mathbb{Z}G) \cong (\mathbb{Z}/p)^{p-1} & \text{if } G^{ab} \cong C_p \times C_p \\ &\cong (\mathbb{Z}/p)^{2(p-1)} & \text{if } G^{ab} \cong C_{p^2} \times C_p \\ &\cong (\mathbb{Z}/p)^{(p+2)(p-1)/2} & \text{if } G^{ab} \cong C_p \times C_p \times C_p. \end{aligned}$$

Proof. Write

$$\mathbb{Q}G = \mathbb{Q}[G^{ab}] \times M \quad \text{and} \quad \mathbb{Q}H = \mathbb{Q}[H/[G, G]] \times M';$$

where M is a product of rank p matrix algebras over fields. Then the inclusion $M' \subseteq M$ is a sum of inclusions of the form

$$\prod_{r=1}^p \mathbb{Q}\zeta_{p^r} \subseteq M_p(\mathbb{Q}\zeta_{p^r}); \quad \mathbb{Q}\zeta_{p^{r+1}} \subseteq M_p(\mathbb{Q}\zeta_{p^r}) \quad (r = 1, 2).$$

In particular, $K_2(\hat{\mathbb{Q}}_p \otimes_{\mathbb{Q}} M')_{(p)}$ surjects onto $K_2(\hat{\mathbb{Q}}_p \otimes_{\mathbb{Q}} M)_{(p)}$. Since $Cl_1(\mathbb{Z}H) = 0$ [10, Theorems 4.4.1 and 5.1.1], this shows that

$$K_2(\hat{\mathbb{Q}}_p \otimes_{\mathbb{Q}} M)_{(p)} \subseteq \text{Im} [\varphi_G : K_2(\hat{\mathbb{Z}}_p G) \rightarrow K_2(\hat{\mathbb{Q}}_p G)_{(p)}].$$

In other words, if $\mathbb{Q}[G^{ab}] \cong \prod_{i=1}^k F_i$, then

$$SK_1(\mathbb{Z}G) = Cl_1(\mathbb{Z}G) \cong \text{Coker} \left[\text{proj} \circ \psi_G : H_1(G; \mathbb{Z}G) \rightarrow \prod_{i=1}^k (\mu_{F_i})_p \right].$$

If $G^{ab} \cong C_p \times C_p$, with basis $\{a, b\}$, then $\text{Im}(\text{proj} \circ \psi_G)$ is generated by the images of $a \otimes 1$ and $b \otimes 1$, and so $SK_1(\mathbb{Z}G)$ has rank $(p+1)-2=p-1$. If $G^{ab} \cong C_p^3$, then there are generators a, b, c such that $c \in Z(G)$, and the computation follows from the table in the proof of Proposition 5.1. The proof when $G^{ab} \cong C_{p^2} \times C_p$ is similar. \square

It is interesting to note that for each of these classes of p -groups, the rank of $Cl_1(\mathbb{Z}G)$ is a polynomial in p . This has already been remarked in the case of abelian p -groups (see [1, Conjecture 5.8]); but is harder to formulate as a precise conjecture in the non-abelian case.

Section 5

As another application of Theorem 1.4, we now study the relationship between the complex Artin cokernel

$$A_{\mathbb{C}}(G) = \text{Coker} \left[\sum \{R_{\mathbb{C}}(H) : H \subseteq G \text{ cyclic}\} \xrightarrow{\text{Ind}} R_{\mathbb{C}}(G) \right]$$

of a finite group G , and $Cl_1(RG)$ for large R .

First, epimorphisms

$$I_{RG} : A_{\mathbb{C}}(G) \twoheadrightarrow Cl_1(RG)$$

are constructed, for G any finite group and R the ring of integers in any number field $K \subseteq \mathbb{C}$ (the identification of K as a subfield of \mathbb{C} is needed when defining I_{RG}). The I_{RG} are shown to be natural with respect to homomorphisms and transfer maps, and then shown to be isomorphisms for sufficiently large R .

The following lemma on norm residue symbols will be needed.

LEMMA 5.1. Fix a prime p , fix extensions $E \supseteq F \supseteq \mathbb{Q}_p$, and let $\hat{\mu} \subseteq E^*$ and $\mu \subseteq F^* \cap \hat{\mu}$ be groups of roots of unity. Then the diagram

$$\begin{array}{ccc} K_2(E) & \xrightarrow{(\cdot, \cdot)_{\hat{\mu}}} & \hat{\mu} \\ \downarrow \text{trf}_F^E & & \downarrow [\hat{\mu} : \mu] \\ K_2(F) & \xrightarrow{(\cdot, \cdot)_{\mu}} & \mu \end{array} \quad (1)$$

commutes; where $(\cdot, \cdot)_{\hat{\mu}}$ and $(\cdot, \cdot)_{\mu}$ are the norm residue symbol homomorphisms.

Proof. Set $n = |\hat{\mu}|$ and $m = |\mu|$. Fix $u \in F^*$ and $v \in E^*$, and let $E(\alpha)/E$ be an extension such that $\alpha^n = u$. The diagram

$$\begin{array}{ccc} E^* & \xrightarrow{\hat{s}} & \text{Gal}(E(\alpha)/E) \\ \downarrow N_{E/F} & & \downarrow \text{res} \\ F^* & \xrightarrow{s} & \text{Gal}(F(\alpha^{n/m})/F) \end{array}$$

commutes by [23, Section XI.3]; where \hat{s} and s are the reciprocity maps and res is induced by restriction. By [23, Proposition XIV.6],

$$\begin{aligned} (u, N_{E/F}(v))_{\mu} &= s(N_{E/F}(v))(\alpha^{n/m})/\alpha^{n/m} \\ &= [\hat{s}(v)(\alpha)/\alpha]^{n/m} = ((u, v)_{\hat{\mu}})^{n/m}. \end{aligned} \quad (2)$$

Since $\text{trf}_F^E(\{u, v\}) = \{u, N_{E/F}(v)\}$ for $u \in F^*$ and $v \in E^*$, this shows that (1) commutes on the subgroup $\{F^*, E^*\} \subseteq K_2(E)$. Furthermore,

$$\text{trf}_F^E(\{F^*, E^*\}) = \{F^*, N_{E/F}(E^*)\} = K_2(F):$$

the last equality is shown in [14, Lemma] when $\text{Gal}(E/F)$ is cyclic, and follows from [6, Chapter VI, §2.2] ($N_{E/F}$ is onto) when $\text{Gal}(E/F)$ is non-abelian simple. Since $K_2(E) \cong \mu_E$ and $K_2(F) \cong \mu_F$ are cyclic [12, Theorem A.14], it follows that $\{F^*, E^*\} \supseteq K_2(E)_{(p)}$ for any prime $p \mid |K_2(F)|$, and hence any $p \mid |\mu|$. So (1) commutes. \square

Now fix a finite group G , and let $K \subseteq \mathbb{C}$ be any splitting field for G : i.e., KG is a product of matrix algebras over K . As in [20, Section 2], we define for each prime p :

$$\begin{aligned} C_p(KG) &= \text{Coker} \left[K_2 \left(\mathfrak{M} \left[\frac{1}{p} \right] \right) \rightarrow K_2(\hat{K}_p G) \right] \quad (\hat{K}_p = \mathbb{Q}_p \otimes_{\mathbb{Q}} K) \\ &\cong \text{Coker} [K_2(\mathfrak{M}) \rightarrow K_2(\hat{\mathfrak{M}}_p)] \quad (\hat{\mathfrak{M}}_p = \hat{\mathbb{Z}}_p \otimes_{\mathbb{Z}} \mathfrak{M}) \end{aligned}$$

where $\mathfrak{M} \subseteq KG$ is any maximal order. Then $C_p(KG)$ is a p -group for all p (since $K_2(\mathfrak{M}_p)$ is a p -group). Finally, set

$$C(KG) = \sum_p C_p(KG).$$

Write $KG = \prod_{i=1}^k B_i$, where $B_i \cong \text{End}_K(V_i)$ for each i , and V_1, \dots, V_k are the irreducible KG -modules. By results going back to Bass, Milnor, and Serre [5], $C(KG) = 0$ if K has a real embedding. If K is purely imaginary, then there is an isomorphism

$$\lambda_{KG}: C(KG) \xrightarrow{\cong} \prod_{i=1}^k (\mu_K)$$

such that for any prime $\mathfrak{p} \subseteq R$, and any units $u \in K^*$ and $v \in (\hat{K}_{\mathfrak{p}}[G])^*$,

$$\lambda_{KG}(\{u, v\}_{\mathfrak{p}}) = [(u, \det_K(v, V_i))]_{i=1}^k$$

Here, $\{u, v\}$ denotes the image of

$$\{u, v\} \in K_2(\hat{K}_{\mathfrak{p}}[G]) \rightarrow C(KG);$$

and

$$(\cdot, \cdot)_{\mathfrak{p}}: (\hat{K}_{\mathfrak{p}})^* \times (\hat{K}_{\mathfrak{p}})^* \rightarrow \mu_K$$

denotes the norm residue symbol with values in μ_K . See [20, Theorem 2.2] for more details.

Thus, when $K \subseteq \mathbb{C}$ is a splitting field for G and has no real embedding and $KG \cong \prod_{i=1}^k B_i$ as above, an isomorphism \tilde{I}_{KG} from $R_{\mathbb{C}}(G)$ to $C(KG)$ can be defined as the composite

$$\tilde{I}_{KG}: R_{\mathbb{C}}(G) \cong \prod_{i=1}^k \mathbb{Z} \xrightarrow{\prod [1 \mapsto \exp(2\pi i/m)]} \prod_{i=1}^k \mu_K \xrightarrow{\lambda_{KG}^{-1}} C(KG) \quad (m = |\mu_K|).$$

In other words, for each $1 \leq i \leq k$, we set

$$I_{KG}([V_i]) = \lambda_{KG}^{-1}([\exp(2\pi i/m)]_i);$$

where $[V_i] \in R_{\mathbb{C}}(G)$ denotes the class of $\mathbb{C} \otimes_K V_i$.

If K is a splitting field for G but has a real embedding, we set $\tilde{I}_{KG} = 0$ ($C(KG) = 0$). If $K \subseteq \mathbb{C}$ is a number field which does not split G , set $n = \exp(G)$ and $L = K(\zeta_n)$, and define

$$\tilde{I}_{KG} = \text{trf}_{KG}^{LG} \circ \tilde{I}_{LG} : R_{\mathbb{C}}(G) \rightarrow C(LG) \rightarrow C(KG).$$

(Note that L is a splitting field for G by [5, Theorem 4.1.1].) This definition of the I_{KG} seems rather artificial; but the following proposition shows that these maps do have all desired naturality properties.

PROPOSITION 5.2. *For any number field $K \subseteq \mathbb{C}$ and any finite group G , \tilde{I}_{KG} is surjective. The \tilde{I}_{KG} are natural in that for any homomorphism $\alpha : \tilde{G} \rightarrow G$ of finite groups, for any $H \subseteq G$, and for any pair $K \subseteq L \subseteq \mathbb{C}$ of number fields, the following diagrams all commute:*

$$\begin{array}{ccc} R_{\mathbb{C}}(G) & & R_{\mathbb{C}}(\tilde{G}) \xrightarrow{R_{\mathbb{C}}(\alpha)} R_{\mathbb{C}}(G) \xrightarrow{\text{Res}_H^G} R_{\mathbb{C}}(H) \\ \swarrow (1) \searrow & & \downarrow i_{KG} \quad (2) \quad \downarrow \tilde{i}_{KG} \quad (3) \quad \downarrow \tilde{i}_{KH} \\ C(LG) \xrightarrow{\text{trf}_{KG}^{LG}} C(KG) & & C(K\tilde{G}) \xrightarrow{C(K\alpha)} C(KG) \xrightarrow{\text{trf}_{KH}^{KG}} C(KH) \end{array}$$

Proof. The proposition will be proven in four steps. For finite G and arbitrary $K \subseteq \mathbb{C}$, we regard $K_0(KG) = R_K(G)$ as a subring of $R_{\mathbb{C}}(G)$ in the usual fashion (identifying $[V] \in R_K(G)$ with $[C \otimes_K V] \in R_{\mathbb{C}}(G)$).

Step 1. By construction, \tilde{I}_{KG} is surjective if K splits G . To see that \tilde{I}_{KG} is surjective in general, we must show for any G , and any number fields $K \subseteq L$, that the transfer map

$$\text{trf}_{KG}^{LG} : K_2(\hat{L}_p G) \rightarrow K_2(\hat{K}_p G)$$

is onto for each prime p ($\hat{L}_p = \hat{\mathbb{Q}}_p \otimes_{\mathbb{Q}} L$, etc.).

Write $\hat{K}_p G \cong \prod_{i=1}^k M_{n_i}(D_i)$, where the D_i are division algebras. For each i , set $F_i = Z(D_i)$, the center, and let $E_i \subseteq D_i$ be a maximal subfield. By [3, Corollary 4.15], $K_2(D_i)$ is generated by symbols $\{F_i^*, D_i^*\}$; and hence $K_2(E_i) \rightarrow K_2(D_i)$ is onto by [14, Proposition].

Consider the following square, for each $1 \leq i \leq k$:

$$\begin{array}{ccc} K_1(L \otimes_K E_i) & \xrightarrow{\text{incl}} & K_2(L \otimes_K D_i) \\ \downarrow t'_i & & \downarrow t_i \\ K_2(E_i) & \longrightarrow \longrightarrow & K_2(D_i). \end{array}$$

Here t_i and t'_i are the transfer maps. The square commutes since the two sides are induced by tensoring with the bimodules

$$D_i \otimes_{E_i} (L \otimes_K E_i) \cong L \otimes_K D_i.$$

The map t'_i is the product of the transfer homomorphisms for the field summands of $L \otimes_K E_i$, each of which is onto by [12, Corollary A.15]. So t_i is also onto. But trf_{KG}^{LG} is isomorphic to the sum of the t_i , and is hence surjective.

Step 2. Fix K and G such that K is a totally imaginary splitting field for G . In particular, $K_0(KG) = R_{\mathbb{C}}(G)$. For any finite dimensional (left) KG -module V , define

$$f_V : C(K) \rightarrow C(KG)$$

to be the homomorphism induced by the functor

$$V \otimes_K : K\text{-mod} \rightarrow KG\text{-mod}.$$

If V is irreducible, then f_V is just the Morita equivalence identifying $C(B)$ with $C(K)$, where $B \subseteq KG$ is the simple summand with irreducible representation V . So by definition,

$$\tilde{I}_{KG}([V]) = f_V(\lambda_K^{-1}(\exp(2\pi i/m))); \quad (m = |\mu_K|) \quad (4)$$

where $\lambda_K : C(K) \xrightarrow{\cong} \mu_K$ is induced by the norm residue symbol. Both sides of (4) are additive ($f_{V \oplus W} = f_V + f_W$), so (4) holds for arbitrary V .

Step 3. We can now show the commutativity of triangle (1) above: that $\tilde{I}_{KG} = \text{trf}_{KG}^{LG} \circ \tilde{I}_{LG}$ for any G and any number fields $K \subseteq L \subseteq \mathbb{C}$. It suffices to do this when K and L both are totally imaginary splitting fields for G . In particular, $K_0(KG) = R_{\mathbb{C}}(G)$.

By (4), for any finite dimensional KG -module V ,

$$\tilde{I}_{KG}([V]) = f_V(\lambda_K^{-1}(\exp(2\pi i/m))), \quad (m = |\mu_K|)$$

$$\text{trf}_{KG}^{LG}(\tilde{I}_{LG}([V])) = \text{trf}_{KG}^{LG} \circ f_{L \otimes_K V}(\lambda_L^{-1}(\exp(2\pi i/n))); \quad (n = |\mu_L|)$$

and it remains to check the commutativity of the following diagram:

$$\begin{array}{ccccc}
 \mu_L & \xrightarrow{\lambda_L^{-1}} & C(L) & \xrightarrow{f_L \otimes v} & C(LG) \\
 \downarrow n/m & (5) & \downarrow \text{trf}_K^L & (6) & \downarrow \text{trf}_{KG}^{LG} \\
 \mu_K & \xrightarrow{\lambda_K^{-1}} & C(K) & \xrightarrow{f_v} & C(KG).
 \end{array}$$

But (5) commutes by Lemma 5.1, while (6) commutes since the two composites are induced by tensoring with the bimodules

$$KG^{LG} \otimes_{LG} (L \otimes_K V)_L = {}_{KG}V \otimes_K L_L.$$

Step 4. Now fix a homomorphism $\alpha: \tilde{G} \rightarrow G$ of finite groups, and a subgroup $H \subseteq G$. We must show that (2) and (3) commute for any number field $K \leq \mathbb{C}$. If $L \supseteq K$ is any pair of number fields, then the squares

$$\begin{array}{ccccc}
 C(L\tilde{G}) & \xrightarrow{C(L\alpha)} & C(LG) & \xrightarrow{\text{trf}_{LG}^{LG}} & C(LH) \\
 \downarrow \text{trf}_{KG}^{L\tilde{G}} & & \downarrow \text{trf}_{KG}^{LG} & & \downarrow \text{trf}_{KH}^{LH} \\
 C(K\tilde{G}) & \xrightarrow{C(K\alpha)} & C(KG) & \xrightarrow{\text{trf}_{KH}^{KG}} & C(KH)
 \end{array}$$

commute (just compare bimodules). So by (1) (Step 3), it suffices to prove the commutativity of (2) and (3) when K is a splitting field for \tilde{G} , G and H (and totally imaginary).

Fix such a K ; in particular, $K_0(K\tilde{G}) = R_{\mathbb{C}}(\tilde{G})$ and $K_0(KG) = R_{\mathbb{C}}(G)$. Fix finite dimensional modules V over $K\tilde{G}$ and W over KG . Set

$$x = \lambda_K^{-1}(\exp(2\pi i/|\mu_k|)) \in C(K).$$

Then, by (4),

$$\begin{aligned}
 \tilde{I}_{KG}(R_{\mathbb{C}}(\alpha)([V])) &= f_{KG \otimes_{K\tilde{G}} V}(x), \\
 C(K\alpha) \circ \tilde{I}_{K\tilde{G}}([V]) &= C(K\alpha) \circ f_V(x); \\
 \tilde{I}_{KH}(\text{Res}_H^G([W])) &= f_{W|H}(x),
 \end{aligned}$$

and

$$\text{trf}_{KH}^{KG} \circ \tilde{I}_{KG}([W]) = \text{trf}_{KH}^{KG} \circ f_W(x).$$

So we will be done upon showing that the following triangles commute:

$$\begin{array}{ccc}
 & C(K\tilde{G}) & \\
 f_V \nearrow & \downarrow C(K\alpha) & \\
 C(K) & & \\
 \downarrow f_{KG} \otimes_{K\tilde{G}} V & & \\
 & C(KG) & \\
 \end{array}
 \quad
 \begin{array}{ccc}
 & C(KG) & \\
 f_W \nearrow & \downarrow \text{trf}_{KH}^{KG} & \\
 C(K) & & \\
 \downarrow f_{WH} & & \\
 & C(KH) & \\
 \end{array}$$

But they are induced by the following pairs of isomorphic bimodules:

$${}_{KG}(KG \otimes_{K\tilde{G}} V)_K \cong {}_{KG}KG \otimes_{K\tilde{G}} V_K \quad \text{and} \quad {}_{KH}W_K \cong {}_{KH}KG \otimes_{KG} W_K;$$

and we are done. \square

Again fix a finite group G and a number field $K \subseteq \mathbb{C}$, and let $R \subseteq K$ be the ring of integers. Then $Cl_1(RG)$ is described by a localization sequence

$$\sum_p K_2(\hat{R}_p G) \xrightarrow{\partial_{RG}} C(KG) \xrightarrow{\partial_{RG}} Cl_1(RG) \rightarrow 0$$

(see [20, Theorem 2.1] for details). We now consider the composite

$$R_C(G) \xrightarrow{\tilde{I}_{KG}} C(KG) \xrightarrow{\partial_{RG}} Cl_1(RG).$$

Both maps are natural with respect to induction from subgroups of G . Hence, since $Cl_1(RH) = SK_1(RH) = 0$ for any cyclic $H \subseteq G$ by [1, Theorem 3.3], $\partial_{RG} \circ \tilde{I}_{KG}$ vanishes on any element of $R_C(G)$ induced up from a cyclic subgroup. Thus, $\partial_{RG} \circ \tilde{I}_{KG}$ factors through a homomorphism

$$I_{RG} : A_C(G) \rightarrow Cl_1(RG),$$

where $A_C(G)$ is the Artin cokernel.

THEOREM 5.3. *For any finite group G , and any number field $K \subseteq \mathbb{C}$ with ring of integers R ,*

$$I_{RG} : A_C(G) \rightarrow Cl_1(RG)$$

is surjective. The I_{RG} are natural in that for any homomorphism $\alpha : \tilde{G} \rightarrow G$ of finite groups, any $H \subseteq G$, and any pair $R \subseteq S$ of rings of integers in number fields, the

following diagrams all commute:

$$\begin{array}{ccccc}
 A_{\mathbb{C}}(G) & & A_{\mathbb{C}}(\tilde{G}) & \xrightarrow{A_{\mathbb{C}}(\alpha)} & A_{\mathbb{C}}(G) \xrightarrow{\text{Res}_H^G} A_{\mathbb{C}}(H) \\
 \swarrow I_{KG} \quad \searrow I_{RG} & & \downarrow I_{R\tilde{G}} & & \downarrow I_{RG} \quad \downarrow I_{RH} \\
 Cl_1(SG) \xrightarrow{\text{trf}_{RG}^{SG}} Cl_1(RG) & & Cl_1(R\tilde{G}) \xrightarrow{Cl_1(A\alpha)} Cl_1(RG) \xrightarrow{\text{trf}_{RH}^{RG}} Cl_1(RH).
 \end{array}$$

Proof. For any R and G , I_{RG} is surjective since \tilde{I}_{KG} and ∂_{RG} both are surjective. The naturality properties follow from the corresponding properties for the \tilde{I}_{KG} (Proposition 5.2), and the naturality of the boundary maps ∂_{RG} in the localization sequences. \square

Now that the I_{RG} have been constructed, we can finally apply Theorem 1.4 to show that they are isomorphisms for sufficiently large R . For any finite G , $a_{\mathbb{C}}(G)$ will denote the complex Artin exponent: the order of $1 \in R_{\mathbb{C}}(G)$ in $A_{\mathbb{C}}(G)$. By Frobenius reciprocity,

$$a_{\mathbb{C}}(G) = \exp(A_{\mathbb{C}}(G));$$

i.e., $a_{\mathbb{C}}(G) \cdot x$ is induced from cyclic subgroups for any $x \in R_{\mathbb{C}}(G)$. By the Artin induction theorem [7, Theorem 39.1], $a_{\mathbb{C}}(G) \mid |G|$.

THEOREM 5.4. *Let G be any finite group, and set $n = a_{\mathbb{C}}(G) \cdot \exp(G)$. Let K be any number field such that $\zeta_n \in K$, and let $R \subseteq K$ be the ring of integers. Then I_{RG} is an isomorphism: $Cl_1(RG) \cong A_{\mathbb{C}}(G)$.*

Proof. This will be shown first for p -groups, then for p -elementary groups, and finally for arbitrary finite groups.

Step 1. Let G be a p -group, and set $p^k = a_{\mathbb{C}}(G)$, $p^m = \exp(G)$, and $q = p^{k+m}$. By Theorem 5.3(1), it will suffice to show that I_{RG} is an isomorphism when $K = \mathbb{Q}\zeta_q$ and $R = \mathbb{Z}\zeta_q$.

Let C_q be a (multiplicative) cyclic group of order q with generator z . Consider the pullback square

$$\begin{array}{ccc}
 \hat{\mathbb{Z}}_p[C_q \times G] & \xrightarrow{\alpha} & \hat{\mathbb{Z}}_p\zeta_q[G] \\
 \downarrow & & \downarrow \\
 \mathbb{Z}_p[(C_q/z^{p^{n+m-1}}) \times G] & \xrightarrow{\beta} & \mathbb{Z}/p[(C_q/z^{p^{n+m-1}}) \times G];
 \end{array} \tag{1}$$

where α is induced by: $\alpha(z) = \zeta_q$. Then $K_2(\beta)$ is onto by [17, Lemma 1.7] if $p > 2$; or if $p = 2$ since the only torsion in $K_1(\hat{\mathbb{Z}}_2[(C_q/z^{p^{n+m-1}}) \times G], 2)$ is (-1) (see [15, Proposition 2]). So by the Mayer–Vietoris sequence for (1), $K_2(\alpha)$ is onto.

Now consider the following commutative diagram:

$$(2) \quad \begin{array}{ccccccc} K_2(\hat{\mathbb{Z}}_p[C_q \times G]) & \longrightarrow & C(\mathbb{Q}[C_q \times G])_{(p)} & R_{\mathbb{C}}(G) \otimes \mathbb{Z}/q \\ \downarrow K_2(\alpha) & & \alpha_* \downarrow & \swarrow \bar{I}_{KG} & \downarrow \partial \circ \bar{I}_{KG} \\ K_2(\hat{\mathbb{Z}}_p \zeta_q[G]) & \xrightarrow{\varphi_{RG}} & C_p(\mathbb{Q}\zeta_q[G]) & \xrightarrow{\partial} & Cl_1(RG)_{(p)} & \longrightarrow 0 \end{array} \quad (2)$$

where the bottom row is exact [20, Theorem 2.1]. By Theorem 1.4,

$$\begin{aligned} K_2(\hat{\mathbb{Z}}_p[C_q \times G]) &= K_2(\hat{\mathbb{Z}}_p G) \oplus K_2(\hat{\mathbb{Z}}_p[C_q \times G], (1-z)) \\ &= K_2(\hat{\mathbb{Z}}_p G) \oplus \langle \{h, 1 - (1-z)^i g\}, \{z, 1 - (1-z)^i g\}: \\ &\quad h \in G, g \in C_q \times G, hg = gh, i \geq 1 \rangle. \end{aligned}$$

It follows that

$$K_2(\hat{\mathbb{Z}}_p \zeta_q[G]) = K_2(\hat{\mathbb{Z}}_p G) + X + Y;$$

where with $\zeta = \zeta_q$:

$$X = \langle \{h, 1 - (1 - \zeta)^i \zeta^j g\}: h, g \in G, hg = gh, i \geq 1, j \in \mathbb{Z} \rangle$$

and

$$Y = \langle \{\zeta, 1 - (1 - \zeta)^i \zeta^j g\}: g \in G, i \geq 1, j \in \mathbb{Z} \rangle.$$

Recall that $p^m = \exp(G)$. Then

$$\exp(X) \mid p^m \quad \text{and} \quad \exp(\varphi_{RG}(K_2(\hat{\mathbb{Z}}_p G))) \mid \exp(C(\mathbb{Q}G)_{(p)}) \mid p^m.$$

Furthermore, by definition,

$$\varphi_{RG}(Y) \subseteq \text{Im} \left[\sum \{C(\mathbb{Q}\zeta_q[H]): H \subseteq G \text{ cyclic}\} \rightarrow C(\mathbb{Q}\zeta_q[G]) \right].$$

So by diagram (2) (recalling that $q = p^{k+m}$):

$$\begin{aligned} \text{Ker}(\partial \circ \bar{I}_{KG}) &\subseteq p^k R_{\mathbb{C}}(G) + \text{Im} \left[\sum \{R_{\mathbb{C}}(H): H \subseteq G \text{ cyclic}\} \rightarrow R_{\mathbb{C}}(G) \right] \\ &= p^k R_{\mathbb{C}}(G) + \text{Ker}[R_{\mathbb{C}}(G) \rightarrow A_{\mathbb{C}}(G)]. \end{aligned}$$

Since $p^k = a_{\mathbb{C}}(G) = \exp(A_{\mathbb{C}}(G))$,

$$\text{Ker } [I_{RG}: A_{\mathbb{C}}(G) \rightarrow Cl_1(RG)] \subseteq p^k A_{\mathbb{C}}(G) = 0;$$

and so I_{RG} is an isomorphism.

Step 2. Now assume that G is p -elementary: $G \cong C_m \times H$ where $p \nmid m$ and H is a p -group. Set $n = a_{\mathbb{C}}(G) \cdot \exp(G)$, fix a number field $K \subseteq \mathbb{C}$ containing ζ_n , and let R be the ring of integers of K . Then

$$\begin{aligned} A_{\mathbb{C}}(G) &\cong \text{Coker} \left[\sum \{R_{\mathbb{C}}(C_m) \otimes R_{\mathbb{C}}(H_0) : H_0 \subseteq H \text{ cyclic}\} \rightarrow R_{\mathbb{C}}(C_m) \otimes R_{\mathbb{C}}(H) \right] \\ &\cong R_{\mathbb{C}}(C_m) \otimes A_{\mathbb{C}}(H) \cong \prod^m A_{\mathbb{C}}(H). \end{aligned}$$

On the other hand, the identification $K[G] \cong \prod^m K[H]$ (each factor corresponding to a character of C_m) induces an inclusion $RG \subseteq \prod^m R[H]$ of index prime to p ; and hence an isomorphism

$$Cl_1(RG)_{(p)} \cong \prod^m Cl_1(RH)_{(p)} \cong \prod^m A_{\mathbb{C}}(H) \cong A_{\mathbb{C}}(G).$$

(see [17, Proposition 1.2]). Since I_{RG} is onto, it must be an isomorphism.

Step 3. Now let G be an arbitrary finite group, set $n = a_{\mathbb{C}}(G) \cdot \exp(G)$, and let R be any ring of integers containing ζ . Let \mathcal{E} be the set of elementary subgroups of G . For any $H \in \mathcal{E}$, $\exp(H) \mid \exp(G)$ and $a_{\mathbb{C}}(H) \mid a_{\mathbb{C}}(G)$, so I_{RH} is an isomorphism by Step 2. Consider the following square, which commutes by Theorem 5.3:

$$\begin{array}{ccc} A_{\mathbb{C}}(G) & \xrightarrow{I_{RG}} & Cl_1(RG) \\ \downarrow \sum \text{Res}_H^G & & \downarrow \sum \text{trf}_H^G \\ \sum_{H \in \mathcal{E}} A_{\mathbb{C}}(H) & \xrightarrow{\cong \sum I_{RH}} & \sum_{H \in \mathcal{E}} Cl_1(RH) \end{array}$$

In the language of [10], $A_{\mathbb{C}}(-)$ is a module over the Frobenius functor $R_{\mathbb{C}}(-)$, and hence is detected by restriction to elementary subgroups. So $\sum \text{Res}_H^G$ is injective in the above square, and I_{RG} is an isomorphism. \square

By [4, Theorem XI.4.7], for any finite G ,

$$a_{\mathbb{C}}(G) = \prod_{p \mid |G|} a_{\mathbb{C}}(G_p),$$

where G_p is a p -Sylow subgroup. Thus, the description of

$$a_C(G) = \exp(A_C(G)) = \max_R (\exp(Cl_1(RG)))$$

reduces immediately to the p -group case.

If G is a non-cyclic p -group, then there is a surjection $G \twoheadrightarrow C_p \times C_p$ and an induced surjection of $A_C(G)$ onto $A_C(C_p \times C_p)$. This last group is easily checked to be non-zero (see [1, Lemma 5.5] for details). Thus, for any finite G , $A_C(G)$ is p -torsion free if and only if G_p is cyclic, $A_C(G) = 0$ if and only if G is metacyclic, and these in turn imply similar statements about the $Cl_1(RG)$ (and $SK_1(RG)$). In fact, for fixed p and R such that $\zeta_p \in R$ (or $\zeta_4 \in R$ if $p = 2$), and any G , $Cl_1(RG)_{(p)} = 0$ if and only if G_p is cyclic (see [1, Theorem 3.5]).

A general description of $a_C(G)$ has been given by Gluck [27]. The formula is much more complicated than that for the rational Artin exponent $a_Q(G)$ given by Lam [11]. If G is non-cyclic, and abelian or of exponent p , then $a_C(G) = a_Q(G) = (1/p)|G|$. On the other hand, if G is a semidihedral 2-group, then $a_C(G) = 2$ ($a_Q(G) = 4$); and if p is odd and G a non-abelian group of order p^3 and exponent p^2 , then $a_C(G) = p$ ($a_Q(G) = p^2$).

To end, we note that Theorem 5.3 allows a new interpretation of the following result in [13] (Theorem 1).

COROLLARY 5.5. *Let G be a finite group, and let R be the ring of integers in some number field. Then $Cl_1(RG)$ is generated by induction from elementary subgroups of G .*

Proof. By the Brauer induction theorem, $R_C(G)$, and hence $A_C(G)$ are generated in induction from elementary subgroups of G . The result follows since $I_{RG}: A_C(G) \rightarrow Cl_1(RG)$ is natural and surjective. \square

REFERENCES

- [1] R. ALPERIN, K. DENNIS, R. OLIVER and M. STEIN, *SK₁ of finite abelian groups: II*, Invent. Math. 87 (1987), 253–302.
- [2] E. ARTIN and H. HASSE, *Die beiden Ergänzungssätze zum Reziprozitätsgesetz der lⁿ-ten Potenzreste im Körper der lⁿ-ten Einheitswurzeln*, Abh. Math. Sem. Hamburg 6 (1928), 146–162.
- [3] A. BAK and U. REHMANN, *The congruence subgroup and metaplectic problems for SL_{n≥2} of division algebras*, J. Algebra 78 (1982), 475–547.
- [4] H. BASS, *Algebraic K-theory*, Benjamin (1968).
- [5] H. BASS, J. MILNOR and J.-P. SERRE, *Solution of the congruence subgroup problem for SL_n (n ≥ 3) and Sp_{2n} (n ≥ 2)*, I.H.E.S., Publications Math. 33 (1967), 59–137.

- [6] J. W. S. CASSELS and A. FRÖHLICH, *Algebraic number theory*, Academic Press (1967).
- [7] C. CURTIS and I. REINER, *Representation theory of finite groups and associative algebras* (Interscience, London, 1962).
- [8] P. HILTON and U. STAMMBACH, *A course in homological algebra*, Springer-Verlag (1971).
- [9] B. HUPPERT, *Endliche Gruppen*, Springer-Verlag (1967).
- [10] T.-Y. LAM, *Induction theorems for Grothendieck groups and Whitehead groups of finite groups*, Ann. Sci. Ecole Norm. Sup. 1 (1968), 91–148.
- [11] T. Y. LAM, *Artin exponent of finite groups*, J. Alg. 9 (1968), 94–119.
- [12] J. MILNOR, *Introduction to Algebraic K-theory*, Princeton Univ. Press (1971).
- [13] R. OLIVER, *SK₁ for finite group rings: I*, Invent. Math. 57 (1980), 183–204.
- [14] R. OLIVER, Correction to: *SK₁ for finite group rings: I*, Invent. Math. 64 (1981), 167–169.
- [15] R. OLIVER, *SK₁ for finite group rings: II*, Math. Scand. 47 (1980), 195–231.
- [16] R. OLIVER, *SK₁ for finite group rings: III*, Lecture Notes in Math. vol. 854, Springer-Verlag (1981), 299–337.
- [17] R. OLIVER, *SK₁ for finite group rings: IV*, Proc. London Math. Soc. 46 (1983), 1–37.
- [18] R. OLIVER, *D($\mathbb{Z}\pi$)⁺ and the Artin cokernel*, Comment. Math. Helv. 58 (1983), 291–311.
- [19] R. OLIVER, *Lower bounds for K₂^{top}($\hat{\mathbb{Z}}_p\pi$) and K₂($\mathbb{Z}\pi$)*, J. Algebra 94 (1985), 425–487.
- [20] R. OLIVER, *The Whitehead transfer homomorphism for oriented S¹-bundles*, Math. Scand. 57 (1985), 51–104.
- [21] R. OLIVER, *K₂ of p-adic group rings of abelian p-groups* Math. Z. (to appear).
- [22] P. ROQUETTE, *Realisierung von Darstellungen endlicher nilpotenter Gruppen*, Arkiv der Math. 9 (1958), 241–250.
- [23] J.-P. SERRE, *Corps Locaux*, Hermann (1968).
- [24] R. SWAN, *Excision in algebraic K-theory*, J. Pure Appl. Alg. 1 (1971), 221–252.
- [25] J. SYLVESTER, *Introduction to Algebraic K-theory*, Chapman and Hall (1981).
- [26] C. T. C. WALL, *Norms of units in groups rings*, Proc. London Math. Soc. 29 (1974), 593–632.
- [27] D. GLUCK, *Artin exponent for arbitrary characters of cyclic subgroups*, J. Algebra 61 (1979), 58–70.

Department of Mathematics
 Aarhus University
 Ny Munkegade
 8000 Aarhus, Denmark

Received May 5, 1985

Buchanzeigen

D. AMIR, Characterizations of Inner Product Spaces. (Operator Theory Vol. 20) Birkhäuser Verlag Basel–Boston–Stuttgart, 1986, 200 pp., SFr. 60.

I. Introduction – II. 2-dimensional Characterization – III. 3-Dimensional Characterizations – IV. Tables and references.

I. GOHBERG and M. A. KAASHOEK, Constructive Methods of Wiener–Hopf Factorization. (Operator-Theory Vol. 21), Birkhäuser-Verlag Basel–Boston–Stuttgart 1986, 409 pp., SFr. 98. Part I: Canonical and minimal factorization (contributions by J. A. Ball/A.C.M. Ran, H. Bart/I. Gohberg/M.A. Kaashoek, I. Gohberg/M.A. Kaashoek/L. Lerer/L. Rodman, L. Roodeman, I. Gohberg/M.A. Kaashoek) – Part II: Non-Canonical Wiener-Hopf Factorization (contributions by H. Bart/I. Gohberg/M.A. Kaashoek, H. Bart/I. Gohberg/M.A. Kaashoek, M.A. Kaashoek/A.C.M. Ran).

V.A. MARCHENKO, Sturm–Liouville Operators and Applications. (Operator-Theory Vol. 22), translated from the Russian by A. Iacob, Birkhäuser–Verlag Basel–Boston–Stuttgart, 1986, 367 pp., 110.

1. The Sturm–Liouville equation and transformation operators – 2. The Sturm–Liouville boundary value problem on the half line – 3. The boundary value problem of scattering theory – 4. Nonlinear equations.

HYO CHUL MYUNG, Malcev-Admissible Algebras. (Progress in Math. Vol. 64), Birkhäuser Verlag Boston–Basel–Stuttgart, 1986, 353 pp., SFr. 118.

1. Flexible Malcev-admissible algebras – 2. Power-associative Malcev-admissible algebras – 3. Invariant operators in simple Lie algebras and flexible Malcev-admissible algebras with A^- simple – 4. Malcev-admissible algebras with the solvable radical of A^- nonzero – 5. Malcev-admissible algebras of low dimension.

G. GRUBB, Functional Calculus of Pseudo-Differential Boundary Problems. (Progress in Math. Vol. 65), Birkhäuser Verlag Boston–Basel–Stuttgart, 1986, 511 pp., SFr. 98.

1. Standard pseudo-differential boundary problems and their realizations – 2. The calculus of parameter-dependent operators – 3. Parametrix and resolvent constructions – 4. Some applications – Appendix: Various prerequisites.

J. GRAY, Linear Differential Equations and Group Theory from Riemann to Poincaré. Birkhäuser–Verlag, Boston–Basel–Stuttgart, 1986, 460 pp. SFr. 98.

1. Hypergeometric equations and modular equations – 2. Lazarus Fuchs – 3. Algebraic Solutions to a Differential Equation – 4. Modular Equations – 5. Some algebraic Curves – 6. Automorphic Functions – Appendices: Notes and Bibliography – Riemann, Schottky, and Schwarz on conformal representation – Riemann's lectures on differential equations – Fuchs's analysis of the n th order equation – On the History of non-Euclidean geometry – Notes on Chapters 1 – 6 and Appendix 4.

H.P. YAP, Some Topics in Graph Theory. (London Math. Soc. Lecture Note Series 108), Cambridge University Press, Cambridge, London etc., 1986, 230 pp., £12.50, \$24.95.

1. Basic Terminology – 2. Edge-colourings of Graphs – 3. Symmetries in Graphs – 4. Packing of Graphs – 5. Computational Complexity of Graph Properties.