

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 62 (1987)

Artikel: Gaps and bands of one dimensional periodic Schrödinger operators, II.
Autor: Garnett, John / Trubowitz, Eugene
DOI: <https://doi.org/10.5169/seals-47338>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Gaps and bands of one dimensional periodic Schrödinger operators, II

JOHN GARNETT and EUGENE TRUBOWITZ

§1. Introduction

Let $q(x) \in L^2_{\mathbf{R}}[0, 1]$, the Hilbert space of square integrable real valued functions on the unit interval. Extend $q(x)$ to the whole line \mathbf{R} by $q(x+1) = q(x)$. The spectrum of the Schrödinger operator $-d^2/dx^2 + q(x)$, acting on $L^2(\mathbf{R})$, is the set of λ such that

$$-y'' + q(x)y = \lambda y \tag{1.1}$$

has a nontrivial solution bounded on \mathbf{R} . The spectrum is contained in \mathbf{R} and it is the union of a sequence of closed intervals $[\lambda_{2n-2}, \lambda_{2n-1}]$, where $\lambda_n = \lambda_n(q)$, $n \geq 0$, satisfies

$$\lambda_0 < \lambda_1 \leq \lambda_2 < \lambda_3 \leq \lambda_4 < \cdots.$$

These intervals are called *bands* and the intervening, possibly void, open intervals are called *gaps*. The possible arrangements of gaps and bands were investigated in [1]. This paper continues that study and includes some applications and simplifications.

Let $\gamma_n(q) = \lambda_{2n}(q) - \lambda_{2n-1}(q)$ be the n -th gap length. It is well known that $\gamma_n(q) \in (l^2)^+$, the space of nonnegative sequences with $\sum \gamma_n^2 < \infty$. Two of the three main results of [1] are:

(a) *Whenever $\gamma_n \in (l^2)^+$, there exists $q \in L^2_{\mathbf{R}}([0, 1])$ such that $\gamma_n(q) = \gamma_n$, $n = 1, 2, \dots$. Moreover, q can be chosen from the even subspace E of $q \in L^2_{\mathbf{R}}[0, 1]$ such that*

$$q(1-x) = q(x)$$

(b) *the spectrum is determined, up to a translation, by the gap lengths $\gamma_n(q)$.*

Let $\mu_n(q)$, $n \geq 1$, be the Dirichlet spectrum of q , that is, the spectrum of (1.1) for the boundary condition

$$y(0) = y(1) = 0,$$

and let $\nu_n(q)$, $n \geq 0$, be it's Neumann spectrum, i.e. the spectrum of (1.1) with boundary condition

$$y'(0) = y'(1) = 0.$$

Then $q \in E$ if and only if $\{\mu_n(q), \nu_n(q)\} = \{\lambda_{2n-1}(q), \lambda_{2n}(q)\}$, so that for q even,

$$\gamma_n(q) = |\mu_n(q) - \nu_n(q)|.$$

As functions on $L^2_{\mathbf{R}}[0, 1]$, $\mu_n(q)$ and $\nu_n(q)$ are real analytic (while λ_{2n} is not analytic at a q for which $\lambda_{2n}(q) = \lambda_{2n-1}(q)$) and hence the *signed gap length* $\sigma_n(q) = \mu_n(q) - \nu_n(q)$, $n \geq 1$, is real analytic in q . Furthermore, the map $\sigma: L^2_{\mathbf{R}}[0, 1] \rightarrow \ell^2$ defined by $\sigma(q) = (\sigma_n(q))$, $n \geq 1$, is a real analytic mapping from the Hilbert space $L^2_{\mathbf{R}}[0, 1]$ to the Hilbert space ℓ^2 . The third main result of [1] is:

(c) *Let E_0 be the space of even potentials in $L^2_{\mathbf{R}}[0, 1]$ satisfying $\int_0^1 q(x) dx = 0$. Then the map*

$$E_0 \ni q \rightarrow \sigma(q) = (\sigma_1(q), \sigma_2(q), \dots)$$

is a real analytic isomorphism between E_0 and ℓ^2 , that is, σ is one-to-one and onto and both σ and σ^{-1} are real analytic maps of Hilbert space.

Of course, since $\gamma_n(q) = |\sigma_n(q)|$, $q \in E_0$, result (c) included result (a).

The proof of (a), (b) and (c) in [1] applied harmonic measure arguments to the identification, due to Marčenko and Ostrovskii [3], of band configurations with certain slit quarter planes. In Section 2 we give a direct proof, using analysis in Hilbert space, that the Jacobian

$$d_q \sigma: E_0 \rightarrow \ell^2$$

is invertible. From this it follows easily that σ is one-to-one, and that, if σ is onto, than by the Inverse Function Theorem, σ^{-1} is real analytic. Consequently, result (c) can be proved without the intricate Section 6 of [1]. We cannot prove σ is onto ℓ^2 using only the method of Section 2 without a still unknown estimate of $\|q\|_2$ in terms of $\|\sigma(q)\|_{\ell^2}$. However, in Sobolev space such an estimate is

available and thus we show in Section 2 that σ is an isomorphism from $E_0 \cap H^k = \{q \in E_0 : q \text{ has } k \text{ derivatives periodic and in } L^2([0, 1])\}$ onto $\ell_k^2 = \{(\sigma_n) : \sum n^{2k} \sigma_n^2 < \infty\}$.

In Section 3 result (c) is used to prove Marčenko's theorem [2] that the finite band potentials (those q with $\gamma_n(q) = 0$ for large n) are norm dense in L^2 , and that q has primitive period $1/k$ if and only if $\gamma_n(q) = 0$ when k does not divide n .

In Section 4 we give some inequalities that band lengths must satisfy and we show that for real analytic potentials the band lengths determine the spectrum up to a translation. Here the harmonic measure methods of [1] reappear.

§2. Signed gap lengths

We need a general interpolation lemma.

LEMMA 2.1. *Suppose $\phi(\lambda)$ is an entire function satisfying*

$$\sup_{|\lambda|=(n+1/2)^2\pi^2} \left| \phi(\lambda) / \frac{\sin \sqrt{\lambda}}{\sqrt{\lambda}} \right| = o(1)$$

as $n \rightarrow \infty$. Then

$$\phi(\lambda) = \sum_{n \geq 1} \phi(\xi_n) \prod_{\substack{m \geq 1 \\ m \neq n}} \frac{\xi_m - \lambda}{\xi_m - \xi_n}$$

for any sequence ξ_n , $n \geq 1$, of distinct complex numbers satisfying $\xi_n = n^2\pi^2 + o(1)$.

Proof. If ξ_m , $m \geq 1$, is a distinct sequence with $\xi_m = m^2\pi^2 + o(1)$, then

$$\prod \frac{\xi_m - z}{m^2\pi^2} = \frac{\sin \sqrt{z}}{\sqrt{z}} \left(1 + o\left(\frac{\log n}{n}\right) \right)$$

uniformly on the circles $\Gamma_n = \{|z| = (n + 1/2)^2\pi^2\}$. Hence the meromorphic function

$$f(z) = \frac{\phi(z)}{z - \lambda} \prod_{m \geq 1} \frac{m^2\pi^2}{\xi_m - z}$$

satisfies $\sup_{\Gamma_n} |f(z)| = o(n^{-2})$, $n \rightarrow \infty$; and the sum of its residues inside Γ_n has

limit 0 as $n \rightarrow \infty$. But $f(z)$ has simple poles of λ and at ξ_n , $n \geq 1$, and $f(z)$ is regular elsewhere. Summing the residues, we obtain

$$0 = \phi(\lambda) \prod_{m \geq 1} \frac{m^2 \pi^2}{\xi_m - \lambda} - \sum_{n=1}^{\infty} \phi(\xi_n) \frac{n^2 \pi^2}{\lambda_n - \lambda} \prod_{m \neq n} \frac{m^2 \pi^2}{\xi_m - \xi_n},$$

which is the assertion of the lemma. \square

We turn to the main result of this section.

THEOREM 2.2. *For all $q \in E_0$, the Jacobian $d_q \sigma: E_0 \rightarrow \ell^2$ is an isomorphism onto ℓ^2 .*

Proof. See Chapter 2 of [6] for the facts used in this proof.

The components of $d_q \sigma$ are

$$d_q \sigma_n = d_q \mu_n - d_q \nu_n = g_n^2 - h_n^2,$$

where

$$g_n^2(t) = 2 \sin^2 n\pi t + o\left(\frac{1}{n}\right)$$

and

$$h_n^2(t) = 2 \cos^2 n\pi t + o\left(\frac{1}{n}\right)$$

are the respective squares of the n -th Dirichlet and Neumann eigenfunctions. Hence the operator $d_q \sigma$ is the sum of the isomorphic Fourier series operator

$$E_0 \ni f \rightarrow (-2 \langle \cos 2n\pi t, f \rangle, n \geq 1)$$

and the compact operator

$$E_0 \ni f \rightarrow \left(\left\langle o\left(\frac{1}{n}\right), f \right\rangle, n \geq 1 \right),$$

and $d_q \sigma: E_0 \rightarrow \ell^2$ is a Fredholm operator.

When q is even the vectors $g_m^2 - 1$, $m \geq 1$, form a basis for E_0 with dual basis $-2a'_m(x)$, where

$$a_m(x) = y_1(x, \mu_m) y_2(x, \mu_m)$$

and where $y_1(x, \mu_m), y_2(x, \mu_m)$ are the fundamental solutions of (1.1) for $\lambda = \mu_m$ with

$$\begin{pmatrix} y_1(0) & y_2(0) \\ y_1'(0) & y_2'(0) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

That is, if $f, g \in E_0$, then

$$\langle f, -2a'_m \rangle \in \ell^2$$

$$\langle g_m^2 - 1, g \rangle \in \ell^2$$

and

$$\int fg \, dx = \sum_{m=1}^{\infty} \langle f, -2a'_m \rangle \langle g_m^2 - 1, g \rangle.$$

Therefore it is sufficient to prove that the matrix

$$a_{n,m} = \langle g_n^2 - h_n^2, -2a'_m \rangle$$

is invertible in $B(\ell^2, \ell^2)$.

We have $\langle g_n^2, -2a'_m \rangle = \delta_{n,m}$, and because $a_m(0) = a_m(1) = 0$,

$$\begin{aligned} \langle -h_n^2, -2a'_m \rangle &= 2 \int h_n^2 a'_m \, dx = -4 \int h'_n h_n y_1(x, \mu_m) y_2(x, \mu_m) \, dx \\ &= \int y_1 h_n [h_n, y_2] + y_2 h_n [h_n, y_1] \, dx \end{aligned}$$

where $[f, g] = fg' - f'g$. But by (1.1),

$$\frac{d}{dx} [h_n, y_j] = (v_n - \mu_m) h_n y_j.$$

So if $v_n \neq \mu_m$, then

$$\begin{aligned} \langle -h_n^2, -2a'_m \rangle &= \frac{1}{v_n - \mu_m} ([h_n, y_1][h_n, y_2])|_0^1 \\ &= \frac{1}{v_n - \mu_m} h_n^2(1) y_1'(1) y_2'(1) = \frac{(-1)^m}{v_n - \mu_m} h_n^2(1) y_1'(1, \mu_m) \end{aligned}$$

since $h'_n(0) = h'_n(1) = 0$, since $y'_1(0) = 0$ and since, when q is even, $y'_2(1, \mu_m) = (-1)^m$. Also

$$h_n^2(1) = \frac{y_1^2(1, \nu_n)}{\|y_1(\cdot, \nu_n)\|_2^2} = \frac{(-1)^{n+1}}{\dot{y}'_1(1, \nu_n)}$$

where $\dot{y} = \partial y / \partial \lambda$, because $y_1(1, \nu_n) = (-1)^n$ when q is even and because $\|y_1(\cdot, \nu_n)\|_2^2 = -\dot{y}'_1(1, \nu_n)y_1(1, \nu_n)$. From the product formulas

$$y'_1(1, \mu_m) = (\nu_0 - \mu_m) \prod_{k \geq 1} \frac{\nu_k - \mu_m}{k^2 \pi^2}$$

$$\dot{y}'_1(1, \nu_n) = \frac{-(\nu_0 - \nu_n)}{n^2 \pi^2} \prod_{1 \leq k \neq n} \frac{\nu_k - \nu_n}{k^2 \pi^2}$$

we conclude that

$$\langle -h_n^2, -2a'_m \rangle = (-1)^{n+m} \prod_{0 \leq k \neq n} \frac{\nu_k - \mu_m}{\nu_k - \nu_n} \quad (2.1)$$

when $\nu_n \neq \mu_m$. If $\nu_n = \mu_m$, then $n = m$ and $[h_n, y_j] = \|y_1\|_2^{-1} \delta_{j,2}$ because the Wronskian $[y_1, y_2] = 1$. Consequently $\langle -h_n^2, -2a'_m \rangle = 1$ and (2.1) also holds when $\nu_n = \mu_m$. Thus our matrix is

$$a_{n,m} = \delta_{n,m} + (-1)^{n+m} \prod_{0 \leq k \neq n} \frac{\nu_k - \mu_m}{\nu_k - \nu_n},$$

and $(a_{n,m})$ is Fredholm because $d_q \sigma$ is a Fredholm operator.

By the Fredholm alternative, $d_q \sigma$ is an isomorphism of E_0 onto ℓ^2 if the transpose $(a_{m,n})$ is one-to-one. Now suppose $\tau = (\tau_n, n \geq 1) \in \ell^2$ lies in the kernel of $(a_{m,n})$. Then

$$0 = (-1)^n \frac{\tau_n}{\nu_0 - \mu_n} + \sum_{m \geq 1} (-1)^m \frac{\tau_m}{\nu_0 - \nu_m} \prod_{1 \leq k \neq m} \frac{\nu_k - \mu_n}{\nu_k - \nu_m}.$$

Consider the function

$$\phi(\lambda) = \sum_{m \geq 1} (-1)^m \frac{\tau_m}{\nu_0 - \nu_m} \prod_{1 \leq k \neq m} \frac{\nu_k - \lambda}{\nu_k - \nu_m}.$$

We will show in a moment that $\phi(\lambda)$ is an entire function of λ satisfying

$$\left| \phi(\lambda) / \frac{\sin \sqrt{\lambda}}{\sqrt{\lambda}} \right| = o(1) \quad (2.2)$$

uniformly on the circles $|\lambda|(n + 1/2)^2\pi^2$ as $n \rightarrow \infty$. But since

$$\phi(\mu_n) = (-1)^{n+1} \frac{\tau_n}{\nu_0 - \mu_n}, \quad \phi(\nu_n) = (-1)^n \frac{\tau_n}{\nu_0 - \nu_n},$$

$\phi(\xi_n) = 0$ at some point ξ_n in the n -th gap. Consequently $\phi \equiv 0$ by Lemma 2.1 and $\tau_n = 0$, $n \geq 1$. That means the transpose $(a_{m,n})$ is one-to-one and $d_q\sigma$ is an isomorphism.

It remains to prove (2.2). Since $\nu_n - n^2\pi^2 \in \ell^2$,

$$\begin{aligned} \prod_{1 \leq k \neq m} \frac{\nu_n - \lambda}{\nu_k - \nu_m} &= \left(\prod_{1 \leq k \neq m} \frac{k^2\pi^2 - \lambda}{k^2\pi^2 - m^2\pi^2} \right) \left(1 + o\left(\frac{\log n}{n}\right) \right) \\ &= \frac{m^2\pi^2}{m^2\pi^2 - \lambda} \left(\prod_{1 \leq k \neq m} \frac{k^2}{k^2 - m^2} \right) \frac{\sin \sqrt{\lambda}}{\sqrt{\lambda}} \left(1 + o\left(\frac{\log n}{n}\right) \right) \\ &= 2(-1)^{m+1} \frac{m^2\pi^2}{m^2\pi^2 - \lambda} \frac{\sin \sqrt{\lambda}}{\sqrt{\lambda}} \left(1 + o\left(\frac{\log n}{n}\right) \right) \end{aligned}$$

on $|\lambda| = (n - 1/2)^2\pi^2$. Hence for such λ ,

$$\left| \phi(\lambda) / \frac{\sin \sqrt{\lambda}}{\sqrt{\lambda}} \right| \leq \text{Const.} \sum_m \frac{\tau_m}{|m^2 - (n + 1/2)^2|} = o(1). \quad \square$$

It will be convenient to replace E_0 by

$$\mathcal{E}_0 = \{q \in E : \lambda_0(q) = 0\}.$$

Since even potentials are determined by their Dirichlet spectra and since $\mu_n(q + c) = \mu_n(q) + c$, and $\nu_n(q + c) = \nu_n(q) + c$, the map $q \rightarrow q - [q]$, where $[q] = \int_0^1 q(x) dx$, is an isomorphism from \mathcal{E}_0 to E_0 preserving signed gap lengths. Let

$$\mathcal{E}^n = \{q \in \mathcal{E}_0 : \gamma_m(q) = \sigma_m(q) = 0, m > n\}.$$

Because, by Theorem 2.2, σ is local analytic isomorphism on \mathcal{E}_0 , \mathcal{E}^n is a real analytic submanifold of \mathcal{E}_0 of dimension n .

COROLLARY 2.3. *For each $n \geq 1$, the signed gap length map is a real analytic isomorphism of \mathcal{E}^n onto \mathbf{R}^n .*

Proof. The image $\sigma(\mathcal{E}^n)$ is an open subset \mathbf{R}^n because $\sigma: \mathcal{E}_0 \rightarrow \ell^2$ is a local homeomorphism and $\mathcal{E}^n = \sigma^{-1}(\mathbf{R}^n) \cap \mathcal{E}_0$. We next show $\sigma(\mathcal{E}^n)$ is closed. The identity from [7],

$$q(t) = \lambda_0 + \sum_{m \geq 1} \{\lambda_{2m} + \lambda_{2m-1} - 2\mu_m(T_t q)\},$$

where $T_t q(x) = q(x + t)$, yields

$$|q(t)| \leq \sum_{m=1}^n \gamma_m(q), \quad q \in \mathcal{E}^n \quad (2.3)$$

Hence the preimage in \mathcal{E}^n of any compact subset of \mathbf{R}^n is bounded in L^2 . It is also weakly closed because the functions $\sigma_m(q) = \mu_m(q) - \nu_m(q)$ are weakly continuous. Thus the preimage of a compact subset of \mathbf{R}^n is a weakly compact subset of \mathcal{E}^n , and it follows that the map $\sigma: \mathcal{E}^n \rightarrow \mathbf{R}^n$ is proper and that $\sigma(\mathcal{E}^n)$ is a nonempty, closed subset of \mathbf{R}^n . Therefore σ maps \mathcal{E}^n onto \mathbf{R}^n .

Now let M be the set of points in \mathbf{R}^n having more than one preimage. Then M is open because σ is a local homeomorphism. But M is also closed. Indeed, if there are distinct points q_j and p_j in \mathcal{E}^n such that $\sigma(p_j) = \sigma(q_j) \rightarrow \sigma \in \mathbf{R}^n$, then because the map is proper there are subsequences such that $p_j \rightarrow p \in \mathcal{E}^n$ and $q_j \rightarrow q \in \mathcal{E}^n$. If $p = q$ then $p_j = q_j$ for j large because the map σ is homeomorphic on a neighborhood of p . So $p \neq q$ and M is closed. But $0 \notin M$ by (2.3). Thus $M \neq \emptyset$ and the mapping is one-to-one.

The map $\sigma: \mathcal{E}^n \rightarrow \mathbf{R}^n$ is real analytic because μ_m and ν_m are real analytic on $L^2_{\mathbf{R}}[0, 1]$. The inverse map is real analytic because $d_q \sigma$ is invertible. \square

It is now easy to show that the map σ is one-to-one on \mathcal{E}_0 (and hence on E_0).

COROLLARY 2.4. *The signed gap length map is one-to-one on \mathcal{E}_0 .*

Proof. Suppose not. Then some point $\tau \in \ell^2$ has at least two preimages. Since σ is a local homeomorphism, the same is true for each point in some neighborhood of τ , so it is also true at

$$\tau^{(N)} = (\tau_1, \dots, \tau_N, 0, 0, \dots)$$

for N sufficiently large. But that contradicts Corollary 2.3. \square

Write ℓ_k^2 for the space of sequences (a_n) with $\sum n^{2k} |a_n|^2 < \infty$. From the asymptotics for $y_2(1, \lambda, q)$ and $y_1'(1, \lambda, q)$ we have

$$\begin{aligned}\mu_n(q) &= n^2\pi^2 + [q] - \langle \cos 2n\pi x, q \rangle + \ell_1^2 \\ \nu_n(q) &= n^2\pi^2 + [q] + \langle \cos 2n\pi x, q \rangle + \ell_1^2.\end{aligned}$$

Hence for $q \in E_0$, $\sigma_n(q) \in \ell_1^2$ if and only if $\langle \cos 2n\pi x, q \rangle + \ell_1^2$; i.e. if and only if q is in the Sobolev space

$$H^1 = \{q \in L_{\mathbf{R}}^2[0, 1] : q' \in L_{\mathbf{R}}^2[0, 1]\}.$$

THEOREM 2.5. *The signed gap length map from $E_0 \cap H^1$ to ℓ_1^2 is one-to-one and onto.*

Proof. By Corollary 2.4 σ is one-to-one. To prove it is onto fix $\tau \in \ell_1^2$ and let $\tau^{(N)} = (\tau_1, \tau_2, \dots, \tau_N, 0, 0, \dots)$. By Corollary 2.3 there is $q_N \in \varepsilon^N$ such that $\sigma(q_N) = \tau^{(N)}$, and by (2.3)

$$|q_N(t)| \leq \sum_{n=1}^N |\tau_n| \leq \left(\sum_{n=1}^N n^2 \tau_n^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} n^{-2} \right)^{1/2},$$

so that $\|q_N\|_2 \leq \text{Const.} \|\tau\|_{\ell_1^2}$. Let $q \in \varepsilon_0$ be a weak limit of the sequence $\{q_N\}$. Then

$$\sigma_n(q - [q]) = \tau_n$$

for all n , and $q - [q] \in H^1 \cap E_0$ since $\tau \in \ell_1^2$. \square

Remark 2.6. We are unable to prove the full result that σ maps E_0 onto ℓ^2 by this method. What is needed is an estimate of $\|q\|_2$ in terms of $\gamma_n(q)$ more powerful than (2.3). Such an estimate should be useful for other problems.

Remark 2.7. It is possible, by refining the proof of Theorem 2.2, to show that $\sigma : E_0 \cap H^1 \rightarrow \ell_1^2$ is an analytic isomorphism. We omit the details.

Remark 2.8. It is known [3, p. 534] that $\gamma_n(q) \in \ell_k^2$ if and only if $q \in H^k$, i.e. if and only if q has k derivatives which are periodic and lie in $L_{\mathbf{R}}^2[0, 1]$. Thus the proof of Theorem 2.5 shows that

$$\sigma : E_0 \cap H^k \rightarrow \ell_k^2$$

is one-to-one and onto. We have not verified the likely statement that this map is bianalytic.

§3. Two applications

The potential $q \in L^2_{\mathbf{R}}[0, 1]$ is called a *finite band* potential if $\gamma_n(q) = 0$ for all but finitely many n . Marčenko [2, p. 258] proved that the set of finite band potentials is norm dense in $L^2_{\mathbf{R}}[0, 1]$. Here we derive that Theorem from result (c), stated in the introduction.

THEOREM 3.1 (Marčenko). *The set of finite band potentials is norm dense in $L^2_{\mathbf{R}}[0, 1]$.*

For $q \in E$, Theorem 3.1 is immediate from results (c). To prove it for arbitrary q we need two additional theorems. Define

$$\kappa_n(q) = \log((-1)^n y'_2(1, \mu_u, q)).$$

In [6] it is proved that $\kappa_n(q) \in \ell^2_1$, i.e. that $\sum n^2 \kappa_n^2(q) < \infty$, and that the correspondence

$$q \rightarrow (\mu_n(q) - [q], \kappa_n(q))$$

is a homeomorphism from $L^2_{\mathbf{R}}[0, 1]$ onto $\ell^2 \times \ell^2_1$. That is the first theorem.

The second theorem is the description of the isospectral manifold

$$L(q) = \{p \in L^2_{\mathbf{R}}[0, 1] : \lambda_n(p) = \lambda_n(q), \text{ all } n\}$$

given in [4]. The parameters

$$\mu_n(p) \in [\lambda_{2n-1}, \lambda_{2n}]$$

and

$$\text{sign } \kappa_n(p)$$

uniquely determine $p \in L(q)$. Although true generally, this theorem will only be used for finite band potentials, and such potentials satisfy the smoothness assumptions of [4].

Proof of Theorem 3.1. Fix $q \in L^2_{\mathbf{R}}[0, 1]$. Since $\lambda_n(q + c) = \lambda_n(q)$, we may suppose $\lambda_0(q) = 0$. By result (c) there exist, for $N = 1, 2, \dots$, $e_N \in \varepsilon_0$ such that

$$\gamma_n(e_0) = \begin{cases} \gamma_n(q) & n \leq N \\ 0 & n > N \end{cases}$$

and

$$\mu_n(e_N) = \lambda_{2n-1}(e_N), \quad n = 1, 2, \dots$$

Since $\mu_n(q) \in [\lambda_{2n-1}(q), \lambda_{2n}(q)]$ there exists $t_n \in [0, 1]$ such that

$$\mu_n(q) = t_n \lambda_{2n}(q) + (1 - t_n) \lambda_{2n-1}(q),$$

and by the second theorem just cited there exists $q_N \in L(e_N)$ such that for all n ,

$$\mu_n(q_N) = t_n \lambda_{2n}(e_N) + (1 - t_n) \lambda_{2n-1}(e_N)$$

and

$$\text{sign } \kappa_n(q_N) = \text{sign } \kappa_n(q).$$

By the first cited theorem $\|q_N - q\|_2 \rightarrow 0$ if

$$\|\mu(q_N) - \mu(q)\|_{\ell^2} \rightarrow 0 \tag{3.1}$$

and

$$\|\kappa_n(q_N) - \kappa_n(q)\|_{\ell^2_1} \rightarrow 0. \tag{3.2}$$

By the second theorem there exists $e \in \mathcal{E}_0$ such that for all n ,

$$\lambda_n(e) = \lambda_n(q)$$

$$\mu_n(e) = \lambda_{2n-1}(q).$$

Then $\|\gamma_n(e_N) - \gamma_n(e)\|_{\ell^2} \rightarrow 0$ and $\sigma_n(e_N)$ and $\sigma_n(e)$ have the same sign, so that

$$\|\sigma_n(e_N) - \sigma_n(e)\|_{\ell^2} \rightarrow 0. \tag{3.3}$$

Hence by result (c), $\|e_N - e\|_2 \rightarrow 0$ and by the first theorem

$$\|\mu_n(e_N) - \mu_n(e)\|_{\ell^2} \rightarrow 0. \tag{3.4}$$

But then by the choices of q_N , e_N and e ,

$$\mu_n(q_N) - \mu_n(q) = -t_n(\sigma_n(e_N) - \sigma_n(e)) + \mu_n(e_N) - \mu_n(e),$$

and (3.3) and (3.4) imply (3.1).

To prove (3.2) we use the identity

$$2 \cosh \kappa_n(q) = (-1)^n \Delta(\mu_n(q), q),$$

where $\Delta(\lambda, q)$ is the discriminant function

$$\Delta(\lambda, q) = y_1(1, \lambda, q) + y_2'(1, \lambda, q)$$

and the inequality

$$|x - y|^2 \leq 2 |\cosh x - \cosh y|,$$

valid when x and y have the same sign. They give

$$\begin{aligned} n^2 |\kappa_n(q_N) - \kappa_n(q)|^2 &\leq n^2 |\Delta(\mu_n(q_N), q_N) - \Delta(\mu_n(q), q)| \\ &= n^2 |\Delta(\mu_n(q_N), e_N) - \Delta(\mu_n(q), e)|. \end{aligned}$$

Since $\|e_N - e\|_2 \rightarrow 0$, $\Delta(\lambda, e_N) \rightarrow \Delta(\lambda, e)$ uniformly on compact sets. Thus by (3.1)

$$|\Delta(\mu_n(q_N), e_N) - \Delta(\mu_n(q), e)| \rightarrow 0$$

for each n . Moreover,

$$\begin{aligned} &|\Delta(\mu_n(q_N), e_N) - \Delta(\mu_n(q), e)| \\ &\leq |\Delta(\mu_n(q_N), e_N) - 2(-1)^n| + |\Delta(\mu_n(q), e) - 2(-1)^n| \end{aligned}$$

since $\Delta(\lambda_{2n-1}, e_N) = \Delta(\lambda_{2n-1}, e) = 2(-1)^n$. Because $\dot{\Delta}(\lambda) = \partial \Delta / \partial \lambda$ is an entire function of order $\frac{1}{2}$, having one zero $\dot{\lambda}_n$ in each gap $[\lambda_{2n-1} \leq \lambda \leq \lambda_{2n}]$ and no other zeros, the product representation

$$\dot{\Delta}(\lambda) = \prod_{n \geq 1} \frac{\dot{\lambda}_n - \lambda}{n^2 \pi^2}$$

shows that

$$\sup_{\lambda_{2n-1} \leq \lambda \leq \lambda_{2n}} |\dot{\Delta}(\lambda, q)| \leq c \frac{\gamma_n(q)}{n^2}.$$

Hence by (3.5)

$$n^2 |\Delta(\mu_n(q_N), e_N) - \Delta(\mu_n(q), e)| \leq c\gamma_n^2(q)$$

and by dominated convergence

$$\lim_{N \rightarrow \infty} \sum_n n^2 |\kappa_n(q_N) - \kappa_n(q)|^2 = 0. \quad \square$$

Our second application concerns the subspace $L_k^2 \subset L_R^2[0, 1]$ of all functions whose periodic extensions have primitive period $1/k$, $k = 1, 2, \dots$. First of all, if $q \in L_k^2$, $\gamma_n(q) = 0$ whenever $k \nmid n$. To see this, recall from [7] that $\mu_n(t) = \mu_n(T_t q)$, where $T_t q(x) = q(x - t)$, then as t runs from 0 to 1, $\mu_n(t)$ makes n complete trips between $\lambda_{2n-1}(q)$ and $\lambda_{2n}(q)$, when $\gamma_n(q) \neq 0$. By assumption, $\mu_n(t + 1/k) = \mu_n(t)$. Therefore, n is equal to k times the number of complete trips in time $1/k$, and so k divides n , $k \mid n$.

Now let $E_0^k = E_0 \cap L_k^2$. It follows from the observation above that the restriction of σ to E_0^k maps into $\ell^2(k) = \{\sigma \in \ell^2 : \sigma_n = 0 \text{ whenever } k \nmid n\}$. Without any change in the argument of [1] or that of Section 2, one can show that σ is a real analytic isomorphism between E_0^k and $\ell^2(k)$ or between $E_0^k \cap H^1$ and $\ell_1^2(k)$.

Suppose $q \in E$ and $\sigma(q) \in \ell^2(k)$. Then there is a $p \in E_0^k$ such that $\sigma(p) = \sigma(q)$. However, σ is globally one-to-one on E_0 so that $p = q - [q]$. In other words, $q \in E_0$ has primitive period $1/k$ if and only if $\gamma_n(q) = 0$ whenever $k \nmid n$.

It is easy to extend this observation to all of $L_R^2[0, 1]$. Let $q \in L_R^2[0, 1]$ and $L(q) = \{r \in L_R^2[0, 1] \mid \lambda_i(r) = \lambda_i(q) \text{ for } i \geq 0\}$, i.e., the isospectral set of q . It is not hard to see that $L(q) \cap E \neq \emptyset$ and that all points in $L(q)$ have the same primitive period. See [4]. Thus we have proved

THEOREM 3.2. *The potential q has primitive period $1/k$ if and only if $\gamma_n(q) = 0$ whenever k does not divide n .*

§4. Band lengths

Let $\alpha_n(q) = \alpha_n = \lambda_{2n-1} - \lambda_{2n}$ be the length of the n -th band. It is well known that

$$\alpha_n(q) - (2n - 1)\pi^2 \in \ell^2 \tag{4.1}$$

and in [1] and [5] it was shown that

$$\tilde{\alpha}_n = (2n - 1)\pi^2 - \alpha_n \geq 0, \quad (4.2)$$

with equality holding for some n if and only if q is constant.

THEOREM 4.1. *For all n and all q ,*

$$\alpha_n - \alpha_{n-1} + \alpha_{n-2} \mp \cdots > 0, \quad (4.3)$$

and

$$\beta_n = \tilde{\alpha}_n - \tilde{\alpha}_{n-1} + \tilde{\alpha}_{n-2} \mp \cdots \geq 0. \quad (4.4)$$

Moreover, if $\beta_n = 0$ for some n , then q is constant and $\beta_k = 0$ for all k .

Note that by (4.1), (4.3) has content only for small n . By (4.4) $\beta_n \leq \tilde{\alpha}_n$, so that by (4.1)

$$\beta_n \in \ell^2_+$$

and by (4.4) and (4.3),

$$0 \leq \beta_n < n\pi^2. \quad (4.5)$$

We shall show that (4.5) is sharp for every n and that, properly interpreted, the Jacobian $d_g \beta_n : E_0 \rightarrow \ell^2$ is invertible at $q = 0$. A simple characterization of band lengths thus seems unlikely.

Proof. Recall from [1] that there exists $h_n = h_n(q) \geq 0$, such that $\sum n^2 h_n^2 < \infty$, and such that

$$\delta(\lambda, q) = \cos^{-1} \left(\frac{\Delta(\lambda, q)}{2} \right)$$

is a conformal mapping from the half plane $\{\text{Im } \lambda > 0\}$ onto the slit quarter plane

$$\Omega(h) = \{x > 0, y > 0\} \setminus \bigcup_{n=1}^{\infty} T_n$$

where

$$T_n = \{n\pi + iy : 0 < y \leq h_n\}.$$

Under $\delta(\lambda, q)$ the n -th band is mapped onto the segment

$$B_n = [(n-1)\pi-, n\pi+] \subset \partial\Omega(h),$$

and if

$$u_n(z, h) = \omega(z, B_n, \Omega(h)) = u_n(z)$$

is the harmonic measure of B_n in $\Omega(h)$, then

$$\alpha_n = \lim_{x \rightarrow \infty} 2\pi x^2 u_n(x + ix, h). \quad (4.6)$$

Let $k \leq n$ and let $z = x + ix$ with $x > n\pi$. Then $u_k(z)$ is the probability that a Brownian path starting at z makes its first exit from $\Omega(h)$ through B_k . Letting S_k be the set of such paths, we write

$$u_k(z) = P_z(S_k)$$

Brownian paths can be assumed continuous. Thus every path in S_k must cross the half line $J_k = \{x = k\pi, y > 0\}$ before it leaves $\Omega(h)$. Let R_k be those paths in S_k which, before leaving $\Omega(h)$, last meet $J_k \cup J_{k-1}$ in J_k , and let L_k be those whose last contact with $J_k \cup J_{k-1}$, before departing from $\Omega(h)$, is in J_{k-1} . Then R_k and L_k are P_z measurable, $R_k \cap L_k = \emptyset$ and

$$P_z(S_k) = P_z(R_k) + P_z(L_k).$$

But

$$P_z(L_k) = P_z(R_{k-1})$$

by a reflection. Since $L_1 = \emptyset$, we conclude that

$$u_n(z) - u_{n-1}(z) \pm \cdots = P_z(R_n) > 0$$

which by (4.6) yields (4.3). To prove (4.4), let

$$V_n(z) = u_n(z, 0) - u_n(z, h).$$

Then

$$\tilde{\alpha}_n = \lim_{x \rightarrow \infty} 2\pi x^2 V_n(x + ix).$$

On $\partial\Omega(h)$,

$$V_n(\zeta) = \sum_{k=1}^{\infty} u_n(\zeta, 0) \chi_{T_k}(\zeta), \quad (4.7)$$

and the argument above shows

$$\sum_{k=1}^n (-1)^{n-k} u_k(\zeta, 0) > 0$$

on $\bigcup T_k$. Hence for x large

$$\sum_{k=t}^n (-1)^{n-k} V_k(x + ix) \geq 0$$

and (4.4) holds. If equality holds in (4.4) then by (4.7), $\bigcup T_k$ has zero harmonic measure in $\Omega(h)$. That means all gap lengths are zero and q is constant. \square

To see that (4.5) is sharp, note that $q \rightarrow h_n(q)$ maps onto ℓ_1^2 and that by (4.6),

$$\lim_{h_n \rightarrow \infty} \alpha_k = 0, \quad 1 \leq k \leq n.$$

For $q \in E$, define

$$a_n(q) = \mu_n(q) - \nu_{n-1}(q), \quad n \geq 1$$

and

$$b_n(q) = \sum_{k=1}^n (-1)^{n-k} ((2n-1)\pi^2 - a_n(q)) = n\pi^2 - \sum_{k=1}^n (-1)^{n-k} a_n(q).$$

Then for $q \in E$

$$b_n(q) = \beta_n(q) + \text{Max}(\sigma_n(q), 0),$$

and for each potential $q \in L^2$ there is $q^+ \in E$ with $\mu_n(q^+) \leq \nu_n(q^+)$ and $\lambda_n(q^+) = \lambda_n(q)$, so that $b_n(q^+) = \beta_n(q)$.

THEOREM 4.2. *At $q=0$ the Jacobian $d_q(b_n): E_0 \rightarrow \ell^2$ is an isomorphism onto ℓ^2 .*

Proof. At $q = 0$, $f \in E_0$,

$$\langle d_q a_n, f \rangle = \langle 2 \sin^2 n\pi t - 2 \cos^2 (n-1)\pi t, f \rangle$$

and

$$\langle d_q b_n, f \rangle = -2 \langle \sin^2 n\pi t, f \rangle = \langle \cos 2n\pi t, f \rangle,$$

and $(\cos 2n\pi t)_{n \geq 1}$ is a complete orthonormal system in E_0 . \square

THEOREM 4.3. (a) *If q and \tilde{q} are finite band potentials and if $\alpha_n(q) = \alpha_n(\tilde{q})$ for infinitely many n , then the periodic spectra of q and \tilde{q} agree up to a translation.*

(b) *If q and \tilde{q} are real analytic, and if $\alpha_n(q) = \alpha_n(\tilde{q})$ for all large n , then q and \tilde{q} have the same periodic spectrum up to a translation.*

Proof. Let $\phi(z, q)$ be the inverse of the mapping $\delta(\lambda, q)$. If q is a finite band potential then $h_n(q) = 0$, $n > N$ and $\phi(z, q)$ reflects to be analytic in the complement of the finite union of vertical slits $\{|x| = n\pi, |y| \leq h_n(q), 1 \leq n \leq N\}$. For z large we have

$$\phi(z, q) = z^2 + o\left(\frac{1}{z}\right).$$

By the hypothesis of (a),

$$\phi(z + \pi, q) - \phi(z, q) = \phi(z + \pi, \tilde{q}) - \phi(z, \tilde{q}) \quad (4.8)$$

holds for an infinite sequence of integers tending to ∞ . Hence (4.8) holds for all z , and $\phi(z, q)$ and $\phi(z, \tilde{q})$ have the same singularities. Therefore $h_n(q) = h_n(\tilde{q})$ for all n , which means the spectra of q and \tilde{q} differ by at most a translation.

To prove (b), set $f(z) = \phi(z, q) - \phi(z, \tilde{q})$. By reflection $f(z)$ is analytic in

$$\Omega^* = \mathbf{C} \setminus \bigcup_{n=1}^{\infty} (S_n \cup S_{-n})$$

where $S_n = \{x = n\pi, |y| \leq \max(h_{|n|}(q), h_{|n|}(\tilde{q}))\}$, and by the asymptotics for $\Delta(\lambda, q)$, $f(z)$ is bounded on Ω^* . Since q and \tilde{q} are real analytic, we have by [7],

$$\begin{aligned} \max(h_n(q), h_n(\tilde{q})) &\leq C \max(\gamma_n(q), \gamma_n(\tilde{q})) \\ &\leq C e^{-an} \end{aligned}$$

for constants a and C . Viewing S_n as two-sided, we see that $f(z)$ has continuous boundary values on S_n and that for $n \geq 1$,

$$\sup_{z \in \bar{S}_n} |f(z) - f(n\pi+)| \leq \gamma_n(q) + \gamma_n(\bar{q}) \leq Ce^{-an}.$$

By hypothesis there is N so that

$$f((n+1)\pi-) - f(n\pi+) = \alpha_n(q) - \alpha_n(\bar{q}) = 0$$

for $n \geq N$, and hence

$$\sup_{z \in \bar{S}_n} |f(z)| \leq Ce^{-an}, \quad n \geq N. \quad (4.9)$$

Set $h^* = \sup_n \{h_n(q), h_n(\bar{q})\}$. We shall prove

$$|f(x + Ch^*)| \leq Ce^{-a'x}, \quad x > x_0 \quad (4.10)$$

First assume (4.10). Then because $f(z)$ is bounded and analytic in $\{y > h^*\}$,

$$\log |f(z)| \leq C_1 + C_2 \int_{x > x_0} \frac{-a'x}{1+x^2} dx = -\infty$$

on $|z - i(h^* + 1)| < \frac{1}{2}$. Therefore $f = 0$ and (b) is proved.

We turn to the proof of (4.10). Let Δ_n be the disc $\{|z - n\pi| < 2Ae^{-a|n|}\}$ with A so large that $\text{dist}(S_n, \partial\Delta_n) \geq Ae^{-an}$ for all $n \geq 1$. Then by (4.9) and the three circles theorem,

$$\sup_{\partial\Delta_n} |f(z)| \leq C_2 e^{-an} \quad (4.11)$$

Now let $\Omega = \mathbb{C} \setminus \bigcup_{n=1}^{\infty} (\bar{\Delta}_n \cup \bar{\Delta}_{-n})$ and for $\delta > 0$ fixed and x large, set

$$E_x = \bigcup \{\Delta_n : |n\pi - x| < \delta x\}.$$

LEMMA 4.4. *There is $C(h^*, a)$ such that for x large,*

$$\omega(x + ih^*, E_x, \Omega) \geq C(h^*, a)$$

Note that by the subharmonicity of $\log |f|$, this lemma and (4.11) imply (4.10) and hence the theorem.

Proof of Lemma 4.9. Fix δ_1 , $0 < \delta_1 < \delta$, to be determined later, and let $N_x \sim 2\delta_1 x / \pi$ be the number of n such that $|n\pi - x| < \delta_1 x$. Set

$$u(z) = \frac{1}{N_x} \sum_{|n\pi - x| < \delta_1 x} \log \frac{1}{|z - n\pi|}.$$

Then $u(z)$ is harmonic and bounded above in Ω , and

$$\sup_{z \in \partial\Omega \setminus E_x} u(z) \leq \log \frac{1}{(\delta - \delta_1)x} + c = \alpha.$$

But if $z \in E_x$ then

$$\begin{aligned} u(z) &\leq \frac{1}{N_x} \log (Ae^{-an}) + \frac{1}{N_x} \sum_{i=2}^{N_x/2} \log \frac{1}{i\pi} \\ &\leq \log \frac{1}{\delta_1 x} + c'(a) = \beta, \end{aligned}$$

and by a similar calculation,

$$u(x + ih^*) \geq \beta + c(h^*, a).$$

We choose δ_1 so that $\beta - \alpha = c'' > 1$. Then by the maximum principle,

$$\omega(z, E_x, \Omega) \geq \frac{u(z) - \alpha}{\beta - \alpha}$$

and

$$\omega(x + ih^*, E_x, \Omega) \geq \frac{c(h^*, a)}{c''} \quad \square.$$

We thank L. Carleson and P. Jones for suggesting the proof of Lemma 4.4, which actually shows that E_x has logarithmic capacity comparable to its diameter.

REFERENCES

- [1] J. GARNETT and E. TRUBOWITZ, "*Gaps and bands of one dimensional periodic Schrödinger operators*," Comment. Math. Helv. 59 (1984), 258–312.
- [2] V. A. MARČENKO, *Sturm Liouville Operators and Applications*, Kiev 1977 (Russian).
- [3] V. A. MARČENKO and I. O. OSTROVSKII, *A Characterization of the spectrum of Hill's operator*, Math. USSR-Sbornik, 97 (1975), 493–554.
- [4] H. P. MCKEAN and E. TRUBOWITZ, *Hill's operator and Hyperelliptic function theory in the presence of infinitely many branch points*, Comm. Pure Appl. Math 29 (1976), 143–226.
- [5] J. MOSER, *Integrable Hamiltonian Systems and Spectral Theory*, Lezioni Fermiane, Scuola Normale Superiore, Pisa, 1981.
- [6] J. POSCHEL and E. TRUBOWITZ, *Lectures on Inverse Spectral Theory*, (to appear).
- [7] E. TRUBOWITZ, *The Inverse Problem for Periodic Potentials* Comm, Pure Appl. Math. 30 (1977), 321–337.

University of California, Los Angeles
and
E.T.H. Zurich, Switzerland

Received April 23, 1986