

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 63 (1988)

Artikel: Normal subgroups of classical groups over Banach algebras.
Autor: Vaserstein, Leonid N.
DOI: <https://doi.org/10.5169/seals-48197>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Normal subgroups of classical groups over Banach algebras

LEONID N. VASERSTEIN

Introduction

All subgroups H of the general linear group $GL_n A$, $n \geq 3$, over a Banach algebra A with 1, which are normalized by elementary matrices were described in [2] (when $n = 2$, [2] shows that the situation becomes more complicated and gives partial results). In this paper, we obtain similar result for unitary groups. In the next two paragraphs we give Wall's [4] definition of unitary groups which include symplectic and orthogonal groups.

Let A be an associative ring with 1, $*: A \rightarrow A$ an anti-automorphism of A (i.e. $*$ is a bijection on A such that $(x - y)^* = x^* - y^*$ and $(xy)^* = y^*x^*$ for all x and y in A). We assume that $x^{**} = \varepsilon x \varepsilon^*$ for some unit $\varepsilon = \varepsilon^{*-1}$ of A and every x in A .

Set $F_n = e_{1,2} + \varepsilon e_{2,1} + \cdots + e_{2n-1,2n} + \varepsilon e_{2n,2n-1} \in GL_{2n} A$, where $e_{i,j}$ are the matrix units. Let $U_{2n}(A, *, \varepsilon)$ or just $U_{2n} A$ for short denote the group of all g in $GL_{2n} A$ such that $g^* F_n g = F_n$ ($*$ is extended to anti-automorphisms of the matrix rings in the usual way: $(g^*)_{i,j} = (g_{j,i})^*$). Since $F_n^* = F_n^{-1}$, $F_n \in U_{2n} A$ and $U_{2n} A$ is invariant under $*$.

Here are classical examples.

EXAMPLE 1. $*$ is identical on A and $\varepsilon = -1$. Then A is commutative and $U_{2n} A$ is the standard symplectic group $Sp_{2n} A$.

EXAMPLE 2. $*$ are as in Example 1, but $\varepsilon = 1$. If 2 is not a zero divisor in A , then $U_{2n} A$ is the orthogonal group $O_{2n} A$ of a quadratic form in $2n$ variables of Witt index n .

EXAMPLE 3. A be the complex numbers, $*$ the complex conjugation, $\varepsilon = 1$ or -1 . Then $U_{2n} A$ is isomorphic to the standard unitary group.

EXAMPLE 4. $A = D \times D^{\text{op}}$, where D is an associative ring with 1, D^{op} is the opposite ring, and $(d, d')^* = (d', d)$ for any (d, d') in A . Then, for any ε , $U_{2n} A$ is isomorphic to $GL_{2n} D$.

The unitary group $U_{2n}A$ is a normal subgroup of the group $GU_{2n}A$ of unitary *similitudes*, i.e. of all matrices g in $GL_{2n}A$ such that $g^*F_ngF_n^{-1}$ is a scalar matrix over the center of A (the scalar matrix depends on g , and it is 1_{2n} if and only if g is in $U_{2n}A$). For any ideal $B = B^*$ of A , let $GU_{2n}(A, B)$ denote the group of g in $GU_{2n}A$ which reduce to scalar matrices over the center of A/B modulo B . This is a normal subgroup of $GU_{2n}A$.

Now we will define elementary unitary matrices. First, we define a bijection ' of the natural numbers by $(2i)' = 2i - 1$ and $(2i - 1)' = 2i$ for any integer $i \geq 1$. For any a in A and any integers i, j such that $1 \leq i \neq j \leq 2n$ we set $E_{i,j}(a) = 1_{2n} + ae_{i,j} - \varepsilon^c a^* \varepsilon^{c'} e_{j',i'}$, where $c = ((-1)^i - 1)/2$ and $c' = (1 - (-1)^j)/2$. It is easy to check that all these $E_{i,j}(a)$ belong to $U_{2n}A$. In particular, $E_{2k-1,2k}(a) = 1_{2n} + (a - \varepsilon^* a^*)e_{2k-1,2k}$ and $E_{2k,2k-1}(a) = 1_{2n} + (a - a^* \varepsilon)e_{2k,2k-1}$.

For any ideal $B = B^*$ of A , let $EU_{2n}B$ denote the subgroup of $GU_{2n}B$ generated by all $E_{i,j}(b)$ with b in B , and let $EU_{2n}(A, B)$ denote the normal subgroup of $EU_{2n}A$ generated by $EU_{2n}B$.

THEOREM. *Suppose that A is a Banach algebra with 1 and $n \geq 2$. Then,*

(1) $EU_{2n}(A, B) = [EU_{2n}A, EU_{2n}B] = [EU_{2n}A, GU_{2n}(A, B)] = [EU_{2n}(A, B), GU_{2n}A]$ for any ideal $B = B^*$ of A .

So for every subgroup H of $GU_{2n}(A, B)$ containing $EU_{2n}(A, B)$ we have $[H, EU_{2n}A] = EU_{2n}(A, B)$, hence H is normalized by $EU_{2n}A$. Conversely, if $n \geq 3$, then

(2) *for any subgroup H of $GU_{2n}A$ which is normalized by $EU_{2n}A$ there is an ideal $B = B^*$ such that $EU_{2n}(A, B) \subset H \subset GU_{2n}(A, B)$.*

Remarks. 1. When $n = 1$, (2) fails in the case of ordinary orthogonal groups, because then the group EU_2A is trivial. The conclusions (1) with $n = 1$ and (2) with $n = 2$ hold under additional conditions on A , $*$, ε (for example, when A is commutative and $\varepsilon = \pm 1$), but the situation in general is unclear.

2. When $*$ is the identity (so A is commutative) and $\varepsilon = 1$ or -1 , our theorem is contained in results of [3].

3. When A has no proper ideals $B = B^*$ (for example, A is simple), our theorem says that the group $EU_{2n}A$ modulo its center $GU_{2n}(A, 0) \cap EU_{2n}A$ is simple for $n \geq 3$. Compare this with results of [1] about simplicity of unitary groups over some factors A .

4. The group $EU_{2n}(A, B)$ is contained in the identity component $GU_{2n}(A, B)^0$ of $GU_{2n}(A, B)$. On the other hand, this component is contained in the subgroup $GEU_{2n}(A, B)$ of $GU_{2n}(A, B)$ generated by $EU_{2n}(A, B)$ and diagonal matrices. Therefore, when $n \geq 2$, $EU_{2n}(A, B) = [(GU_{2n}A)^0, GU_{2n}(A, B)^0]$.

Proof of (1)

Evidently, $EU_{2n}(A, B) \supset [EU_{2n}A, EU_{2n}B]$. Let us prove the inverse inclusion, i.e. that every elementary unitary matrix $E_{i,j}(a)$ in $EU_{2n}B$ belongs to the commutator subgroup $[EU_{2n}A, EU_{2n}B]$. When $i \neq j'$, $E_{i,j}(a)$ is the image of an elementary matrix under a monomorphism $s: GL_n A \rightarrow U_{2n}A$ such that $s(E_n A) \subset EU_{2n}A$ and $s(E_n B) \subset EU_{2n}B$. (For a monomial matrix f in $GL_n A$, depending on i and j , with a non-zero entry 1 or ε in each row and column, we have

$$F_n = f^* \begin{pmatrix} 0 & 1_n \\ \varepsilon 1_n & 0 \end{pmatrix} f \quad \text{and} \quad s(g) = f^{-1} \begin{pmatrix} g & 0 \\ 0 & g^{*-1} \end{pmatrix} f \quad \text{for all } g \text{ in } GL_n A.$$

Since $E_n B \subset [E_n A, E_n B]$ (see [2]), $E_{i,j}(a) \in s([E_n A, E_n B]) \subset [EU_{2n}A, EU_{2n}B]$.)

When $i = j'$, we pick an integer k in the interval $1 \leq k \leq 2n$ such that $k \neq i$ and $k + i$ is even. Then $E_{i,j}(a) = E_{k,j}(x)[E_{k,k'}(-a), E_{i,k}(-1)] \in [EU_{2n}A, EU_{2n}B]$, where $x := a - a^*a^*$. (By our definition, $[g, h] := ghg^{-1}h^{-1}$, so $[E_{k,k'}(-a), E_{i,k}(-1)] = 1_{2n} - xe_{k,i} - xe_{i,k} + xe_{i,i'}$.)

The first equality in (1) is proved.

To prove the second one, pick an arbitrary g in $GU_{2n}(A, B)$. For any elementary $E_{i,j}(a)$ in $EU_{2n}A$ any rational number r , we set $h(r) = [g, E_{i,j}(ra)]$. We want to prove that $h(1) \in EU_{2n}(A, B)$. When r is close to 0, $h(r)$ is close to $h(0) = 1_{2n}$, hence it is the product of a diagonal matrix from $U_{2n}B$ and a matrix from $EU_{2n}B$. Let $GEU_{2n}B$ be the group of all such products. Evidently, it is normalized by $EU_{2n}A$. So $h(1) \in GEU_{2n}B$. Hence $\varphi(h') = [g, h'] \in GEU_{2n}B$ for any h' in $EU_{2n}A$. Note that $h' \rightarrow \varphi(h')EU_{2n}(A, B)$ is a group homomorphism from $EU_{2n}A$ to $GEU_{2n}B/EU_{2n}(A, B)$. Since the first group here is perfect (see above) and the second group is commutative (because it is a factor group of the group $GE_n B/E_n(A, B)$ which is commutative by the Whitehead lemma that allows us to permute diagonal matrices modulo elementary matrices provided that $n \geq 2$), we conclude that the homomorphism is trivial. That is, $\varphi(h') \in EU_{2n}(A, B)$ for all h' in $EU_{2n}A$. In particular, when $h' = E_{i,j}(a)$, we obtain that $h(1)$ is in $EU_{2n}(A, B)$. The second equality in (1) is proved.

Using this with $B = A$, we conclude that $EU_{2n}A = EU_{2n}(A, A)$ is a normal subgroup of $GU_{2n}A = GU_{2n}(A, A)$. Since $GU_{2n}(A, B)$ is also a normal subgroup of $GU_{2n}A$, we conclude that $EU_{2n}(A, B) = [EU_{2n}A, GU_{2n}(A, B)]$ is normal in $GU_{2n}A$ too. That is, we obtain the third equality in (1).

Proof of (2)

Let H be a subgroup of $GU_{2n}A$, and let H be normalized by $EU_{2n}A$. For any integers i and j such that $1 \leq i \neq j \leq 2n$, we set $X_{i,j} = \{a \in A : E_{i,j}(a) \in H\}$. Clearly,

they are additive subgroups of A . The identity of the form $[E_{i,j}(a), E_{j,k}(b)] = E_{i,k}(ab)$, where $i \neq j \neq i'$, $k \neq j \neq k'$, and $k \neq j \neq k'$, show that $X_{i,j} = X_{1,3}$ whenever $i \neq j$ and that $X_{1,3} =: B$ is an ideal of A . Since $X_{1,3} = (X_{4,2})^*$, $B = B^*$. Now it is easy to check that $X_{2i-1,2i} = (X_{2i,2i-1})^* = \{b \in B : b^* = -\varepsilon b\}$ for $i = 1, \dots, n$.

Let us show now that the image of H modulo B belongs to the center of $GU_{2n}(A/B)$. Otherwise, an element g of H does not commute with an elementary unitary matrix (note that the centralizer of $EU_{2n}(A/B)$ in $GU_{2n}(A/B)$ consists of scalar matrices over the center of A/B). Then g does not commute with an elementary unitary matrix $E_{i,j}(a)$ modulo $GU_{2n}(A, B)$ (otherwise we would obtain a non-trivial homomorphism from the perfect group $EU_{2n}A$ to the commutative group $GU_{2n}(A, B)/GU_{2n}B$). So $h = [g, E_{i,j}(a/N)] \in H$ and h is outside of $GU_{2n}(A, B)$ for any natural number N . Taking a large N , we obtain a matrix h in H outside $GU_{2n}(A, B)$ which is arbitrarily close to the identity matrix 1_{2n} .

Now, after a permutation of the basis, we will think about $U_{2n}A$ as $U_2(M_nA)$, where M_nA is the ring of n by n matrices over A . Since h is close to the identity, we can write $h = E_{2,1}(c) \text{diag}(d, d^{*-1})E_{1,2}(b)$ where $d \in GL_nA$, $B = -\varepsilon^*b^* \in M_nA$, $c = -c^*\varepsilon \in M_nA$. If $c \notin M_nB$, we can replace h by $b'^{1,2}h(-b')^{1,2}$ with a matrix $b' = -\varepsilon^*b'^*$ in M_nA (which results in replacing d by $d + b'c$) to get a non-diagonal entry of d outside B . Moreover the matrix b' above can be taken to be small, so the d -entry stays invertible. Similarly, if $b \notin M_nB$, we can replace h by $(-c')^{2,1}hc'^{2,1}$ with $c' = -c'^*\varepsilon$ in M_nA to reach the same objective.

Thus, we can assume that H contains an element of the form $h = E_{2,1}(c) \text{diag}(d, d^{*-1})E_{1,2}(b)$ with $d \in GL_nA \setminus G_n(A, B)$ (i.e. the image of d in $GL_n(A/B)$ is not a scalar matrix over the center of A/B).

To complete our proof we need to show that H contains an elementary unitary matrix outside $GU_{2n}(A, B)$.

Consider the set T of all triples $(c', d', b') \in M_nA \times GL_nA \times M_nA$ such that $t(c', d', b') := E_{2,1}(c') \text{diag}(d', d'^{*-1})E_{1,2}(b') \in H$. Note that if $(c', d', b'), (c'', d'', b'') \in T$, then $(c' - c'', d'd''^{-1}, d''(b' - b'')d''^*)$, $(d''^{-1}(c' - c'')d''^{*-1}, d''^{-1}d', d' - d'') \in T$. Indeed, $E_{2,1}(c'')^{-1}(t(c', d', b')t(c'', d'', b'')^{-1}E_{2,1}(c'') = t((c' - c'', d'd''^{-1}, d''(b' - b'')d''^*) \in H$ and $E_{2,1}(c'')(t(c', d', b')^{-1}t(c'', d'', b'')E_{2,1}(c'')^{-1} = t(d''^{-1}(c' - c'')d''^{*-1}, d''^{-1}d', d' - d'') \in H$. Therefore, the projection of $T \subset M_nA \times GL_nA \times M_nA$ on each of 3 factors is a subgroup there.

Since the set $t(T)$ is normalized by all elements of the form $\text{diag}(u, u^{-1})$, where $u \in E_nA$, the group E_nA acts on T . We use this action to define an operation $[,]' : E_nA \times T \rightarrow T$ as follows: $[u, (c', d', b')]' = (u^{*-1}c'u^{-1} - c', [u, d'], d'(ub'u^* - b')d'^*)$, where $u \in E_nA$ and $(c', d', b') \in T$.

Note that the second projection H' of T is a subgroup of GL_nA and H' is

normalized by $E_n A$. By [2], H' contains an elementary matrix $z^{1,2} = 1_n + ze_{1,2}$ with z outside of B . So we can assume that $d = z^{1,2}$ with z in $A \setminus B$. We will need no conditions on A anymore.

We have $[1^{3,1}, (c, d, b)]' = (c', z^{3,2}, \cdot) = x_1 \in T$ with the matrix $c' = c - (-1)^{1,3}c(-1^{3,1})$ in $M_n A$ having non-zero entries only in the first column and row.

Next we consider the triple $x_2 = [(-1)^{2,1}x_1]' = (c'', z^{3,1}, \cdot)$ in T . Here $c'' = 1^{1,2}c'(1^{2,1}) - c'$ can have non-zero entries only at position $(1, 1)$.

Set $x_3 = [1^{2,3}, x_2]' = (0, z^{2,1}, \cdot) \in H$. Then

$$\begin{aligned} [t(x_3), (e_{1,3} - \varepsilon^* e_{3,1})^{1,2}] &= [\text{diag}(z^{2,1}, (-z^*)^{1,2}), (e_{1,3} - \varepsilon^* e_{3,1})^{1,2}] \\ &= (z^{2,1}(e_{1,3} - \varepsilon^* e_{3,1})z^{*1,2} - (e_{1,3} - \varepsilon^* e_{3,1})^{1,2} \\ &= (ze_{2,3} - \varepsilon^* z^* e_{3,2})^{1,2} \in H. \end{aligned}$$

Thus, H contains the elementary unitary matrix $t(0, 1_n, ze_{2,3} - \varepsilon^* z^* e_{3,2}) = (ze_{2,3} - \varepsilon^* z^* e_{3,2})^{1,2}$ (which is $E_{3,4}(z)$ in the original notation) outside $GU_{2n}(A, B)$. Our proof is completed.

Here are some entries of the first 3 columns of the matrices h , $t(x_1)$, $t(x_2)$, $t(x_3)$ (namely, the positions (i, j) of the c - and d -parts with $1 \leq i, j \leq 3$).

1	z	0	1	0	0	1	0	0	1	0	0
0	1	0	0	1	0	0	1	0	z	1	0
0	0	1	0	z	1	0	0	1	0	0	1
\cdot	\cdot	\cdot	\cdot	\cdot	\cdot	z	0	1	0	0	0
\cdot	\cdot	\cdot	\cdot	0	0	0	0	0	0	0	0
\cdot	\cdot	\cdot	\cdot	0	0	0	0	0	0	0	0

REFERENCES

- [1] PIERRE DE LA HARPE, *Simplicity of the projective unitary groups defined by simple factors*, Comm. Math. Helv. **54** (1979), 334–345.
- [2] L. N. VASERSTEIN, *Normal subgroups of the general linear groups over Banach algebras*, J. Pure Appl. Algebra **41** (1986), 99–112.
- [3] ——, *On normal subgroups of Chevalley groups over commutative rings*, Tôhoku Math. J. **38** (1986), 219–230.
- [4] C. T. C. WALL, *On the axiomatic foundations of the theory of Hermitian forms*, Proc. Cambridge Phil. Soc. **67** (1970), 243–250.

Department of Mathematics
The Pennsylvania State University
University Park, PA 16802, USA.

Received September 10, 1986.