Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 63 (1988)

Artikel: A non-immersion theorem for spaceforms.
Autor: Pedit, F.J.
DOI: https://doi.org/10.5169/seals-48225

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Comment. Math. Helvetici 63 (1988) 672–674 0010–2571/88/040672–03$01.50 + 0.20/0
© 1988 Birkhäuser Verlag, Basel

A non-immersion theorem for spaceforms

F. J. PEDIT

Introduction

This note gives a uniform treatment of some non-existence results concerning isometric immersions of spaceforms into spaceforms. Let M, $\tilde{M}$ be spaceforms of $\dim M = m$, $\dim \tilde{M} = 2m - 1$ and curvatures c, $\tilde{c}$. Then we prove the following

THEOREM. *Let M be complete. Then there exists no isometric immersion $f: M \to \tilde{M}$ if*

(1) $c < \tilde{c}$, $c < 0$ *and M is fuchsian,*

(2) $0 < c < \tilde{c}$.

A hyperbolic spaceform is called fuchsian if the limit set of the fundamental group $\Pi_1(M)$ in the sphere at infinity of the universal cover contains more than two (and hence infinitely many) points [2]. Part (1) of our result extends recent work of F. Xavier [6] where the same statement is proven by a different method in case $\tilde{M} = \mathbb{R}^{2m-1}$ with the standard flat metric. Since every compact hyperbolic spaceform is fuchsian our result also relates to early work of S. S. Chern and N. H. Kuiper [1] where, in order to use compactness of M, $\tilde{M}$ has to be diffeomorphic to $\mathbb{R}^{2m-1}$. Behind all of this one clearly has in mind the problem of whether m-dimensional hyperbolic space H^m can be isometrically immersed into $\mathbb{R}^{2m-1}$. The case $m = 2$ was treated by D. Hilbert who showed that there are no isometric immersions of H^2 into $\mathbb{R}^3$ [4]. Part (2) proves again a result of J. D. Moore [3] and we only add it since it fits canonically into our approach.

It is well known [4] that in the case $c \geq \tilde{c}$ one always has isometric immersions $f: M \to \tilde{M}$ for complete M.

Our method is based on the observation that an isometric immersion $f: M \to \tilde{M}$ always comes with a flat metric if $\tilde{c} > c$. If M is complete this metric will be complete and hence put restrictions on the fundamental group of M.

Basic facts

Let I, $\tilde{I}$ be the riemannian metrics of constant curvatures c, $\tilde{c}$ on M, $\tilde{M}$ and assume $\kappa = \tilde{c} - c > 0$. If $f: M \to \tilde{M}$ is an isometric immersion then $f^*\tilde{I} = I$ and we

denote by $II \in \Gamma(\odot^2 T^*M, \perp_f M)$ the 2nd fundamental form of f. Applying E. Cartan's Theorem on exteriorly orthogonal symmetric bilinear forms to $II \oplus \kappa I \in \Gamma(\odot^2 T^*M, \perp_f M \oplus \mathbb{R})$ one gets [3]:

For every $p \in M$ there exists an open neighbourhood $p \in U \subset M$, an orthonormal coframe $(\omega^1, \ldots, \omega^m)$ on U and an orthonormal frame $(e_{m+1}, \ldots, e_{2m-1})$ of $\perp_f M$ over U such that

$$I = \sum_i (\omega^i)^2$$

$$II = \sum_{i,\alpha} b_i^\alpha (\omega^i)^2 e_\alpha$$

with smooth functions $b_i^\alpha : U \to \mathbb{R}$. Moreover, setting

$$a_i = \frac{1}{\sqrt{\kappa}} \sqrt{\kappa + b_i^2}, \; b_i^2 = \sum_\alpha (b_i^\alpha)^2,$$

the 1-forms $a_i \omega^i \in \Omega^1(U)$ are closed. Hence there exist (principal curvature) coordinates $x = (x^1, \ldots, x^m) : U \to \mathbb{R}^m$ with $dx^i = a_i \omega^i$.

Let $III \in \Gamma(\odot^2 T^*M, \mathbb{R})$ be the 3rd fundamental form of f, i.e.,

$$III(X, Y) = \mathrm{tr}_I \tilde{I}(II(X, —), II(—, Y)), \qquad X, Y \in TM,$$

then we have

LEMMA. *Let M be complete. Then $I_0 = \frac{1}{\kappa} III + I$ is a complete flat riemannian metric on M.*

Proof. Since $\kappa > 0$ and III is positive semi-definite $I_0 \geqslant I$ and so I_0 is complete. To show flatness of I_0 we express I_0 in the above coordinates on $U \subset M$:

$$III = \sum_{i,\alpha} (b_i^\alpha)^2 (\omega^i)^2 = \sum_i b_i^2 (\omega^i)^2$$

and so

$$I_0 = \frac{1}{\kappa} III + I = \frac{1}{\kappa} \sum_i (b_i^2 + \kappa)(\omega^i)^2 = \sum_i (a_i \omega^i)^2 = \sum_i (dx^i)^2.$$

Hence I_0 is flat. □

Proof of the theorem

Assume there is an isometric immersion $f : M \to \tilde{M}$ with M complete and $\tilde{c} - c > 0$. Then by the Lemma (M, I_0) is a flat complete riemannian manifold.

Hence its fundamental group satisfies the exact sequence

$$0 \to \mathbb{Z}^r \to \Pi_1(M) \to G \to 1$$

where $0 \leq r \leq m$ and G is finite [5].

Now let $c < 0$ and M be fuchsian. Then $\Pi_1(M)$ contains a free group on at least 2 generators $a, b \in \Pi_1(M)$ [2], [6]. Since G is finite there are $p, q \in \mathbb{N}$ with $a^p, b^q \in \mathbb{Z}^r$. But $\mathbb{Z}^r$ is abelian, so $a^p b^q = b^q a^p$ which contradicts that a, b generate a free group.

In case $c > 0$ we know that the universal cover of M is $S^m(1/c)$. But we also have that the universal cover is $\mathbb{R}^m$ which gives a contradiction. □

REFERENCES

[1] CHERN, S. S. and KUIPER, N. H., *Some theorems on the isometric embedding of compact riemannian manifolds in euclidean space,* Ann. of Math. (2), *56* (1952), 422–430.

[2] EBERLEIN, P., *Some properties of the fundamental group of a fuchsian manifold,* Inventiones Math. *19* (1973), 5–13.

[3] MOORE, J. D., *Isometric immersions of spaceforms in spaceforms,* Pacific J. Math. *40* (1972), 157–166.

[4] SPIVAC, M., *A Comprehensive Introduction to Differential Geometry,* Vol. IV, V, Publish or Perish, Inc., Berkeley, 1979.

[5] WOLF, J., *Spaces of Constant Curvature,* 3rd edition, Publish or Perish, Boston, 1974.

[6] XAVIER, F., *A non-immersion theorem for hyperbolic manifolds.* Comm. Math. Helv. *60* (1985), 280–283.

Department of Mathematics
University of North Carolina
Chapel Hill, North Carolina 27514 USA

Received November 4, 1987