

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 64 (1989)

Artikel: A linearity theorem for group actions on spheres with applications to homotopy representations.
Autor: Bauer, Stefan
DOI: <https://doi.org/10.5169/seals-48938>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A linearity theorem for group actions on spheres with applications to homotopy representations

STEFAN BAUER

Abstract. Let G be a finite group and X an equivariant $\mathbf{Z}/|G|$ -homology sphere. By Smith-theory the fixed point set X^H for a p -subgroup H is a \mathbf{Z}/p -homology sphere of dimension $d(H)$.

THEOREM. *There exists a virtual representation $V - W \in RO(G)$ such that $d(H) = \dim V^H - \dim W^H$ holds for any subgroup H of G of prime power order.*

This applies to describing the Grothendieck group $\mathcal{D}(G)$ of dimension functions of homotopy representations for compact Lie groups G in algebraic terms.

Let G be a finite group and $T \subset \mathbf{Q}$ denote a subring which is contained in the local ring $\mathbf{Z}_{(p)}$ for any prime divisor p of the order $|G|$ of G . If X is a finitistic G -space, which is a T -homology sphere, then the fixed point set X^H is a $\mathbf{Z}_{(p)}$ -homology sphere of dimension $d(H)$ for any p -subgroup H of G .

The purpose of this paper is to give a proof and an application of the following

THEOREM. *There exists a virtual representation $V - W \in RO(G)$ such that $d(H) = \dim V^H - \dim W^H$ holds for any subgroup H of G of prime power order.*

In particular d can be characterized by the combinatorial congruences in (1.1). Related linearity theorems are known for special cases: Homotopy representations of finite nilpotent groups do have stably linear dimension functions [tD1]. The theorem above is known for p -groups [Do–Ha].

Taking into account tom Dieck’s analysis of group actions on homotopy spheres (compare [tD4] and [tD5]), the linearity theorem also gives sufficient conditions for the existence of group actions on homotopy spheres. An application to homotopy representations will be discussed in the second part.

Let \mathcal{P} denote the open family of subgroups (i.e. \mathcal{P} is a union of conjugacy classes, such that for any $K \in \mathcal{P}$ all subgroups of K are also contained in \mathcal{P}) of prime power order and \mathcal{P}' the subfamily of cyclic ones.

(1.1) **DEFINITION.** Let $\mathcal{D}_{\mathcal{P}}(G)$ denote the group of functions $f: \mathcal{P} \rightarrow \mathbf{Z}$,

constant on conjugacy classes, which satisfy the following relations: For every triple $H \triangleleft K \triangleleft M$ of subgroups of G with $K/H \cong \mathbf{Z}/p$ and H a p -group,

- (1) $f(H) \equiv f(K) \pmod 2$, if p is odd or if $M/H \cong \mathbf{Z}/4$
- (2) $f(H) \equiv f(K) \pmod 4$, if $M/H \cong$ quaternion group
- (3) $f(H) + p \cdot f(M) = \sum f(K_i)$, if $M/H \cong \mathbf{Z}/p \times \mathbf{Z}/p$. Here of course the sum is over all K_i with $H \triangleleft K_i \triangleleft M$ and $K_i/H \cong \mathbf{Z}/p$
- (4) $f(H) \equiv f(K) \pmod{q^{r-l}}$, if $M/K \cong \mathbf{Z}/q^r$, acting on K/H with kernel of prime power order q^l .

(1.2) PROPOSITION. If $d: \mathcal{P} \rightarrow \mathbf{Z}$ denotes the dimension function of a finitistic G -space X which is a T -homology sphere, then $d \in \mathcal{D}_\varphi(G)$.

Proof. (1) holds by Smith-theory; for (2) compare [tD1] or [Do-Ha]. (3) is the Borel formula and (4) follows from a spectral sequence argument (the proof is almost verbatim the same as the proof of [tD2; 4.1], dealing the case $q = 2$). For the reader's convenience I will recall it here. It may be assumed $H = 1$. Look at cohomology with \mathbf{Z}/p -coefficients of the Borel-fibration

$$(X, X^K) \rightarrow (EM \times_K X, EM \times_K X^K) \rightarrow BK.$$

By Smith theory it may be assumed $d(1) = m > n = d(K)$. Fix generators $z \in \text{Coker}(H^n(X) \rightarrow H^n(X^K))$ and $u \in H^m(X)$ to obtain from the exact sequence $H^{j-1}(X^K) \xrightarrow{\delta} H^j(X, X^K) \xrightarrow{i} H^j(X)$ generators $y \in H^{n+1}(X, X^K)$ and $v \in H^m(X, X^K)$ such that $\delta z = y$ and $\iota v = u$. The spectral sequence of the Borel fibration has as E_2 -term $E_2^{ij} = H^i(BK) \otimes H^j(X, X^K)$. The transgression $d: E_2^{0,m} \rightarrow E_2^{m-n, n+1}$ gives a relation $d(1 \otimes \underset{\psi}{u}) = A \otimes y$ for a unique element $0 \neq A \in H^{m-n}(BK)$.

Consider the following map of relative fibrations:

$$\begin{array}{ccc} (X, X^K) & \xrightarrow{l_g} & (X, X^K) \\ \downarrow & & \downarrow \\ EM \times_K (X, X^K) & \xrightarrow{L_g} & EM \times_K (X, X^K) \\ \downarrow & & \downarrow \\ BK & \xrightarrow{a_g} & BK \end{array}$$

Here l_g denotes left translation by an element $g \in M$ of order q^k . L_g is given by $(e, x) \mapsto (eg^{-1}, gx)$ and a_g is the induced map on BK ; it is induced by the automorphism of K , given by conjugation by g . The induced map of spectral sequences gives a relation $A \otimes (\text{deg } l_g) \cdot y = d(1 \otimes (\text{deg } l_g) \cdot u) = d(L_g^*(1 \otimes u)) =$

$L_g^*d(1 \otimes u) = L_g^*(A \otimes y) = a_g^*A \otimes l_g^*y = a_g^*A \otimes (\deg l_g^K)y$. Therefore $a_g^*A = \varepsilon \cdot A$ with $\varepsilon = (\deg l_g)(\deg l_g^K)^{-1}$.

For a generator $t \in H^2(BK)$ the element A takes the form $a \cdot t^{(m-n)/2}$. The action of g on $H^2(BK)$ is by multiplication with $\gamma \in Z/p^*$ such that $\gamma^{q^r} = 1$ iff $(k-l)$ divides r . From $a_g^*A = \gamma^{(m-n)/2} \cdot A = \varepsilon A$ one obtains that $q^{(k-l)}$ divides $(m-n)$. ■

(1.3) THEOREM. *The dimension homomorphism $RO(G) \rightarrow \mathcal{D}_p(G)$ is surjective for any finite group G .*

Proof. For an open subfamily \mathcal{F} of \mathcal{P} let $\mathcal{D}_{\mathcal{F}}(G)$ denote the functions in $\mathcal{D}_p(G)$, restricted to \mathcal{F} . Note that restriction $\mathcal{D}_p(G) \rightarrow \mathcal{D}_{\mathcal{F}}(G)$ is injective. Hence it suffices to show that $RO(G) \rightarrow \mathcal{D}_{\mathcal{F}}(G)$ is surjective for any open subfamily \mathcal{F} of \mathcal{P} . I will do this inductively. Let $\mathcal{F}' \supset \mathcal{F}$ be adjacent families, i.e. $\mathcal{F}' \setminus \mathcal{F}$ is a conjugacy class (H) , H being a cyclic p -group. Suppose $RO(G) \rightarrow \mathcal{D}_{\mathcal{F}}(G)$ is surjective and let $f \in \ker(\mathcal{D}_p(G) \rightarrow \mathcal{D}_{\mathcal{F}}(G))$. I will show that for any prime q there exists a virtual representation $V_q \in \ker(RO(G) \rightarrow \mathcal{D}_{\mathcal{F}}(G))$ with $\dim V_q^H = n_q \cdot f(H)$ such that n_q divides $|G|$ and is prime to q . As the g.c.d. of the n_q is 1, a linear combination V of the V_q will satisfy $\dim V^H = f(H)$, thus showing $RO(G) \rightarrow \mathcal{D}_{\mathcal{F}}(G)$ is surjective.

So let's fix a prime q and choose K such that K/H is a q -Sylow subgroup of NH/H .

First case. $q = p$. Note that K is a p -group. Hence by [tD1] or [Do-Ha] there exists $\tilde{V}_p \in RO(K)$ with $\dim \tilde{V}_p^L = f(L)$ for $L \leq K$. Setting $V_p = \text{ind}_K^G \tilde{V}_p$, one has $\dim V_p^L = n_p \cdot f(L)$ for $L \in \mathcal{F}'$, where $n_p = |NH:K|$ is prime to p .

Second case. $q \neq p$. Note that K is a split extension by the Zassenhaus theorem. The homomorphism $\rho: K \rightarrow \text{Aut } H$, induced by conjugation, has cyclic image. Let $K' < K$ be the preimage of $\{\pm 1\} < \text{Aut } H$. I will construct $\tilde{U} \in RO(K')$ with $\dim \tilde{U}^L = 0$ for $L < K'$, $L \in \mathcal{F}$, and $\dim \tilde{U}^H = 2$.

Note that $\ker \rho = K''$ is nilpotent. Let W be an irreducible representation of H with $|H:\ker W| = p$ and let R denote the trivial representation. These H -representations can be viewed as K'' -representations with the q -Sylow subgroup K''_q acting trivially. In case $K'' = K'$, set $\tilde{U} = 2R - W$ (resp. $\tilde{U} = 2R - 2W$, if $p = 2$). Otherwise take $\tilde{W} = 2R - W$ and note that \tilde{W} has complex structure, thus so does $\text{ind}_{K''}^{K'} \tilde{W}$. But all representations of the dihedral group K'/K''_2 are of real type (here K''_2 denotes the 2-Sylow subgroup of K''). Hence there is $\tilde{U} \in RO(K')$ with $\mathbf{C} \otimes_{\mathbf{R}} \tilde{U} = \text{ind}_{K''}^{K'} \tilde{W}$. Now for $U = \text{ind}_K^G \tilde{U}$ one has $\dim U^L = 0$ for $L \in \mathcal{F}$ and $\dim U^H = 2 \cdot |NH:K'| = 2 \cdot |NH:K| \cdot |K:K'| = 2n_q |K:K'|$. Hence the virtual representation $(f(H)/2 |K:K'|) \cdot U$ can serve as V_q . ■

Note that $\mathcal{D}_\varphi(G)$ is a Mackey-functor, with induction map $i_H^G: \mathcal{D}_\varphi(H) \rightarrow \mathcal{D}_\varphi(G)$ defined by $i_H^G f(K) = \sum f(gKg^{-1} \cap H)$, where the sum is over double cosets $KgH \in K \backslash G/H$ (compare [tD–Pe; 9]). In the sense of Serre [Se; 12] I have just proved a “Brauer”-like statement:

(1.4) COROLLARY. *The induction map $\bigoplus_{(H)} \mathcal{D}_\varphi(H) \rightarrow \mathcal{D}_\varphi(G)$ is surjective, where the sum is taken over all $\Gamma_{\mathbb{R}}$ -elementary subgroups H of G .*

From this view G -spheres behave like real representations.

2. Applications to homotopy representations

A *homotopy representation* of a compact Lie group G is a G -CW-complex X of finite orbit type such that the fixed point sets X^H are homotopy equivalent to spheres $S^{d(H)-1}$ of the same topological dimension. The function d on the set of subgroups of G is called *dimension function*. An addition law (up to G -homotopy equivalence) is defined by taking the join, with Grothendieck group $V(G)$. A thorough discussion of homotopy representations can be found in [tD–Pe].

Let $\Psi(G)$ denote the set of conjugacy classes of closed subgroups of G , topologized via the Hausdorff-metric (compare [tD6], 5.6). Let $C(G)$ denote the group of functions $f: \Psi(G) \rightarrow \mathbb{Z}$. Associating to a homotopy representation its dimension function gives a group homomorphism $d: V(G) \rightarrow C(G)$. There are a couple of partial results on the image of d . The linearity theorem applies to give a complete description of the image of d in terms of dimension functions of linear representations as well as in combinatorial terms. The following definition introduces the candidate $\mathcal{D}(G)$ for this image, which will turn out to be the right guess.

(2.1) DEFINITION. The group $\mathcal{D}(G)$ of dimension functions of G consists of all continuous functions $f: \Psi(G) \rightarrow \mathbb{Z}$ subject to the following conditions: For any finite subquotient K/H of G there exists an element $V - W \in RO(K/H)$ such that for L with $H \triangleleft L < K$,

- (i) if L/H is a p -group, then $f(L) = \dim V^L - \dim W^L$
- (ii) if the p -Sylow subgroup $(L/H)_p$ is normal in L/H with quotient a 2-group, then $f(L) \equiv \dim V^L - \dim W^L \pmod{2}$.

Remarks. –The continuity condition corresponds to the finite orbit type condition in the definition of homotopy representations, compare [Ba; (2.4)].

– For a homotopy representation X the fixed point set X^H is a K/H -equivariant sphere. The existence of $V - W \in RO(K/H)$ satisfying (i) is a consequence of the main theorem.

- The relations in (ii) are a consequence of an Artin relation (mod 2), as given in [Do]. The Euler characteristic of X^L is determined by the dimensions $\dim X^M$ for $M < L$ and M/H a p -group, if L is as in (ii). The Lefschetz fixed point theorem gives a tight connection between this Artin relation and the orientation behaviour of group actions on homotopy spheres, compare [tD2, (1.6), (4.1) and the following remarks].
- There also is a description of $\mathcal{D}(G)$ in terms of combinatorial relations. As to (i) these are stated in (1.1). The relation corresponding to (ii) is stated in [Do].

Here are some conditions that, together, are sufficient for a function $d: \Psi(G) \rightarrow \mathbf{Z}$ to be the dimension function of a homotopy representation:

- (2.2) (a) $d \in \mathcal{D}(G)$ with $d(H) \geq 0$ for $H < G$
- (b) For any subgroup $H < G$ there is a unique maximal *formal isotropy group* $\hat{H} > H$ such that $d(\hat{H}) = d(H)$.
- (c) The set $\text{Iso}(d)$ of formal isotropy groups is closed under intersection, i.e. if $K, H \in \text{Iso}(d)$, then $H \cap K \in \text{Iso}(d)$.
- (d) If $K > H$ then $d(H) \geq d(K) + \dim(G/\hat{K})^H - \dim W\hat{K}$
- (e) There exists a G -complex A with
- (i) $\text{Iso}(d) \supset \text{Iso}(A) \supset (S_1 \cup S_2)$, where
 $S_1 = \{H \in \text{Iso}(d) \mid d(H) \leq \dim WH + 2\}$ and
 $S_2 = \{H \in \text{Iso}(d) \mid \text{there exists } K \in \text{Iso}(d), K \geq H \text{ such that } d(H) = d(K) + \dim(G/K)^H\}$
- (ii) If $K \leq H \leq G$ and $K \in \text{Iso}(A)$, then A^H is a WH -homotopy representation.

(2.3) THEOREM. *If the conditions (2.2) are satisfied, then there exists a homotopy representation X with dimension function d and $\text{Iso}(X) \subset \text{Iso}(d)$.*

Proof. By [Ba; 5.7] one only has to prove it for finite G . To do so, it suffices to show that for each $H < G$ there is a WH -representation sphere Y , with $\dim Y^{K/H} = d(K)$ for any p -group K/H in WH (compare [tD2; 1.7]). By [tD1; 3.6], Y need only exist stably, i.e. there is an element $V - W \in RO(WH)$ with $\dim V^K - \dim W^K = d(K)$ for any p -group K/H in WH . But this is true by the definition of $\mathcal{D}(G)$. Note that the orientation behaviour of X^H is determined by (2.1)(ii) (compare also [Ba; 5.4]). ■

Remark. All conditions in (2.2) are necessary ones, except condition (c) (compare [Ba] and [La]). The latter paper gives an example (2.14) in which condition (c) does not hold.

Stably the picture is more pleasant: Combining with results in [tD–Pe] and [tD3], one gets the structure theorem:

(2.4) THEOREM. *There exists an exact sequence*

$$0 \rightarrow \text{Pic}(A(G)) \rightarrow V(G) \xrightarrow{d} \mathcal{D}(G) \rightarrow 0.$$

Here $\text{Pic}(A(G))$ denotes the Picard group of the Burnside ring $A(G)$.

Proof. The kernel $v(G)$ of d was computed in [tD–Pe] for the finite group case and in [tD3] for the compact Lie case: For an appropriate G -map $f: X \rightarrow Y$ between two homotopy representations with the same dimension function the collection of degrees $(\deg f^H)$ defines an element in $\text{Pic}(A(G))$. This results in an isomorphism $v(G) \cong \text{Pic}(A(G))$. By the remarks following the definition of $\mathcal{D}(G)$ the homomorphism d maps into $\mathcal{D}(G)$. Given a dimension function $d \in \mathcal{D}(G)$, all of the conditions (2.2) can be satisfied by adding the dimension function of an appropriate real representation, compare [Ba, (2.4)]. This proves surjectivity. ■

REFERENCES

- [Ba] BAUER, S., *Dimension functions of homotopy representations for compact Lie groups* (to appear in "Mathematische Annalen").
- [tD1] TOM DIECK, T., *Homotopiedarstellungen endlicher Gruppen: Dimensionsfunktionen*, Invent. Math. 67, 231–252, (1982).
- [tD2] TOM DIECK, T., *Dimension functions of homotopy representations*, Mathemata Gottingensis, Heft 31 (1985).
- [tD3] TOM DIECK, T., *Die Picard-Gruppe des Burnside-Ringes einer kompakten Lieschen Gruppe I*. Mathemata Gottingensis, Heft 46 (1986).
- [tD4] TOM DIECK, T., *The singular set of group actions on homotopy spheres*. Arch. Math. 43, 551–558 (1984).
- [tD5] TOM DIECK, T., *The homotopy type of group actions on homotopy spheres*. Arch. Math. 45, 174–179 (1985).
- [tD6] TOM DIECK, T., *Transformation Groups and Representation Theory*. LNM 766, Springer 1979.
- [tD–Pe] TOM DIECK, T. and PETRIE, T., *Homotopy representations of finite groups*. Publ. Math. I.H.E.S. 56, 129–169 (1983).
- [Do] DOTZEL, R. M., *An Artin relation (mod 2) for finite group actions on spheres*, Pacif J. Math. 114, 335–343 (1984).
- [Do–Ha] DOTZEL, R. M. and HAMRICK, G., *p-group actions on homology spheres*. Invent. Math. 62, 437–442 (1981).
- [La] LAITINEN, E., *Unstable homotopy theory of homotopy representations*, in: LNM 1217, 210–248 (1987).
- [Se] SERRE, J.-P., *Linear Representations of Finite Groups*. GTM 42, Springer (1977).

SONDERFORSCHUNGSBEREICH 170: "Geometrie und Analysis" Mathematisches Institut, Bunsenstr, 3–5, D-3400 Göttingen

University of Chicago, Department of Mathematics, 5734 University Avenue, Chicago, IL 60637

Received March 10, 1987.